



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 18, 2026 – 04:14 AM UTC

PDB ID : 5OV7 / pdb_00005ov7
Title : tubulin - rigosertib complex
Authors : Menchon, G.; Protá, A.E.; Steinmetz, M.; Jost, M.
Deposited on : 2017-08-28
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	451	2% 74% 22% ••
1	C	451	% 76% 21% •
2	B	445	% 70% 24% 6%
2	D	445	2% 69% 26% 5%
3	E	143	3% 66% 19% • 14%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	F	384	5% 60% 29% 10%

2 Entry composition

There are 13 unique types of molecules in this entry. The entry contains 18026 atoms, of which 48 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tubulin alpha-1B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	437	Total	C	N	O	S	0	2	0
			3428	2171	584	651	22			
1	C	440	Total	C	N	O	S	0	2	0
			3449	2183	585	659	22			

- Molecule 2 is a protein called Tubulin beta-2B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	420	Total	C	N	O	S	0	4	0
			3325	2092	565	640	28			
2	D	422	Total	C	N	O	S	0	0	0
			3314	2083	563	641	27			

- Molecule 3 is a protein called Stathmin-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	123	Total	C	N	O	S	0	1	0
			1021	630	185	201	5			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	3	MET	-	initiating methionine	UNP P63043
E	4	ALA	-	expression tag	UNP P63043


- Molecule 4 is a protein called Uncharacterized protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	F	347	Total	C	N	O	S	0	0	0
			2830	1811	486	519	14			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	379	HIS	-	expression tag	UNP E1BQ43
F	380	HIS	-	expression tag	UNP E1BQ43
F	381	HIS	-	expression tag	UNP E1BQ43
F	382	HIS	-	expression tag	UNP E1BQ43
F	383	HIS	-	expression tag	UNP E1BQ43
F	384	HIS	-	expression tag	UNP E1BQ43

- Molecule 5 is GUANOSINE-5'-TRIPHOSPHATE (CCD ID: GTP) (formula: $C_{10}H_{16}N_5O_{14}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
5	C	1	Total	C	N	O	P	0	0
			32	10	5	14	3		

- Molecule 6 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Mg	0	0
			1	1		
6	B	1	Total	Mg	0	0
			1	1		
6	C	1	Total	Mg	0	0
			1	1		
6	D	1	Total	Mg	0	0
			1	1		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	F	1	Total	Mg	0	0
			1	1		

- Molecule 7 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Ca	0	0
			1	1		

- Molecule 8 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			6	3	3		
8	A	1	Total	C	O	0	0
			6	3	3		
8	B	1	Total	C	O	0	0
			6	3	3		
8	B	1	Total	C	O	0	0
			6	3	3		
8	C	1	Total	C	O	0	0
			6	3	3		
8	C	1	Total	C	O	0	0
			6	3	3		

- Molecule 9 is GUANOSINE-5'-DIPHOSPHATE (CCD ID: GDP) (formula: C₁₀H₁₅N₅O₁₁P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
9	B	1	Total 28	C 10	N 5	O 11	P 2	0	0
9	D	1	Total 28	C 10	N 5	O 11	P 2	0	0

- Molecule 10 is N-[2-methoxy-5-({[(E)-2-(2,4,6-trimethoxyphenyl)ethenyl]sulfonyl}methyl)phenyl]glycine (CCD ID: 6FS) (formula: C₂₁H₂₅NO₈S).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
10	B	1	Total	C	H	N	O	S	0	0
			55	21	24	1	8	1		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
10	D	1	Total	C	H	N	O	S	
			55	21	24	1	8	1	
								0	0

- Molecule 11 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
11	B	1	Total	C	N	O	S		
			12	6	1	4	1		
								0	0

- Molecule 12 is PHOSPHOMETHYLPHOSPHONIC ACID ADENYLATE ESTER (CCD ID: ACP) (formula: C₁₁H₁₈N₅O₁₂P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
12	F	1	Total 31	C 11	N 5	O 12	P 3	0	0


- Molecule 13 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
13	A	50	Total O 50 50	0	0
13	B	88	Total O 88 88	0	0
13	C	133	Total O 133 133	0	0
13	D	43	Total O 43 43	0	0
13	E	9	Total O 9 9	0	0
13	F	21	Total O 21 21	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tubulin alpha-1B chain




- Molecule 1: Tubulin alpha-1B chain




- Molecule 2: Tubulin beta-2B chain







• Molecule 2: Tubulin beta-2B chain




• Molecule 3: Stathmin-4



• Molecule 4: Uncharacterized protein





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	104.73Å 156.76Å 182.63Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.08 – 2.40 48.08 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.0 (48.08-2.40) 99.0 (48.08-2.40)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.10 (at 2.39Å)	Xtriage
Refinement program	PHENIX (dev_2863: ???)	Depositor
R, R_{free}	0.190 , 0.235 0.193 , 0.237	Depositor DCC
R_{free} test set	5844 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	54.3	Xtriage
Anisotropy	0.169	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 54.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	18026	wwPDB-VP
Average B, all atoms (Å ²)	77.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACP, GDP, MG, MES, CA, GTP, 6FS, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.21	0/3512	0.45	3/4767 (0.1%)
1	C	0.30	1/3533 (0.0%)	0.46	0/4797
2	B	0.30	0/3410	0.44	1/4616 (0.0%)
2	D	0.22	0/3387	0.40	0/4588
3	E	0.15	0/1033	0.34	0/1371
4	F	0.11	0/2893	0.33	0/3906
All	All	0.24	1/17768 (0.0%)	0.42	4/24045 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	6	SER	C-O	-5.66	1.17	1.23

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	160	ASP	N-CA-C	6.68	118.65	111.36
1	A	164	LYS	N-CA-C	6.29	119.37	110.50
2	B	97	SER	N-CA-C	5.95	117.77	111.28
1	A	133	GLN	N-CA-C	5.07	116.89	111.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3428	0	3350	85	0
1	C	3449	0	3363	69	0
2	B	3325	0	3222	96	0
2	D	3314	0	3194	105	0
3	E	1021	0	1036	28	0
4	F	2830	0	2800	107	0
5	A	32	0	12	0	0
5	C	32	0	12	1	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
6	C	1	0	0	0	0
6	D	1	0	0	0	0
6	F	1	0	0	0	0
7	A	1	0	0	0	0
8	A	12	0	16	2	0
8	B	12	0	16	1	0
8	C	12	0	16	2	0
9	B	28	0	12	1	0
9	D	28	0	12	3	0
10	B	31	24	0	8	0
10	D	31	24	0	4	0
11	B	12	0	12	1	0
12	F	31	0	14	2	0
13	A	50	0	0	5	0
13	B	88	0	0	5	0
13	C	133	0	0	7	0
13	D	43	0	0	2	1
13	E	9	0	0	0	0
13	F	21	0	0	2	1
All	All	17978	48	17087	470	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 470 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:248:LEU:HD21	2:D:352:LYS:CE	1.74	1.16
4:F:87:LEU:O	4:F:87:LEU:HD12	1.45	1.16
2:B:352:LYS:HE2	10:B:504:6FS:C20	1.77	1.14

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:F:81:ILE:HA	4:F:87:LEU:HD11	1.27	1.12
4:F:320:MET:HG2	4:F:330:ILE:HD11	1.35	1.02

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
13:D:634:HOH:O	13:F:520:HOH:O[3_545]	2.12	0.08

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	437/451 (97%)	427 (98%)	10 (2%)	0	100	100
1	C	440/451 (98%)	428 (97%)	12 (3%)	0	100	100
2	B	420/445 (94%)	411 (98%)	9 (2%)	0	100	100
2	D	418/445 (94%)	408 (98%)	10 (2%)	0	100	100
3	E	120/143 (84%)	117 (98%)	3 (2%)	0	100	100
4	F	341/384 (89%)	328 (96%)	12 (4%)	1 (0%)	36	50
All	All	2176/2319 (94%)	2119 (97%)	56 (3%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	F	235	ASP

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	370/379 (98%)	368 (100%)	2 (0%)	81	91
1	C	373/379 (98%)	368 (99%)	5 (1%)	61	80
2	B	368/383 (96%)	365 (99%)	3 (1%)	73	86
2	D	364/383 (95%)	360 (99%)	4 (1%)	65	82
3	E	111/127 (87%)	107 (96%)	4 (4%)	31	52
4	F	309/342 (90%)	304 (98%)	5 (2%)	55	76
All	All	1895/1993 (95%)	1872 (99%)	23 (1%)	63	81

5 of 23 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	E	15	THR
3	E	133	VAL
3	E	74	GLU
4	F	12	SER
1	C	71	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	91	ASN
3	E	129	HIS
4	F	306	HIS
4	F	38	ASN
3	E	51	GLN

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 6 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
8	GOL	A	505	-	5,5,5	0.91	0	5,5,5	1.11	0
8	GOL	C	501	-	5,5,5	0.92	0	5,5,5	1.09	0
11	MES	B	505	-	12,12,12	2.24	1 (8%)	15,16,16	2.23	7 (46%)
5	GTP	A	501	6	33,34,34	1.05	4 (12%)	50,54,54	1.54	9 (18%)
8	GOL	B	506	-	5,5,5	0.91	0	5,5,5	1.09	0
10	6FS	D	503	-	31,32,32	1.05	3 (9%)	40,44,44	1.74	7 (17%)
12	ACP	F	401	6	31,33,33	1.56	7 (22%)	47,52,52	1.96	11 (23%)
8	GOL	C	504	-	5,5,5	0.93	0	5,5,5	1.09	0
10	6FS	B	504	-	31,32,32	1.15	3 (9%)	40,44,44	1.69	8 (20%)
9	GDP	B	501	6	29,30,30	1.64	7 (24%)	45,47,47	1.86	9 (20%)
5	GTP	C	502	6	33,34,34	0.97	2 (6%)	50,54,54	1.56	9 (18%)
8	GOL	B	503	-	5,5,5	0.16	0	5,5,5	0.45	0
9	GDP	D	501	6	29,30,30	1.16	2 (6%)	45,47,47	1.75	6 (13%)
8	GOL	A	504	-	5,5,5	0.92	0	5,5,5	1.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	GOL	A	505	-	-	0/4/4/4	-
8	GOL	C	501	-	-	0/4/4/4	-
11	MES	B	505	-	-	5/6/14/14	0/1/1/1
5	GTP	A	501	6	-	8/22/38/38	0/3/3/3
8	GOL	B	506	-	-	0/4/4/4	-
10	6FS	D	503	-	-	8/24/24/24	0/2/2/2
12	ACP	F	401	6	-	8/19/38/38	0/3/3/3
8	GOL	C	504	-	-	0/4/4/4	-
10	6FS	B	504	-	-	16/24/24/24	0/2/2/2
9	GDP	B	501	6	-	4/16/32/32	0/3/3/3
5	GTP	C	502	6	-	9/22/38/38	0/3/3/3
8	GOL	B	503	-	-	0/4/4/4	-
9	GDP	D	501	6	-	5/16/32/32	0/3/3/3
8	GOL	A	504	-	-	0/4/4/4	-

The worst 5 of 29 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
11	B	505	MES	C8-S	-7.47	1.67	1.77
9	B	501	GDP	C6-N1	-3.82	1.31	1.38
12	F	401	ACP	C5-N7	-3.31	1.33	1.39
10	B	504	6FS	O4-S1	3.20	1.46	1.44
12	F	401	ACP	C5-C4	3.12	1.44	1.39

The worst 5 of 66 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	B	501	GDP	C5-C4-N3	-6.32	118.33	128.39
10	D	503	6FS	C20-N1-C16	-6.07	114.96	123.94
12	F	401	ACP	C5-C4-N3	-6.04	118.40	126.72
9	D	501	GDP	C5-C4-N3	-6.03	118.79	128.39
9	B	501	GDP	C2-N3-C4	5.37	121.55	112.30

There are no chirality outliers.

5 of 63 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	501	GTP	PB-O3B-PG-O2G
5	A	501	GTP	C5'-O5'-PA-O3A
5	A	501	GTP	C5'-O5'-PA-O1A
5	A	501	GTP	C5'-O5'-PA-O2A

Continued on next page...

Continued from previous page...



Mol	Chain	Res	Type	Atoms
5	C	502	GTP	C5'-O5'-PA-O3A

There are no ring outliers.


10 monomers are involved in 25 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
11	B	505	MES	1	0
10	D	503	6FS	4	0
12	F	401	ACP	2	0
8	C	504	GOL	2	0
10	B	504	6FS	8	0
9	B	501	GDP	1	0
5	C	502	GTP	1	0
8	B	503	GOL	1	0
9	D	501	GDP	3	0
8	A	504	GOL	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.






Ligand ACP F 401



Ligand 6FS B 504







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	437/451 (96%)	0.04	7 (1%) 70 66	41, 68, 110, 171	2 (0%)
1	C	440/451 (97%)	-0.26	3 (0%) 84 81	28, 50, 86, 121	2 (0%)
2	B	420/445 (94%)	-0.05	4 (0%) 79 76	30, 59, 101, 158	5 (1%)
2	D	422/445 (94%)	0.29	9 (2%) 63 59	44, 87, 129, 168	6 (1%)
3	E	123/143 (86%)	0.32	5 (4%) 41 37	37, 84, 138, 180	1 (0%)
4	F	347/384 (90%)	0.50	18 (5%) 33 29	54, 103, 173, 225	0
All	All	2189/2319 (94%)	0.10	46 (2%) 63 59	28, 71, 136, 225	16 (0%)

The worst 5 of 46 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	285	ALA	4.6
2	B	371	LEU	4.3
4	F	155	ALA	4.0
2	D	248	LEU	4.0
1	A	437	VAL	4.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	MG	F	402	1/1	0.71	0.20	155,155,155,155	0
8	GOL	B	506	6/6	0.79	0.15	95,105,107,112	0
8	GOL	C	504	6/6	0.82	0.20	94,96,99,103	0
8	GOL	C	501	6/6	0.84	0.21	70,80,81,82	0
8	GOL	A	505	6/6	0.85	0.15	90,97,105,114	0
12	ACP	F	401	31/31	0.88	0.11	82,104,152,159	0
8	GOL	A	504	6/6	0.90	0.12	81,83,85,86	0
10	6FS	D	503	31/31	0.91	0.14	64,106,140,143	0
9	GDP	D	501	28/28	0.92	0.11	62,77,96,111	0
6	MG	D	502	1/1	0.94	0.06	66,66,66,66	0
10	6FS	B	504	31/31	0.94	0.12	54,77,118,134	0
8	GOL	B	503	6/6	0.95	0.14	81,87,91,92	0
11	MES	B	505	12/12	0.95	0.10	49,70,89,102	0
7	CA	A	503	1/1	0.95	0.06	91,91,91,91	0
9	GDP	B	501	28/28	0.97	0.08	33,47,62,100	0
5	GTP	A	501	32/32	0.98	0.06	34,49,58,65	0
5	GTP	C	502	32/32	0.98	0.06	31,42,48,59	0
6	MG	B	502	1/1	0.98	0.05	49,49,49,49	0
6	MG	C	503	1/1	0.99	0.02	42,42,42,42	0
6	MG	A	502	1/1	1.00	0.02	47,47,47,47	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ACP F 401:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 6FS D 503:**

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GDP D 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 6FS B 504:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GDP B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GTP A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around GTP C 502:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.