



Full wwPDB NMR Structure Validation Report ⓘ

Mar 10, 2026 – 08:49 AM UTC

PDB ID : 5OVM / pdb_00005ovm
BMRB ID : 34175
Title : Solution structure of lipase binding domain LID1 of foldase from *Pseudomonas aeruginosa*
Authors : Viegas, A.; Jaeger, K.-E.; Etzkorn, M.; Gohlke, H.; Verma, N.; Dollinger, P.; Kovacic, F.
Deposited on : 2017-08-29

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 82%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	89	 85% . . 8%

2 Ensemble composition and analysis

This entry contains 20 models. Model 9 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:67-A:146 (80)	1.90	9

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 19, 20
2	4, 18

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1291 atoms, of which 641 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Lipase chaperone.

Mol	Chain	Residues	Atoms						Trace
1	A	82	Total	C	H	N	O	S	0
			1291	411	641	112	126	1	

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	58	MET	-	initiating methionine	UNP Q01725
A	59	GLY	-	expression tag	UNP Q01725
A	60	HIS	-	expression tag	UNP Q01725
A	61	HIS	-	expression tag	UNP Q01725
A	62	HIS	-	expression tag	UNP Q01725
A	63	HIS	-	expression tag	UNP Q01725
A	64	HIS	-	expression tag	UNP Q01725
A	65	HIS	-	expression tag	UNP Q01725

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Lipase chaperone




4.2 Scores per residue for each member of the ensemble

Colouring as in section [4.1](#) above.


4.2.1 Score per residue for model 1

- Molecule 1: Lipase chaperone




4.2.2 Score per residue for model 2

- Molecule 1: Lipase chaperone




4.2.3 Score per residue for model 3

- Molecule 1: Lipase chaperone




4.2.4 Score per residue for model 4

- Molecule 1: Lipase chaperone




4.2.5 Score per residue for model 5

- Molecule 1: Lipase chaperone




4.2.6 Score per residue for model 6

- Molecule 1: Lipase chaperone




4.2.7 Score per residue for model 7

- Molecule 1: Lipase chaperone




4.2.8 Score per residue for model 8

- Molecule 1: Lipase chaperone




4.2.9 Score per residue for model 9 (medoid)

- Molecule 1: Lipase chaperone




4.2.10 Score per residue for model 10

- Molecule 1: Lipase chaperone




4.2.11 Score per residue for model 11

- Molecule 1: Lipase chaperone




4.2.12 Score per residue for model 12

- Molecule 1: Lipase chaperone




4.2.13 Score per residue for model 13

- Molecule 1: Lipase chaperone




4.2.14 Score per residue for model 14

- Molecule 1: Lipase chaperone




4.2.15 Score per residue for model 15

- Molecule 1: Lipase chaperone




4.2.16 Score per residue for model 16

- Molecule 1: Lipase chaperone




4.2.17 Score per residue for model 17

- Molecule 1: Lipase chaperone




4.2.18 Score per residue for model 18

- Molecule 1: Lipase chaperone




4.2.19 Score per residue for model 19

- Molecule 1: Lipase chaperone



4.2.20 Score per residue for model 20

- Molecule 1: Lipase chaperone



5 Refinement protocol and experimental data overview

The models were refined using the following method: *molecular dynamics*.

Of the 20 calculated structures, 20 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure calculation	
Amber	geometry optimization	
Amber	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	929
Number of shifts mapped to atoms	929
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	82%

6 Model quality [i](#)

6.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.88±0.03	0±0/642 (0.0± 0.1%)	1.36±0.03	1±1/866 (0.1± 0.1%)
All	All	0.88	2/12840 (0.0%)	1.36	21/17320 (0.1%)

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	112	SER	C-N	6.29	1.42	1.33	18	1
1	A	118	ALA	C-N	5.70	1.41	1.33	18	1

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	98	ASP	CA-CB-CG	8.38	120.98	112.60	20	2
1	A	92	ASP	CA-CB-CG	7.18	119.78	112.60	20	1
1	A	86	ASN	CA-CB-CG	6.84	119.44	112.60	18	6
1	A	100	PHE	CA-CB-CG	6.00	119.80	113.80	8	2
1	A	138	GLN	OE1-CD-NE2	-5.79	116.81	122.60	11	2
1	A	76	ASP	CA-CB-CG	5.71	118.31	112.60	4	1
1	A	111	GLN	OE1-CD-NE2	-5.65	116.95	122.60	5	2
1	A	86	ASN	OD1-CG-ND2	-5.23	117.37	122.60	9	1
1	A	137	GLN	OE1-CD-NE2	-5.22	117.38	122.60	16	3
1	A	110	GLN	OE1-CD-NE2	-5.13	117.47	122.60	8	1

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	632	623	623	0±1
All	All	12640	12460	12460	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:114:ASP:OD1	1:A:117:ARG:NE	0.59	2.32	12	1
1:A:79:PHE:CE1	1:A:96:LEU:HD13	0.48	2.43	10	1
1:A:93:ILE:HA	1:A:96:LEU:HD12	0.47	1.87	10	1
1:A:93:ILE:HG22	1:A:97:PHE:CE1	0.46	2.45	13	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	79/89 (89%)	72±2 (91±3%)	6±2 (8±2%)	1±1 (2±1%)	11	57
All	All	1580/1780 (89%)	1433 (91%)	123 (8%)	24 (2%)	11	57

All 15 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	78	SER	3
1	A	89	ILE	3
1	A	70	PHE	2
1	A	75	VAL	2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	106	GLU	2
1	A	104	VAL	2
1	A	105	GLY	2
1	A	72	GLY	1
1	A	144	LYS	1
1	A	76	ASP	1
1	A	103	ALA	1
1	A	68	THR	1
1	A	77	GLY	1
1	A	80	SER	1
1	A	67	PRO	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	67/75 (89%)	58±12 (86±17%)	6±2 (9±2%)	11	56
All	All	1283/1500 (86%)	1158 (90%)	125 (10%)	10	55

All 35 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	89	ILE	19
1	A	137	GLN	18
1	A	138	GLN	10
1	A	142	TYR	8
1	A	111	GLN	7
1	A	110	GLN	7
1	A	116	LEU	6
1	A	143	LYS	5
1	A	86	ASN	4
1	A	114	ASP	4
1	A	87	LEU	4
1	A	144	LYS	2
1	A	113	LEU	2
1	A	117	ARG	2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	101	LEU	2
1	A	98	ASP	2
1	A	107	GLU	2
1	A	92	ASP	2
1	A	145	GLU	2
1	A	71	ARG	2
1	A	94	ARG	1
1	A	106	GLU	1
1	A	115	ARG	1
1	A	90	THR	1
1	A	79	PHE	1
1	A	88	LEU	1
1	A	81	VAL	1
1	A	78	SER	1
1	A	102	SER	1
1	A	73	THR	1
1	A	91	ARG	1
1	A	96	LEU	1
1	A	129	ARG	1
1	A	109	LEU	1
1	A	136	MET	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 82% for the well-defined parts and 82% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: MD1_3.1_6.str

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	929
Number of shifts mapped to atoms	929
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	76	-0.34 ± 0.13	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	72	0.21 ± 0.15	None needed (< 0.5 ppm)
$^{13}\text{C}'$	67	-0.31 ± 0.33	None needed (< 0.5 ppm)
^{15}N	60	0.24 ± 0.24	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 82%, i.e. 904 atoms were assigned a chemical shift out of a possible 1099. 0 out of 14 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	340/399 (85%)	142/162 (88%)	139/160 (87%)	59/77 (77%)
Sidechain	531/624 (85%)	357/405 (88%)	166/191 (87%)	8/28 (29%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	33/76 (43%)	33/36 (92%)	0/40 (0%)	0/0 (—%)
Overall	904/1099 (82%)	532/603 (88%)	305/391 (78%)	67/105 (64%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 82%, i.e. 929 atoms were assigned a chemical shift out of a possible 1132. 0 out of 15 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	349/409 (85%)	146/166 (88%)	143/164 (87%)	60/79 (76%)
Sidechain	547/640 (85%)	368/416 (88%)	171/196 (87%)	8/28 (29%)
Aromatic	33/83 (40%)	33/40 (82%)	0/42 (0%)	0/1 (0%)
Overall	929/1132 (82%)	547/622 (88%)	314/402 (78%)	68/108 (63%)

7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:

