



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 03:27 PM UTC

PDB ID : 7OV7 / pdb_00007ov7
Title : The hybrid cage formed between Pizza6-S and Cu(II)-substituted trilacunary Keggin
Authors : Vandebroek, L.; Noguchi, H.; Anyushin, A.; Van Meervelt, L.; Voet, A.R.D.; Parac-Vogt, T.N.
Deposited on : 2021-06-14
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	256	12% 83% 14% .
1	B	256	32% 84% 13% .

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3987 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Pizza6-S.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	250	Total	C	N	O	0	0	0
			1770	1092	315	363			
1	B	249	Total	C	N	O	0	1	0
			1764	1090	313	361			

- Molecule 2 is COPPER (II) ION (CCD ID: CU) (formula: Cu) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	6	Total	Cu	0	0
			6	6		

- Molecule 3 is Trilacunary Keggin (CCD ID: PW9) (formula: O₃₄PW₉) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	O	P	W	0	1
			88	68	2	18		
3	B	1	Total	O	P	W	0	1
			88	68	2	18		

- Molecule 4 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	6	Total	K	0	0
			6	6		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	165	Total	O	0	0
			165	165		
5	B	100	Total	O	0	0
			100	100		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Pizza6-S



• Molecule 1: Pizza6-S



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	65.79Å 74.15Å 79.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.21 – 1.80 49.21 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.8 (49.21-1.80) 94.7 (49.21-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.29 (at 1.79Å)	Xtriage
Refinement program	PHENIX 1.15.2_3472	Depositor
R, R_{free}	0.232 , 0.287 0.244 , 0.296	Depositor DCC
R_{free} test set	1824 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	23.0	Xtriage
Anisotropy	0.531	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 41.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3987	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CU, K, PW9

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.09	0/1799	0.26	0/2477
1	B	0.08	0/1793	0.25	0/2469
All	All	0.09	0/3592	0.26	0/4946

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1770	0	1748	26	0
1	B	1764	0	1739	23	0
2	A	6	0	0	0	0
3	A	88	0	0	12	0
3	B	88	0	0	12	0
4	A	6	0	0	0	0
5	A	165	0	0	0	0
5	B	100	0	0	1	0
All	All	3987	0	3487	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:134:PRO:HG2	1:A:165:LEU:HD21	1.83	0.60
1:B:162:VAL:HG21	1:B:196:VAL:HG21	1.85	0.58
1:A:18:VAL:HB	1:A:239:THR:HG23	1.87	0.57
1:A:29:THR:HG23	1:A:60:VAL:HB	1.86	0.55
1:A:155:THR:HG23	1:A:186:VAL:HB	1.88	0.55
1:A:113:THR:HG23	1:A:144:VAL:HB	1.89	0.54
1:B:18:VAL:HB	1:B:239:THR:HG23	1.89	0.54
1:B:55:ASN:ND2	5:B:504:HOH:O	2.40	0.54
1:A:28:VAL:HG21	1:A:246:VAL:HG21	1.90	0.54
1:A:120:VAL:HG21	1:A:154:VAL:HG21	1.90	0.53
1:B:166:ALA:HB3	1:B:169:SER:HB2	1.89	0.53
1:A:50:PRO:HG2	1:A:81:LEU:HD21	1.91	0.52
3:B:401[A]:PW9:O18	3:B:401[A]:PW9:O4	2.29	0.51
3:B:401[B]:PW9:O18	3:B:401[B]:PW9:O4	2.29	0.51
3:A:307[B]:PW9:O4	3:A:307[B]:PW9:O18	2.30	0.50
3:A:307[A]:PW9:O18	3:A:307[A]:PW9:O4	2.29	0.49
3:B:401[B]:PW9:O19E	3:B:401[B]:PW9:O9E	2.31	0.49
3:A:307[A]:PW9:O9E	3:A:307[A]:PW9:O19E	2.29	0.49
3:A:307[B]:PW9:O19E	3:A:307[B]:PW9:O9E	2.30	0.49
3:B:401[A]:PW9:O9E	3:B:401[A]:PW9:O19E	2.31	0.49
1:B:155:THR:HG23	1:B:186:VAL:HB	1.93	0.49
1:A:162:VAL:HG21	1:A:196:VAL:HG21	1.96	0.48
1:A:223:ASN:ND2	1:B:181:ASN:HD21	2.12	0.48
1:A:8:PRO:HG2	1:A:39:LEU:HD21	1.96	0.48
3:A:307[B]:PW9:O16	3:A:307[B]:PW9:O21	2.33	0.47
3:B:401[A]:PW9:O16	3:B:401[A]:PW9:O21	2.32	0.46
3:B:401[A]:PW9:O22	3:B:401[A]:PW9:O17	2.34	0.46
3:B:401[B]:PW9:O17	3:B:401[B]:PW9:O22	2.34	0.46
3:A:307[B]:PW9:O4	3:A:307[B]:PW9:O20	2.34	0.46
3:B:401[B]:PW9:O16	3:B:401[B]:PW9:O21	2.33	0.46
3:A:307[B]:PW9:O17	3:A:307[B]:PW9:O22	2.34	0.46
1:A:14:THR:HG23	1:A:241:HIS:CD2	2.51	0.46
1:A:197:THR:HG23	1:A:228:VAL:HB	1.97	0.46
3:A:307[A]:PW9:O16	3:A:307[A]:PW9:O21	2.33	0.46
3:A:307[B]:PW9:O22	3:A:307[B]:PW9:O23	2.34	0.46
3:A:307[A]:PW9:O22	3:A:307[A]:PW9:O23	2.34	0.45
3:B:401[B]:PW9:O22	3:B:401[B]:PW9:O23	2.34	0.45
1:A:241:HIS:CD2	1:A:241:HIS:C	2.93	0.45
3:B:401[A]:PW9:O22	3:B:401[A]:PW9:O23	2.34	0.45
3:A:307[A]:PW9:O22	3:A:307[A]:PW9:O17	2.34	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:307[A]:PW9:O4	3:A:307[A]:PW9:O20	2.35	0.45
1:B:185:GLY:H	1:B:197:THR:HG1	1.59	0.45
1:A:115:HIS:CD2	1:A:115:HIS:C	2.94	0.45
1:B:197:THR:HG23	1:B:228:VAL:HB	1.99	0.44
1:B:199:HIS:CE1	1:B:224:THR:HB	2.52	0.44
1:B:157:HIS:CE1	1:B:182:THR:HB	2.52	0.44
1:A:73:HIS:CD2	1:A:73:HIS:C	2.95	0.44
3:B:401[A]:PW9:O4	3:B:401[A]:PW9:O20	2.35	0.44
3:B:401[B]:PW9:O4	3:B:401[B]:PW9:O20	2.35	0.44
1:A:98:THR:O	1:A:115:HIS:HB3	2.18	0.44
1:A:36:VAL:HG21	1:A:70:VAL:HG21	2.00	0.44
1:A:204:VAL:HG21	1:A:238:VAL:HG21	1.99	0.44
1:A:223:ASN:HD21	1:B:181:ASN:HD21	1.66	0.43
1:B:14:THR:HB	1:B:241:HIS:CE1	2.53	0.43
1:B:56:THR:O	1:B:73:HIS:HB3	2.18	0.43
1:B:14:THR:O	1:B:31:HIS:HB3	2.19	0.43
1:A:31:HIS:CD2	1:A:31:HIS:C	2.97	0.43
1:B:140:THR:O	1:B:157:HIS:HB3	2.19	0.42
1:B:134:PRO:HG2	1:B:165:LEU:HD21	2.00	0.42
1:A:140:THR:O	1:A:157:HIS:HB3	2.20	0.42
1:B:85:SER:OG	1:B:86:ASN:N	2.52	0.42
1:B:98:THR:O	1:B:115:HIS:HB3	2.20	0.42
1:A:79:VAL:HG12	1:A:90:VAL:HG22	2.02	0.41
1:A:157:HIS:CD2	1:A:157:HIS:C	2.98	0.41
1:B:59:GLY:H	1:B:71:THR:HB	1.85	0.41
1:B:73:HIS:CE1	1:B:98:THR:HB	2.55	0.41
1:A:14:THR:O	1:A:31:HIS:HB3	2.20	0.41
1:B:80:LYS:HE3	1:B:91:LEU:HD21	2.02	0.40
1:A:197:THR:HG22	1:A:204:VAL:HG22	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/256 (97%)	241 (97%)	7 (3%)	0	100	100
1	B	248/256 (97%)	242 (98%)	6 (2%)	0	100	100
All	All	496/512 (97%)	483 (97%)	13 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	197/201 (98%)	197 (100%)	0	100	100
1	B	195/201 (97%)	195 (100%)	0	100	100
All	All	392/402 (98%)	392 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	44	ASN
1	B	181	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 16 ligands modelled in this entry, 12 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PW9	A	307[B]	4	12,58,58	3.45	8 (66%)	3,171,171	4.02	1 (33%)
3	PW9	A	307[A]	4,2	12,58,58	3.47	8 (66%)	3,171,171	4.03	1 (33%)
3	PW9	B	401[A]	5	12,58,58	3.46	8 (66%)	3,171,171	4.11	1 (33%)
3	PW9	B	401[B]	5	12,58,58	3.50	8 (66%)	3,171,171	4.09	1 (33%)

All (32) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	401[B]	PW9	P1-O21	8.80	1.69	1.53
3	A	307[A]	PW9	P1-O21	8.71	1.69	1.53
3	B	401[A]	PW9	P1-O21	8.71	1.69	1.53
3	A	307[B]	PW9	P1-O21	8.70	1.69	1.53
3	A	307[A]	PW9	P1-O19E	-4.34	1.46	1.53
3	B	401[B]	PW9	P1-O19E	-4.30	1.46	1.53
3	B	401[A]	PW9	P1-O19E	-4.25	1.46	1.53
3	A	307[B]	PW9	P1-O19E	-4.24	1.46	1.53
3	B	401[B]	PW9	W3E-O19E	3.53	2.63	2.18
3	B	401[A]	PW9	W3E-O19E	3.47	2.63	2.18
3	A	307[A]	PW9	W3E-O19E	3.40	2.62	2.18
3	A	307[B]	PW9	W3E-O19E	3.31	2.61	2.18
3	B	401[B]	PW9	P1-O4	-3.17	1.48	1.53
3	A	307[B]	PW9	P1-O4	-3.17	1.48	1.53
3	B	401[B]	PW9	W5E-O4	3.10	2.58	2.18
3	A	307[A]	PW9	P1-O4	-3.10	1.48	1.53
3	B	401[A]	PW9	P1-O4	-3.07	1.48	1.53
3	A	307[A]	PW9	W5E-O4	3.04	2.57	2.18
3	B	401[A]	PW9	W5E-O4	3.02	2.57	2.18
3	A	307[B]	PW9	W5E-O4	2.92	2.56	2.18
3	A	307[B]	PW9	W6-O5	-2.41	1.59	2.16
3	A	307[A]	PW9	W6-O5	-2.41	1.59	2.16

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	401[A]	PW9	W6-O5	-2.41	1.59	2.16
3	B	401[B]	PW9	W6-O5	-2.40	1.59	2.16
3	A	307[A]	PW9	W3-O11	-2.16	1.65	2.16
3	A	307[B]	PW9	W3-O11	-2.15	1.65	2.16
3	B	401[A]	PW9	W3-O11	-2.15	1.65	2.16
3	B	401[B]	PW9	W3-O11	-2.14	1.65	2.16
3	A	307[A]	PW9	W5-O13	-2.04	1.68	2.16
3	A	307[B]	PW9	W5-O13	-2.04	1.68	2.16
3	B	401[B]	PW9	W5-O13	-2.02	1.68	2.16
3	B	401[A]	PW9	W5-O13	-2.02	1.68	2.16

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	401[A]	PW9	O19E-P1-O4	6.80	123.00	111.56
3	B	401[B]	PW9	O19E-P1-O4	6.76	122.92	111.56
3	A	307[B]	PW9	O19E-P1-O4	6.62	122.69	111.56
3	A	307[A]	PW9	O19E-P1-O4	6.58	122.63	111.56

There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.

4 monomers are involved in 24 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	307[B]	PW9	6	0
3	A	307[A]	PW9	6	0
3	B	401[A]	PW9	6	0
3	B	401[B]	PW9	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient


equivalents in the CSD to analyse the geometry.




Ligand PW9 A 307 (A)




Bond lengths



Bond angles




Torsions




Rings


Ligand PW9 B 401 (A)




Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	250/256 (97%)	1.03	32 (12%) 7 6	16, 25, 38, 50	0
1	B	249/256 (97%)	1.76	81 (32%) 1 1	14, 34, 50, 60	4 (1%)
All	All	499/512 (97%)	1.40	113 (22%) 2 1	14, 30, 46, 60	4 (0%)

All (113) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	84	GLY	7.4
1	B	251	ALA	7.4
1	A	84	GLY	7.0
1	B	16	SER	5.4
1	B	182	THR	5.4
1	B	226	SER	4.8
1	B	192	GLY	4.6
1	B	212	ASN	4.4
1	B	166	ALA	4.4
1	B	148	SER	4.2
1	B	142	SER	4.1
1	B	86	ASN	3.8
1	B	170	ASN	3.8
1	B	179	GLY	3.8
1	B	209	ALA	3.7
1	B	188	VAL	3.6
1	B	233	ALA	3.6
1	B	178	THR	3.6
1	B	167	ALA	3.5
1	A	199	HIS	3.5
1	B	191	ALA	3.5
1	B	181	ASN	3.4
1	A	184	SER	3.4
1	B	169	SER	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	171	THR	3.4
1	B	85	SER	3.4
1	B	184	SER	3.4
1	B	232	SER	3.3
1	B	58	SER	3.3
1	A	42	GLY	3.3
1	B	200	GLY	3.3
1	B	250	ALA	3.2
1	A	209	ALA	3.1
1	B	190	SER	3.1
1	A	129	THR	3.1
1	B	23	ALA	3.1
1	A	226	SER	3.1
1	B	17	GLY	3.1
1	B	87	THR	3.0
1	B	213	THR	3.0
1	B	123	LEU	3.0
1	B	230	VAL	2.9
1	B	129	THR	2.9
1	A	212	ASN	2.9
1	B	224	THR	2.9
1	B	14	THR	2.9
1	B	249	LEU	2.9
1	A	223	ASN	2.8
1	B	3	THR	2.8
1	B	197	THR	2.8
1	B	189	ASP	2.8
1	A	250	ALA	2.8
1	B	37	VAL	2.8
1	B	121	VAL	2.8
1	B	234	GLY	2.7
1	A	232	SER	2.7
1	B	65	ALA	2.7
1	B	210	GLY	2.7
1	A	128	ASN	2.7
1	B	238	VAL	2.7
1	B	208	ALA	2.6
1	A	240	ASP	2.6
1	B	52	THR	2.6
1	A	198	ASP	2.6
1	B	177	PHE	2.6
1	A	207	LEU	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	219	PHE	2.5
1	B	207	LEU	2.5
1	A	45	THR	2.5
1	B	217	LEU	2.5
1	B	195	TYR	2.5
1	A	213	THR	2.4
1	B	125	ALA	2.4
1	B	193	THR	2.4
1	A	44	ASN	2.4
1	A	181	ASN	2.4
1	B	100[A]	SER	2.4
1	B	155	THR	2.4
1	A	86	ASN	2.4
1	A	200	GLY	2.4
1	B	174	VAL	2.4
1	B	165	LEU	2.3
1	B	239	THR	2.3
1	B	168	GLY	2.3
1	B	22	SER	2.3
1	A	14	THR	2.3
1	B	124	ALA	2.3
1	B	139	ASN	2.3
1	A	210	GLY	2.2
1	B	176	PRO	2.2
1	B	198	ASP	2.2
1	A	231	ASP	2.2
1	A	215	THR	2.2
1	B	180	LEU	2.2
1	B	15	PRO	2.2
1	B	205	VAL	2.2
1	B	64	SER	2.2
1	B	49	LEU	2.2
1	A	158	GLY	2.1
1	B	9	PHE	2.1
1	A	1	SER	2.1
1	B	220	THR	2.1
1	B	194	VAL	2.1
1	B	218	PRO	2.1
1	A	2	ASN	2.1
1	A	179	GLY	2.1
1	B	24	GLY	2.0
1	B	152	VAL	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	216	VAL	2.0
1	A	165	LEU	2.0
1	A	219	PHE	2.0
1	B	51	PHE	2.0
1	A	224	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	K	A	308	1/1	0.55	0.30	45,45,45,45	0
4	K	A	310	1/1	0.58	0.31	56,56,56,56	0
4	K	A	312	1/1	0.61	0.29	56,56,56,56	0
4	K	A	313	1/1	0.63	0.26	52,52,52,52	0
4	K	A	311	1/1	0.70	0.21	51,51,51,51	0
3	PW9	B	401[A]	44/44	0.75	0.25	48,48,48,48	44
4	K	A	309	1/1	0.75	0.21	40,40,40,40	0
3	PW9	B	401[B]	44/44	0.75	0.25	48,48,48,48	44
3	PW9	A	307[A]	44/44	0.92	0.16	38,38,38,38	44
3	PW9	A	307[B]	44/44	0.92	0.16	38,38,38,38	44
2	CU	A	301	1/1	0.99	0.05	23,23,23,23	0
2	CU	A	302	1/1	0.99	0.02	26,26,26,26	0
2	CU	A	304	1/1	0.99	0.04	31,31,31,31	0
2	CU	A	305	1/1	0.99	0.03	26,26,26,26	0
2	CU	A	306	1/1	0.99	0.02	22,22,22,22	0
2	CU	A	303	1/1	1.00	0.06	33,33,33,33	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around PW9 B 401 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PW9 B 401 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PW9 A 307 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PW9 A 307 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CU A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CU A 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CU A 304:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CU A 305:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around CU A 306:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.