



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 17, 2026 – 06:13 PM UTC

PDB ID : 2OWM / pdb\_00002owm  
Title : Motor domain of Neurospora crassa kinesin-3 (NcKin3)  
Authors : Marx, A.; Muller, J.; Mandelkow, E.-M.; Woehlke, G.; Mandelkow, E.  
Deposited on : 2007-02-16  
Resolution : 3.25 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1605 (3.30-3.22)
Clashscore	190562	1660 (3.30-3.22)
Ramachandran outliers	187476	1630 (3.30-3.22)
Sidechain outliers	187428	1629 (3.30-3.22)
RSRZ outliers	180081	1605 (3.30-3.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	443	 41% 25% 7% 26%
1	B	443	 42% 24% 7% 26%
1	C	443	 39% 26% 8% 26%
1	D	443	 42% 25% 6% 27%

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 10338 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Related to KINESIN-LIKE PROTEIN KIF1C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	328	2560	1609	445	495	11	0	0	0
1	B	327	2561	1609	449	492	11	0	0	0
1	C	328	2560	1609	445	495	11	0	0	0
1	D	325	2545	1600	444	490	11	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	59	GLU	LYS	conflict	UNP Q9C2M3
A	83	ASP	ASN	conflict	UNP Q9C2M3
A	252	GLY	ARG	conflict	UNP Q9C2M3
B	59	GLU	LYS	conflict	UNP Q9C2M3
B	83	ASP	ASN	conflict	UNP Q9C2M3
B	252	GLY	ARG	conflict	UNP Q9C2M3
C	59	GLU	LYS	conflict	UNP Q9C2M3
C	83	ASP	ASN	conflict	UNP Q9C2M3
C	252	GLY	ARG	conflict	UNP Q9C2M3
D	59	GLU	LYS	conflict	UNP Q9C2M3
D	83	ASP	ASN	conflict	UNP Q9C2M3
D	252	GLY	ARG	conflict	UNP Q9C2M3


- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		
2	B	1	Total	Mg	0	0
			1	1		


*Continued on next page...*









● Molecule 1: Related to KINESIN-LIKE PROTEIN KIF1C



● Molecule 1: Related to KINESIN-LIKE PROTEIN KIF1C





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	75.94Å 98.41Å 111.76Å 90.00° 91.86° 90.00°	Depositor
Resolution (Å)	111.80 – 3.25 111.70 – 3.25	Depositor EDS
% Data completeness (in resolution range)	100.0 (111.80-3.25) 100.0 (111.70-3.25)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.45 (at 3.26Å)	Xtrriage
Refinement program	REFMAC refmac_5.2.0005	Depositor
R, $R_{free}$	0.210 , 0.256 0.209 , 0.252	Depositor DCC
$R_{free}$ test set	1329 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	69.8	Xtrriage
Anisotropy	0.329	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 78.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.028 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	10338	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	76.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.89% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.15	3/2610 (0.1%)	1.25	16/3546 (0.5%)
1	B	1.09	1/2611 (0.0%)	1.22	5/3546 (0.1%)
1	C	1.09	2/2610 (0.1%)	1.23	19/3546 (0.5%)
1	D	0.97	0/2595	1.17	7/3525 (0.2%)
All	All	1.08	6/10426 (0.1%)	1.22	47/14163 (0.3%)

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	157	THR	CA-CB	-7.49	1.44	1.54
1	A	213	PRO	CA-C	6.01	1.57	1.52
1	A	182	THR	CA-CB	5.57	1.60	1.53
1	C	182	THR	CA-CB	5.11	1.59	1.53
1	A	337	ALA	CA-CB	-5.07	1.45	1.53

The worst 5 of 47 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	323	ASN	N-CA-C	7.84	123.11	112.68
1	C	226	GLY	CA-C-N	7.66	127.61	120.03
1	C	226	GLY	C-N-CA	7.66	127.61	120.03
1	A	323	ASN	N-CA-C	7.64	123.09	112.93
1	C	94	GLU	N-CA-C	7.19	119.49	108.42

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2560	0	2475	121	1
1	B	2561	0	2482	122	0
1	C	2560	0	2475	121	0
1	D	2545	0	2464	96	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	27	0	12	3	0
3	B	27	0	12	1	0
3	C	27	0	12	1	0
3	D	27	0	12	0	0
All	All	10338	0	9944	444	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 22.

The worst 5 of 444 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:157:THR:HG23	1:A:158:PRO:HD2	1.32	1.10
1:B:305:ARG:HH11	1:B:305:ARG:HG3	1.02	1.07
1:B:412:ARG:HG2	1:B:412:ARG:HH11	1.17	1.05
1:A:132:ASN:HD21	1:A:138:THR:HB	1.22	1.04
1:C:157:THR:HG23	1:C:158:PRO:HD2	1.39	1.04

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:63:GLU:OE2	1:A:220:ARG:NH1[2_646]	1.97	0.23

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	318/443 (72%)	286 (90%)	26 (8%)	6 (2%)	6	28
1	B	317/443 (72%)	282 (89%)	29 (9%)	6 (2%)	6	28
1	C	318/443 (72%)	288 (91%)	24 (8%)	6 (2%)	6	28
1	D	315/443 (71%)	284 (90%)	25 (8%)	6 (2%)	6	28
All	All	1268/1772 (72%)	1140 (90%)	104 (8%)	24 (2%)	6	28

5 of 24 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	67	ALA
1	A	210	PRO
1	A	324	LYS
1	B	304	GLU
1	B	324	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	278/389 (72%)	221 (80%)	57 (20%)	1	5
1	B	278/389 (72%)	219 (79%)	59 (21%)	1	4
1	C	278/389 (72%)	223 (80%)	55 (20%)	1	5
1	D	277/389 (71%)	219 (79%)	58 (21%)	1	5
All	All	1111/1556 (71%)	882 (79%)	229 (21%)	1	5

5 of 229 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	418	VAL
1	D	400	THR
1	C	237	VAL
1	D	386	THR
1	D	237	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	173	GLN
1	D	323	ASN
1	D	280	HIS
1	B	383	ASN
1	C	179	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	ADP	A	1002	-	28,29,29	1.49	5 (17%)	43,45,45	2.07	10 (23%)
3	ADP	B	1002	2	28,29,29	1.32	3 (10%)	43,45,45	2.01	13 (30%)
3	ADP	D	1002	2	28,29,29	1.50	4 (14%)	43,45,45	2.18	12 (27%)
3	ADP	C	1002	2	28,29,29	1.46	4 (14%)	43,45,45	2.17	12 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	A	1002	-	-	0/16/32/32	0/3/3/3
3	ADP	B	1002	2	-	0/16/32/32	0/3/3/3
3	ADP	D	1002	2	-	0/16/32/32	0/3/3/3
3	ADP	C	1002	2	-	0/16/32/32	0/3/3/3

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	1002	ADP	C5-C4	4.51	1.47	1.39
3	A	1002	ADP	C5-N7	-4.11	1.31	1.39
3	C	1002	ADP	C5-C4	4.08	1.46	1.39
3	B	1002	ADP	C5-C4	3.85	1.46	1.39
3	A	1002	ADP	C5-C4	3.80	1.45	1.39

The worst 5 of 47 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1002	ADP	C5-C4-N3	-6.80	117.36	126.72
3	D	1002	ADP	C5-C4-N3	-6.78	117.39	126.72
3	D	1002	ADP	N3-C4-N9	6.29	137.87	127.17
3	A	1002	ADP	N3-C4-N9	6.02	137.41	127.17
3	B	1002	ADP	C5-C4-N3	-5.98	118.48	126.72

There are no chirality outliers.


There are no torsion outliers.



There are no ring outliers.


3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1002	ADP	3	0
3	B	1002	ADP	1	0
3	C	1002	ADP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	328/443 (74%)	-0.03	3 (0%) 81 65	70, 76, 79, 83	0
1	B	327/443 (73%)	0.03	3 (0%) 81 65	68, 76, 79, 90	0
1	C	328/443 (74%)	0.41	8 (2%) 59 40	70, 76, 79, 83	0
1	D	325/443 (73%)	0.57	12 (3%) 45 30	67, 76, 79, 82	0
All	All	1308/1772 (73%)	0.24	26 (1%) 65 46	67, 76, 79, 90	0

The worst 5 of 26 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	281	HIS	4.4
1	C	320	SER	4.4
1	D	288	THR	3.9
1	C	288	THR	3.6
1	A	320	SER	3.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MG	B	1001	1/1	0.83	0.14	61,61,61,61	0
3	ADP	D	1002	27/27	0.88	0.13	72,75,76,77	0
2	MG	D	1001	1/1	0.90	0.13	61,61,61,61	0
3	ADP	B	1002	27/27	0.93	0.10	72,75,77,77	0
3	ADP	C	1002	27/27	0.93	0.10	72,75,77,77	0
2	MG	A	1001	1/1	0.93	0.21	61,61,61,61	0
3	ADP	A	1002	27/27	0.94	0.09	72,75,77,77	0
2	MG	C	1001	1/1	0.94	0.18	60,60,60,60	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around ADP B 1002:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around ADP C 1002:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.