



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2026 – 05:08 PM UTC

PDB ID : 7OW3 / pdb_00007ow3
Title : Crystal structure of HLA-A*11:01 in complex with KRAS peptide (VVVGAG-GVGK)
Authors : Coles, C.H.; Karuppiah, V.; Robinson, R.A.
Deposited on : 2021-06-16
Resolution : 2.46 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.46 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1190 (2.46-2.46)
Clashscore	190562	1229 (2.46-2.46)
Ramachandran outliers	187476	1218 (2.46-2.46)
Sidechain outliers	187428	1218 (2.46-2.46)
RSRZ outliers	180081	1190 (2.46-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	6% 90% 9%
1	D	275	3% 94% 6%
1	G	275	10% 93% 6%
1	J	275	5% 89% 9%
2	B	100	0% 94% 6%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	E	100	% 92% 8%
2	H	100	5% 88% 12%
2	K	100	% 93% 7%
3	C	10	30% 90% 10%
3	F	10	70% 30%
3	I	10	10% 50% 10% 40%
3	L	10	10% 100%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 12705 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MHC class I antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	274	Total	C	N	O	S	0	0	0
			2240	1392	407	432	9			
1	D	274	Total	C	N	O	S	0	0	0
			2240	1392	407	432	9			
1	G	274	Total	C	N	O	S	0	0	0
			2240	1392	407	432	9			
1	J	275	Total	C	N	O	S	0	0	0
			2244	1394	408	433	9			

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			
2	E	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			
2	H	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			
2	K	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	initiating methionine	UNP P61769
E	0	MET	-	initiating methionine	UNP P61769
H	0	MET	-	initiating methionine	UNP P61769
K	0	MET	-	initiating methionine	UNP P61769

- Molecule 3 is a protein called KRAS peptide (VVVGAGGVGK).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	10	Total	C	N	O	0	0	0
			59	37	11	11			
3	F	7	Total	C	N	O	0	0	0
			46	30	8	8			
3	I	6	Total	C	N	O	0	0	0
			42	28	7	7			
3	L	10	Total	C	N	O	0	0	0
			59	37	11	11			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	44	Total	O	0	0
			44	44		
4	B	13	Total	O	0	0
			13	13		
4	D	31	Total	O	0	0
			31	31		
4	E	16	Total	O	0	0
			16	16		
4	F	1	Total	O	0	0
			1	1		
4	G	24	Total	O	0	0
			24	24		
4	H	7	Total	O	0	0
			7	7		
4	J	37	Total	O	0	0
			37	37		
4	K	14	Total	O	0	0
			14	14		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MHC class I antigen




- Molecule 1: MHC class I antigen



- Molecule 1: MHC class I antigen





- Molecule 1: MHC class I antigen






- Molecule 2: Beta-2-microglobulin



- Molecule 2: Beta-2-microglobulin




- Molecule 2: Beta-2-microglobulin



- Molecule 2: Beta-2-microglobulin




- Molecule 3: KRAS peptide (VVVGAGGVGK)




- Molecule 3: KRAS peptide (VVVGAGGVGK)



- Molecule 3: KRAS peptide (VVVGAGGVGK)



- Molecule 3: KRAS peptide (VVVGAGGVGK)



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	71.85Å 117.03Å 110.64Å 90.00° 99.30° 90.00°	Depositor
Resolution (Å)	79.96 – 2.46 79.96 – 2.46	Depositor EDS
% Data completeness (in resolution range)	99.9 (79.96-2.46) 99.9 (79.96-2.46)	Depositor EDS
R_{merge}	0.19	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.86 (at 2.45Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.228 , 0.260 0.236 , 0.269	Depositor DCC
R_{free} test set	3237 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	37.2	Xtriage
Anisotropy	0.161	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 29.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	12705	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.09	1/2301 (0.0%)	1.40	4/3124 (0.1%)
1	D	1.09	0/2301	1.37	7/3124 (0.2%)
1	G	1.07	0/2301	1.40	6/3124 (0.2%)
1	J	1.04	1/2305 (0.0%)	1.38	5/3129 (0.2%)
2	B	1.12	0/860	1.30	0/1162
2	E	1.10	1/860 (0.1%)	1.32	2/1162 (0.2%)
2	H	1.07	0/860	1.29	1/1162 (0.1%)
2	K	1.09	0/860	1.31	0/1162
3	C	1.14	0/58	1.95	2/75 (2.7%)
3	F	1.06	0/44	1.53	0/55
3	I	0.99	0/40	1.27	0/50
3	L	1.20	0/58	1.45	0/75
All	All	1.08	3/12848 (0.0%)	1.37	27/17404 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	197	HIS	CE1-NE2	5.83	1.38	1.32
2	E	13	HIS	CE1-NE2	5.79	1.38	1.32
1	J	192	HIS	CE1-NE2	5.75	1.38	1.32

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	J	111	ARG	CA-C-N	8.17	127.82	122.18
1	J	111	ARG	C-N-CA	8.17	127.82	122.18
1	D	226	GLN	CA-C-O	-7.66	112.96	121.00
1	A	111	ARG	CA-C-N	6.64	126.76	122.18
1	A	111	ARG	C-N-CA	6.64	126.76	122.18
1	J	128	GLU	CB-CG-CD	6.38	123.44	112.60
1	G	14	ARG	N-CA-C	-5.74	102.93	110.39
1	D	110	LEU	N-CA-C	-5.71	106.97	114.04
2	H	83	ASN	CA-CB-CG	-5.66	106.94	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	116	ASP	CA-CB-CG	5.59	118.19	112.60
3	C	4	GLY	CA-C-N	5.56	127.67	120.44
3	C	4	GLY	C-N-CA	5.56	127.67	120.44
1	A	116	ASP	CA-CB-CG	5.41	118.01	112.60
1	G	111	ARG	CA-C-N	5.33	125.86	122.18
1	G	111	ARG	C-N-CA	5.33	125.86	122.18
1	D	82	ARG	CA-C-N	5.30	125.98	119.99
1	D	82	ARG	C-N-CA	5.30	125.98	119.99
1	J	116	ASP	CA-CB-CG	5.28	117.88	112.60
1	A	186	LYS	N-CA-C	-5.22	100.79	109.46
1	D	111	ARG	N-CA-C	5.22	116.35	108.46
1	J	230	LEU	CA-C-O	-5.22	114.92	120.40
1	G	231	VAL	CA-C-N	5.18	128.12	120.82
1	G	231	VAL	C-N-CA	5.18	128.12	120.82
1	D	227	ASP	CB-CA-C	5.10	117.98	111.40
2	E	19	LYS	CA-C-N	5.08	128.06	120.90
2	E	19	LYS	C-N-CA	5.08	128.06	120.90
1	G	18	GLY	CA-C-O	-5.04	117.43	121.77

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2240	0	2085	15	0
1	D	2240	0	2085	4	0
1	G	2240	0	2085	6	0
1	J	2244	0	2091	12	0
2	B	837	0	803	5	0
2	E	837	0	803	3	0
2	H	837	0	803	4	0
2	K	837	0	803	3	0
3	C	59	0	68	0	0
3	F	46	0	56	0	0
3	I	42	0	53	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	L	59	0	68	0	0
4	A	44	0	0	1	0
4	B	13	0	0	0	0
4	D	31	0	0	0	0
4	E	16	0	0	1	0
4	F	1	0	0	0	0
4	G	24	0	0	0	0
4	H	7	0	0	0	0
4	J	37	0	0	0	0
4	K	14	0	0	0	0
All	All	12705	0	11803	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:9:TYR:CE2	1:A:70:GLN:HG2	2.35	0.62
1:D:129:ASP:OD2	2:H:48:LYS:HE3	2.00	0.60
1:J:9:TYR:CE2	1:J:70:GLN:HG2	2.36	0.60
1:A:19:GLU:HG3	1:A:20:PRO:HD2	1.85	0.58
1:G:9:TYR:CE2	1:G:70:GLN:HG2	2.38	0.57
1:A:14:ARG:HB3	1:A:17:ARG:HB2	1.87	0.56
1:J:82:ARG:HH11	1:J:82:ARG:HB3	1.72	0.55
1:D:105:PRO:HD2	1:G:238:ASP:O	2.05	0.55
1:A:238:ASP:O	1:J:105:PRO:HD2	2.08	0.53
2:E:74:GLU:HG2	4:E:112:HOH:O	2.08	0.53
1:G:63:GLU:OE1	3:I:2:VAL:HG22	2.09	0.53
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.46	0.50
1:A:19:GLU:HG3	1:A:20:PRO:CD	2.41	0.50
1:J:117:ALA:HB2	2:K:60:TRP:CE2	2.48	0.49
1:A:63:GLU:OE2	4:A:301:HOH:O	2.20	0.47
1:A:270:LEU:HD11	1:J:267:PRO:HB2	1.96	0.47
1:J:275:GLU:C	1:J:275:GLU:CD	2.82	0.47
1:A:19:GLU:HG3	1:A:20:PRO:N	2.29	0.47
2:H:51:HIS:HA	2:H:65:LEU:O	2.15	0.47
2:B:51:HIS:HA	2:B:65:LEU:O	2.15	0.46
1:J:42:SER:O	1:J:43:GLN:HB2	2.14	0.46
1:A:31:THR:HG23	1:J:108:ARG:HH11	1.79	0.46
1:D:192:HIS:NE2	2:E:98:ASP:OD2	2.48	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:J:220:ASP:OD2	1:J:256:ARG:NH2	2.48	0.46
2:K:51:HIS:HA	2:K:65:LEU:O	2.16	0.46
2:E:51:HIS:HA	2:E:65:LEU:O	2.17	0.45
1:A:49:ALA:O	1:A:52:ILE:HG22	2.18	0.44
2:H:21:ASN:HB3	2:H:70:PHE:CE1	2.53	0.43
1:D:9:TYR:CE2	1:D:70:GLN:HG2	2.54	0.43
1:G:14:ARG:H	1:G:14:ARG:HG2	1.53	0.43
1:J:82:ARG:HA	1:J:87:GLN:HE21	1.85	0.42
2:B:27:VAL:O	2:B:63:TYR:HA	2.20	0.42
1:G:117:ALA:HB2	2:H:60:TRP:CE2	2.55	0.42
1:A:187:THR:HA	1:A:204:TRP:O	2.20	0.42
1:G:187:THR:HA	1:G:204:TRP:O	2.20	0.42
1:A:187:THR:HG21	1:A:270:LEU:HD23	2.02	0.41
2:B:27:VAL:HG11	2:B:35:ILE:HD13	2.03	0.41
2:B:27:VAL:HG11	2:B:35:ILE:CD1	2.50	0.41
1:J:192:HIS:NE2	2:K:98:ASP:OD2	2.50	0.41
1:J:187:THR:HA	1:J:204:TRP:O	2.22	0.40
1:A:220:ASP:OD2	1:A:256:ARG:NH2	2.46	0.40
1:A:43:GLN:C	1:A:44:ARG:HG2	2.47	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	272/275 (99%)	266 (98%)	6 (2%)	0	100	100
1	D	272/275 (99%)	264 (97%)	8 (3%)	0	100	100
1	G	272/275 (99%)	266 (98%)	6 (2%)	0	100	100
1	J	273/275 (99%)	266 (97%)	7 (3%)	0	100	100
2	B	98/100 (98%)	97 (99%)	1 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	E	98/100 (98%)	97 (99%)	1 (1%)	0	100	100
2	H	98/100 (98%)	97 (99%)	1 (1%)	0	100	100
2	K	98/100 (98%)	97 (99%)	1 (1%)	0	100	100
3	C	8/10 (80%)	6 (75%)	2 (25%)	0	100	100
3	F	3/10 (30%)	2 (67%)	1 (33%)	0	100	100
3	I	2/10 (20%)	1 (50%)	1 (50%)	0	100	100
3	L	8/10 (80%)	8 (100%)	0	0	100	100
All	All	1502/1540 (98%)	1467 (98%)	35 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	232/232 (100%)	227 (98%)	5 (2%)	45	60
1	D	232/232 (100%)	227 (98%)	5 (2%)	45	60
1	G	232/232 (100%)	225 (97%)	7 (3%)	36	51
1	J	232/232 (100%)	221 (95%)	11 (5%)	23	35
2	B	95/95 (100%)	95 (100%)	0	100	100
2	E	95/95 (100%)	93 (98%)	2 (2%)	47	62
2	H	95/95 (100%)	90 (95%)	5 (5%)	20	30
2	K	95/95 (100%)	92 (97%)	3 (3%)	34	49
3	C	5/5 (100%)	5 (100%)	0	100	100
3	F	5/5 (100%)	5 (100%)	0	100	100
3	I	5/5 (100%)	5 (100%)	0	100	100
3	L	5/5 (100%)	5 (100%)	0	100	100
All	All	1328/1328 (100%)	1290 (97%)	38 (3%)	37	52

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	17	ARG
1	A	58	GLU
1	A	110	LEU
1	A	222	GLU
1	A	255	GLN
1	D	2	SER
1	D	67	VAL
1	D	113	TYR
1	D	182	THR
1	D	264	GLU
2	E	89	GLN
2	E	92	ILE
1	G	14	ARG
1	G	17	ARG
1	G	58	GLU
1	G	89	GLU
1	G	110	LEU
1	G	255	GLN
1	G	268	LYS
2	H	1	ILE
2	H	27	VAL
2	H	69	GLU
2	H	91	LYS
2	H	92	ILE
1	J	2	SER
1	J	15	PRO
1	J	34	VAL
1	J	82	ARG
1	J	113	TYR
1	J	154	GLU
1	J	165	VAL
1	J	182	THR
1	J	255	GLN
1	J	264	GLU
1	J	275	GLU
2	K	9	VAL
2	K	27	VAL
2	K	92	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	54	GLN
1	A	155	GLN
1	A	188	HIS
1	A	226	GLN
1	A	262	GLN
1	D	43	GLN
1	D	141	GLN
1	D	151	HIS
1	G	62	GLN
1	G	226	GLN
1	G	262	GLN
1	J	87	GLN
1	J	155	GLN
1	J	226	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	274/275 (99%)	0.57	17 (6%) 26 24	18, 35, 52, 76	0
1	D	274/275 (99%)	0.57	7 (2%) 57 58	23, 37, 57, 71	0
1	G	274/275 (99%)	1.01	28 (10%) 12 10	21, 46, 66, 81	0
1	J	275/275 (100%)	0.47	13 (4%) 36 35	23, 33, 54, 66	0
2	B	100/100 (100%)	0.30	1 (1%) 79 80	20, 30, 48, 54	0
2	E	100/100 (100%)	0.31	1 (1%) 79 80	23, 31, 42, 50	0
2	H	100/100 (100%)	0.57	5 (5%) 34 32	23, 39, 59, 65	0
2	K	100/100 (100%)	0.38	1 (1%) 79 80	20, 33, 47, 53	0
3	C	10/10 (100%)	1.28	3 (30%) 1 1	46, 50, 57, 58	0
3	F	7/10 (70%)	0.87	0 100 100	39, 41, 44, 44	0
3	I	6/10 (60%)	1.66	1 (16%) 4 3	50, 52, 55, 59	0
3	L	10/10 (100%)	1.09	1 (10%) 12 10	34, 40, 49, 50	0
All	All	1530/1540 (99%)	0.59	78 (5%) 33 31	18, 36, 59, 81	0

All (78) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	16	GLY	5.2
1	G	17	ARG	4.5
1	G	15	PRO	4.4
1	A	16	GLY	4.1
1	A	15	PRO	4.0
1	A	269	PRO	3.8
1	G	19	GLU	3.8
1	D	223	ASP	3.7
1	A	197	HIS	3.4
1	A	138	MET	3.3
1	G	269	PRO	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	G	20	PRO	3.3
1	D	90	ASP	3.2
1	D	275	GLU	3.2
1	G	58	GLU	3.2
1	D	225	THR	3.1
1	G	83	GLY	3.0
3	L	5	ALA	3.0
1	A	275	GLU	2.9
3	C	5	ALA	2.9
1	J	274	TRP	2.9
1	A	17	ARG	2.8
1	G	133	TRP	2.8
1	J	225	THR	2.8
2	H	69	GLU	2.8
1	G	74	ASP	2.8
3	C	7	GLY	2.7
1	G	90	ASP	2.7
1	G	225	THR	2.7
1	G	138	MET	2.7
1	G	89	GLU	2.7
1	A	196	ASP	2.6
3	C	1	VAL	2.6
1	A	251	SER	2.6
1	D	226	GLN	2.5
1	G	194	ILE	2.5
1	A	194	ILE	2.5
1	G	192	HIS	2.5
1	G	131	ARG	2.5
1	G	150	ALA	2.5
1	G	91	GLY	2.4
2	B	75	LYS	2.4
1	G	54	GLN	2.4
1	J	89	GLU	2.4
1	G	179	LEU	2.4
1	G	65	ARG	2.3
1	G	196	ASP	2.3
1	A	192	HIS	2.3
1	G	18	GLY	2.3
2	H	75	LYS	2.3
3	I	1	VAL	2.3
1	J	181	ARG	2.3
1	J	194	ILE	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	J	275	GLU	2.2
1	A	199	ALA	2.2
1	J	155	GLN	2.2
1	J	182	THR	2.2
1	A	128	GLU	2.2
1	J	224	GLN	2.2
1	D	221	GLY	2.2
1	D	128	GLU	2.2
1	G	268	LYS	2.2
2	H	0	MET	2.1
1	A	212	GLU	2.1
1	A	268	LYS	2.1
1	J	16	GLY	2.1
2	H	47	GLU	2.1
1	G	42	SER	2.1
1	J	196	ASP	2.1
2	H	43	GLY	2.1
1	A	89	GLU	2.1
1	J	177	GLU	2.1
2	K	69	GLU	2.1
1	G	98	MET	2.1
1	A	177	GLU	2.0
2	E	48	LYS	2.0
1	J	197	HIS	2.0
1	G	139	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.