



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 07:41 PM UTC

PDB ID : 2OXH / pdb_00002oxh
Title : The SOXYZ Complex of Paracoccus Pantotrophus
Authors : Sauve, V.; Berks, B.C.; Hemmings, A.M.
Deposited on : 2007-02-20
Resolution : 2.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1596 (2.36-2.36)
Clashscore	190562	1663 (2.36-2.36)
Ramachandran outliers	187476	1646 (2.36-2.36)
Sidechain outliers	187428	1646 (2.36-2.36)
RSRZ outliers	180081	1598 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	107	8% 83% 17%
1	C	107	85% 14% .
1	E	107	13% 83% 17%
1	Z	107	91% 9%
2	B	124	% 77% 15% 7%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	124	
2	F	124	
2	Y	124	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	D	603	-	-	X	-
3	EDO	D	604	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6822 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SoxZ protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Z	107	Total	C	N	O	S	0	1	0
			820	517	136	165	2			
1	A	107	Total	C	N	O	S	0	0	0
			817	515	136	164	2			
1	C	107	Total	C	N	O	S	0	1	0
			823	518	137	166	2			
1	E	107	Total	C	N	O	S	0	0	0
			814	514	136	162	2			

- Molecule 2 is a protein called SoxY protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	Y	111	Total	C	N	O	S	0	2	0
			784	485	135	160	4			
2	B	115	Total	C	N	O	S	0	2	0
			798	492	137	165	4			
2	D	110	Total	C	N	O	S	0	0	0
			764	473	131	156	4			
2	F	114	Total	C	N	O	S	0	0	0
			783	482	135	162	4			

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
Y	-11	MET	-	cloning artifact	UNP Q9LCU9
Y	-10	ARG	-	cloning artifact	UNP Q9LCU9
Y	-9	GLY	-	cloning artifact	UNP Q9LCU9
Y	-8	SER	-	cloning artifact	UNP Q9LCU9
Y	-7	HIS	-	expression tag	UNP Q9LCU9
Y	-6	HIS	-	expression tag	UNP Q9LCU9
Y	-5	HIS	-	expression tag	UNP Q9LCU9
Y	-4	HIS	-	expression tag	UNP Q9LCU9

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
Y	-3	HIS	-	expression tag	UNP Q9LCU9
Y	-2	HIS	-	expression tag	UNP Q9LCU9
Y	-1	GLY	-	cloning artifact	UNP Q9LCU9
Y	0	SER	-	cloning artifact	UNP Q9LCU9
Y	110	CSU	CYS	modified residue	UNP Q9LCU9
B	-11	MET	-	cloning artifact	UNP Q9LCU9
B	-10	ARG	-	cloning artifact	UNP Q9LCU9
B	-9	GLY	-	cloning artifact	UNP Q9LCU9
B	-8	SER	-	cloning artifact	UNP Q9LCU9
B	-7	HIS	-	expression tag	UNP Q9LCU9
B	-6	HIS	-	expression tag	UNP Q9LCU9
B	-5	HIS	-	expression tag	UNP Q9LCU9
B	-4	HIS	-	expression tag	UNP Q9LCU9
B	-3	HIS	-	expression tag	UNP Q9LCU9
B	-2	HIS	-	expression tag	UNP Q9LCU9
B	-1	GLY	-	cloning artifact	UNP Q9LCU9
B	0	SER	-	cloning artifact	UNP Q9LCU9
B	110	CSU	CYS	modified residue	UNP Q9LCU9
D	-11	MET	-	cloning artifact	UNP Q9LCU9
D	-10	ARG	-	cloning artifact	UNP Q9LCU9
D	-9	GLY	-	cloning artifact	UNP Q9LCU9
D	-8	SER	-	cloning artifact	UNP Q9LCU9
D	-7	HIS	-	expression tag	UNP Q9LCU9
D	-6	HIS	-	expression tag	UNP Q9LCU9
D	-5	HIS	-	expression tag	UNP Q9LCU9
D	-4	HIS	-	expression tag	UNP Q9LCU9
D	-3	HIS	-	expression tag	UNP Q9LCU9
D	-2	HIS	-	expression tag	UNP Q9LCU9
D	-1	GLY	-	cloning artifact	UNP Q9LCU9
D	0	SER	-	cloning artifact	UNP Q9LCU9
D	110	CSU	CYS	modified residue	UNP Q9LCU9
F	-11	MET	-	cloning artifact	UNP Q9LCU9
F	-10	ARG	-	cloning artifact	UNP Q9LCU9
F	-9	GLY	-	cloning artifact	UNP Q9LCU9
F	-8	SER	-	cloning artifact	UNP Q9LCU9
F	-7	HIS	-	expression tag	UNP Q9LCU9
F	-6	HIS	-	expression tag	UNP Q9LCU9
F	-5	HIS	-	expression tag	UNP Q9LCU9
F	-4	HIS	-	expression tag	UNP Q9LCU9
F	-3	HIS	-	expression tag	UNP Q9LCU9
F	-2	HIS	-	expression tag	UNP Q9LCU9
F	-1	GLY	-	cloning artifact	UNP Q9LCU9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	0	SER	-	cloning artifact	UNP Q9LCU9
F	110	CSU	CYS	modified residue	UNP Q9LCU9

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	Y	1	Total C O 4 2 2	0	0
3	Y	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	Z	78	Total O 78 78	0	0
4	Y	50	Total O 50 50	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	27	Total 27	O 27	0	0
4	B	42	Total 42	O 42	0	0
4	C	60	Total 60	O 60	0	0
4	D	54	Total 54	O 54	0	0
4	E	29	Total 29	O 29	0	0
4	F	55	Total 55	O 55	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: SoxZ protein

Chain Z:  91% 9%



- Molecule 1: SoxZ protein

Chain A:  8% 83% 17%




- Molecule 1: SoxZ protein

Chain C:  85% 14% .



- Molecule 1: SoxZ protein

Chain E:  13% 83% 17%




- Molecule 2: SoxY protein


Chain Y:  % 77% 11% . 10%




- Molecule 2: SoxY protein



- Molecule 2: SoxY protein



- Molecule 2: SoxY protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	40.22Å 54.49Å 104.87Å 103.13° 95.21° 90.01°	Depositor
Resolution (Å)	50.00 – 2.35 50.00 – 2.35	Depositor EDS
% Data completeness (in resolution range)	96.4 (50.00-2.35) 96.4 (50.00-2.35)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.17 (at 2.34Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.194 , 0.244 0.195 , 0.243	Depositor DCC
R_{free} test set	1754 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å ²)	16.6	Xtriage
Anisotropy	0.195	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 36.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.104 for -h,k,-k-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	6822	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, CSU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.70	0/833	0.88	0/1130
1	C	0.78	0/839	0.86	0/1138
1	E	0.71	0/830	0.86	0/1126
1	Z	0.76	0/836	0.84	0/1136
2	B	0.78	1/798 (0.1%)	0.86	1/1085 (0.1%)
2	D	0.84	0/762	0.88	0/1041
2	F	0.83	0/780	0.90	0/1061
2	Y	0.83	0/788	0.87	0/1075
All	All	0.78	1/6466 (0.0%)	0.87	1/8792 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	103	VAL	C-N	5.23	1.40	1.33

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	83	GLN	N-CA-C	5.58	116.61	108.14

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	108	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	817	0	798	9	0
1	C	823	0	802	13	0
1	E	814	0	796	10	0
1	Z	820	0	795	5	0
2	B	798	0	807	11	0
2	D	764	0	777	7	0
2	F	783	0	791	15	0
2	Y	784	0	801	9	0
3	A	4	0	6	1	0
3	B	4	0	6	2	0
3	D	8	0	12	11	0
3	Y	8	0	12	3	0
4	A	27	0	0	0	0
4	B	42	0	0	0	0
4	C	60	0	0	2	0
4	D	54	0	0	1	0
4	E	29	0	0	4	0
4	F	55	0	0	1	0
4	Y	50	0	0	1	0
4	Z	78	0	0	2	0
All	All	6822	0	6403	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 78 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:48:MET:HE2	2:F:88:LEU:HD12	1.22	1.17
3:D:603:EDO:H11	2:F:63:ASN:ND2	1.67	1.08
2:B:48:MET:HE2	2:B:88:LEU:HD12	1.35	1.07

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:603:EDO:C1	2:F:63:ASN:HD21	1.76	0.98
3:D:603:EDO:H11	2:F:63:ASN:HD21	0.81	0.97

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	105/107 (98%)	98 (93%)	7 (7%)	0	100	100
1	C	106/107 (99%)	101 (95%)	5 (5%)	0	100	100
1	E	105/107 (98%)	99 (94%)	6 (6%)	0	100	100
1	Z	106/107 (99%)	103 (97%)	3 (3%)	0	100	100
2	B	114/124 (92%)	109 (96%)	5 (4%)	0	100	100
2	D	108/124 (87%)	105 (97%)	3 (3%)	0	100	100
2	F	111/124 (90%)	107 (96%)	4 (4%)	0	100	100
2	Y	111/124 (90%)	107 (96%)	4 (4%)	0	100	100
All	All	866/924 (94%)	829 (96%)	37 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	88/89 (99%)	84 (96%)	4 (4%)	24	32
1	C	89/89 (100%)	86 (97%)	3 (3%)	32	43
1	E	87/89 (98%)	83 (95%)	4 (5%)	24	31
1	Z	88/89 (99%)	85 (97%)	3 (3%)	32	43
2	B	77/84 (92%)	74 (96%)	3 (4%)	28	39
2	D	74/84 (88%)	71 (96%)	3 (4%)	27	36
2	F	75/84 (89%)	73 (97%)	2 (3%)	39	52
2	Y	77/84 (92%)	73 (95%)	4 (5%)	21	25
All	All	655/692 (95%)	629 (96%)	26 (4%)	28	37

5 of 26 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	25	LEU
2	D	27	GLU
2	F	0	SER
1	C	61	VAL
2	D	81	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	81	GLN
2	F	31	ASN
2	F	81	GLN
2	F	63	ASN
2	B	31	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	CSU	Y	110	2	7,9,10	1.21	1 (14%)	4,12,14	0.62	0
2	CSU	B	110	2	7,9,10	0.80	0	4,12,14	1.18	1 (25%)
2	CSU	D	110	2	7,9,10	1.10	1 (14%)	4,12,14	0.69	0
2	CSU	F	110	2	7,9,10	0.71	0	4,12,14	1.35	1 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CSU	Y	110	2	-	2/4/8/10	-
2	CSU	B	110	2	-	0/4/8/10	-
2	CSU	D	110	2	-	2/4/8/10	-
2	CSU	F	110	2	-	2/4/8/10	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	Y	110	CSU	OD3-S	2.24	1.52	1.45
2	D	110	CSU	OD3-S	2.04	1.51	1.45

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	110	CSU	OD2-S-OD1	-2.11	105.70	112.74
2	B	110	CSU	OD2-S-OD1	-2.00	106.06	112.74

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	Y	110	CSU	OD2-S-SG-CB
2	D	110	CSU	OD1-S-SG-CB
2	D	110	CSU	OD2-S-SG-CB

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	F	110	CSU	OD2-S-SG-CB
2	Y	110	CSU	OD3-S-SG-CB

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	110	CSU	1	0
2	F	110	CSU	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	D	603	-	3,3,3	0.64	0	2,2,2	0.21	0
3	EDO	Y	601	-	3,3,3	0.62	0	2,2,2	0.14	0
3	EDO	B	602	-	3,3,3	0.42	0	2,2,2	0.11	0
3	EDO	Y	605	-	3,3,3	0.33	0	2,2,2	0.49	0
3	EDO	A	606	-	3,3,3	0.70	0	2,2,2	0.17	0
3	EDO	D	604	-	3,3,3	0.51	0	2,2,2	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	D	603	-	-	1/1/1/1	-
3	EDO	Y	601	-	-	1/1/1/1	-
3	EDO	B	602	-	-	1/1/1/1	-
3	EDO	Y	605	-	-	1/1/1/1	-
3	EDO	A	606	-	-	1/1/1/1	-
3	EDO	D	604	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	Y	601	EDO	O1-C1-C2-O2
3	Y	605	EDO	O1-C1-C2-O2
3	A	606	EDO	O1-C1-C2-O2
3	D	604	EDO	O1-C1-C2-O2
3	D	603	EDO	O1-C1-C2-O2

There are no ring outliers.

6 monomers are involved in 17 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	603	EDO	7	0
3	Y	601	EDO	2	0
3	B	602	EDO	2	0
3	Y	605	EDO	1	0
3	A	606	EDO	1	0
3	D	604	EDO	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	107/107 (100%)	0.53	9 (8%) 17 19	6, 18, 46, 58	0
1	C	107/107 (100%)	-0.19	0 100 100	4, 10, 28, 40	1 (0%)
1	E	107/107 (100%)	0.72	14 (13%) 7 8	6, 21, 52, 58	0
1	Z	107/107 (100%)	-0.16	0 100 100	3, 11, 27, 38	1 (0%)
2	B	114/124 (91%)	-0.04	1 (0%) 81 84	4, 12, 24, 30	2 (1%)
2	D	109/124 (87%)	-0.30	0 100 100	6, 9, 15, 26	0
2	F	113/124 (91%)	-0.05	0 100 100	7, 11, 19, 32	0
2	Y	110/124 (88%)	-0.14	1 (0%) 81 84	4, 9, 16, 25	2 (1%)
All	All	874/924 (94%)	0.04	25 (2%) 53 60	3, 11, 34, 58	6 (0%)

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	38	ALA	5.9
1	E	1	ALA	4.5
1	A	38	ALA	3.9
1	E	4	ALA	3.5
2	B	-1	GLY	3.2

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CSU	Y	110	10/11	0.78	0.21	29,32,43,43	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CSU	D	110	10/11	0.80	0.19	30,34,42,43	0
2	CSU	B	110	10/11	0.90	0.11	23,24,28,28	0
2	CSU	F	110	10/11	0.93	0.10	24,25,27,27	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	EDO	A	606	4/4	0.80	0.14	21,22,22,24	0
3	EDO	Y	601	4/4	0.81	0.17	19,20,20,20	0
3	EDO	D	603	4/4	0.81	0.12	23,23,25,27	0
3	EDO	D	604	4/4	0.83	0.13	25,27,27,27	0
3	EDO	B	602	4/4	0.84	0.14	23,25,25,27	0
3	EDO	Y	605	4/4	0.88	0.13	23,25,26,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.