



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 2, 2026 – 12:53 AM UTC

PDB ID : 8OXF / pdb_00008oxf
Title : Enantioselective synthesis of fluorine-containing building blocks by employing mutant HHDH in a two-phase system
Authors : Majeric Elenkov, M.; Stefanic, Z.
Deposited on : 2023-05-02
Resolution : 1.77 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1365 (1.78-1.78)
Clashscore	190562	1395 (1.78-1.78)
Ramachandran outliers	187476	1382 (1.78-1.78)
Sidechain outliers	187428	1382 (1.78-1.78)
RSRZ outliers	180081	1365 (1.78-1.78)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	243	 91% 9%
1	B	243	 88% 12%
1	C	243	 93% 7%
1	D	243	 88% 12%
1	E	243	 89% 10%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	243	2% 91% 9%
1	G	243	% 89% 10%
1	H	243	% 86% 14%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 16731 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Halohydrin dehalogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	243	Total	C	N	O	S	0	0	0
			1855	1182	322	347	4			
1	B	243	Total	C	N	O	S	0	0	0
			1855	1182	322	347	4			
1	C	243	Total	C	N	O	S	0	0	0
			1855	1182	322	347	4			
1	D	243	Total	C	N	O	S	0	0	0
			1855	1182	322	347	4			
1	E	243	Total	C	N	O	S	0	0	0
			1855	1182	322	347	4			
1	F	243	Total	C	N	O	S	0	0	0
			1855	1182	322	347	4			
1	G	243	Total	C	N	O	S	0	0	0
			1855	1182	322	347	4			
1	H	243	Total	C	N	O	S	0	0	0
			1855	1182	322	347	4			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	178	ALA	ASN	conflict	UNP Q93MS3
B	178	ALA	ASN	conflict	UNP Q93MS3
C	178	ALA	ASN	conflict	UNP Q93MS3
D	178	ALA	ASN	conflict	UNP Q93MS3
E	178	ALA	ASN	conflict	UNP Q93MS3
F	178	ALA	ASN	conflict	UNP Q93MS3
G	178	ALA	ASN	conflict	UNP Q93MS3
H	178	ALA	ASN	conflict	UNP Q93MS3

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Cl 1 1	0	0
2	B	1	Total Cl 1 1	0	0
2	C	1	Total Cl 1 1	0	0
2	D	1	Total Cl 1 1	0	0
2	E	1	Total Cl 1 1	0	0
2	F	1	Total Cl 1 1	0	0
2	G	1	Total Cl 1 1	0	0
2	H	1	Total Cl 1 1	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	229	Total O 229 229	0	0
3	B	253	Total O 253 253	0	0
3	C	240	Total O 240 240	0	0
3	D	252	Total O 252 252	0	0
3	E	247	Total O 247 247	0	0
3	F	213	Total O 213 213	0	0
3	G	213	Total O 213 213	0	0
3	H	236	Total O 236 236	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Halohydrin dehalogenase

Chain A:  91% 9%



- Molecule 1: Halohydrin dehalogenase

Chain B:  88% 12%



- Molecule 1: Halohydrin dehalogenase

Chain C:  93% 7%



- Molecule 1: Halohydrin dehalogenase

Chain D:  88% 12%




- Molecule 1: Halohydrin dehalogenase



Chain E:  89% 10%





- Molecule 1: Halohydrin dehalogenase



- Molecule 1: Halohydrin dehalogenase



- Molecule 1: Halohydrin dehalogenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	60.16Å 60.06Å 137.11Å 81.73° 80.26° 81.72°	Depositor
Resolution (Å)	56.61 – 1.77 56.61 – 1.77	Depositor EDS
% Data completeness (in resolution range)	91.2 (56.61-1.77) 91.1 (56.61-1.77)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.90 (at 1.76Å)	Xtriage
Refinement program	PHENIX 1.19	Depositor
R, R_{free}	0.197 , 0.243 0.197 , 0.243	Depositor DCC
R_{free} test set	1991 reflections (1.09%)	wwPDB-VP
Wilson B-factor (Å ²)	20.7	Xtriage
Anisotropy	0.264	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 29.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.105 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	16731	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.34	0/1899	0.56	0/2589
1	B	0.35	0/1899	0.56	0/2589
1	C	0.36	0/1899	0.55	0/2589
1	D	0.35	0/1899	0.56	0/2589
1	E	0.35	0/1899	0.56	0/2589
1	F	0.34	0/1899	0.55	0/2589
1	G	0.73	4/1899 (0.2%)	0.84	4/2589 (0.2%)
1	H	0.36	0/1899	0.54	0/2589
All	All	0.42	4/15192 (0.0%)	0.60	4/20712 (0.0%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	187	PRO	CB-CG	18.22	2.40	1.49
1	G	187	PRO	CG-CD	-17.66	0.90	1.50
1	G	187	PRO	N-CD	8.42	1.59	1.47
1	G	187	PRO	CA-CB	-7.19	1.43	1.53

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	187	PRO	CB-CG-CD	-19.40	44.01	106.10
1	G	187	PRO	N-CD-CG	-16.40	78.60	103.20
1	G	187	PRO	N-CA-CB	-14.26	88.40	103.38
1	G	187	PRO	CA-CB-CG	-10.94	83.72	104.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1855	0	1841	17	0
1	B	1855	0	1841	19	0
1	C	1855	0	1841	10	0
1	D	1855	0	1841	18	0
1	E	1855	0	1841	23	0
1	F	1855	0	1841	18	0
1	G	1855	0	1841	16	0
1	H	1855	0	1841	25	0
2	A	1	0	0	0	0
2	B	1	0	0	1	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	H	1	0	0	0	0
3	A	229	0	0	3	1
3	B	253	0	0	5	1
3	C	240	0	0	4	0
3	D	252	0	0	6	1
3	E	247	0	0	9	2
3	F	213	0	0	5	2
3	G	213	0	0	2	3
3	H	236	0	0	9	2
All	All	16731	0	14728	141	6

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 141 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:139:LYS:NZ	3:C:401:HOH:O	2.00	0.94
1:A:84:ARG:O	1:A:87:ASN:ND2	2.06	0.89
1:C:101:GLN:NE2	3:C:402:HOH:O	2.10	0.85

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:3:ILE:HD11	1:D:73:ILE:HG12	1.68	0.75
1:D:68:GLN:NE2	3:D:403:HOH:O	2.23	0.71

The worst 5 of 6 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:F:549:HOH:O	3:H:576:HOH:O[1_545]	1.98	0.22
3:B:569:HOH:O	3:D:549:HOH:O[1_455]	2.01	0.19
3:A:622:HOH:O	3:G:583:HOH:O[1_556]	2.12	0.08
3:E:494:HOH:O	3:G:462:HOH:O[1_545]	2.12	0.08
3:F:447:HOH:O	3:H:439:HOH:O[1_655]	2.15	0.05

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	241/243 (99%)	230 (95%)	11 (5%)	0	100	100
1	B	241/243 (99%)	233 (97%)	8 (3%)	0	100	100
1	C	241/243 (99%)	231 (96%)	10 (4%)	0	100	100
1	D	241/243 (99%)	233 (97%)	8 (3%)	0	100	100
1	E	241/243 (99%)	232 (96%)	9 (4%)	0	100	100
1	F	241/243 (99%)	232 (96%)	9 (4%)	0	100	100
1	G	241/243 (99%)	231 (96%)	10 (4%)	0	100	100
1	H	241/243 (99%)	231 (96%)	10 (4%)	0	100	100
All	All	1928/1944 (99%)	1853 (96%)	75 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	191/191 (100%)	191 (100%)	0	100	100
1	B	191/191 (100%)	189 (99%)	2 (1%)	68	55
1	C	191/191 (100%)	187 (98%)	4 (2%)	47	27
1	D	191/191 (100%)	189 (99%)	2 (1%)	68	55
1	E	191/191 (100%)	188 (98%)	3 (2%)	55	38
1	F	191/191 (100%)	190 (100%)	1 (0%)	81	73
1	G	191/191 (100%)	187 (98%)	4 (2%)	47	27
1	H	191/191 (100%)	188 (98%)	3 (2%)	55	38
All	All	1528/1528 (100%)	1509 (99%)	19 (1%)	63	49

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	188	THR
1	H	85	PRO
1	H	121	LEU
1	H	46	SER
1	E	21	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	48	ASN
1	F	101	GLN
1	H	181	ASN
1	F	144	ASN
1	D	68	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	243/243 (100%)	0.10	1 (0%) 88 92	15, 22, 35, 41	0
1	B	243/243 (100%)	0.04	2 (0%) 82 87	15, 21, 35, 45	0
1	C	243/243 (100%)	0.02	1 (0%) 88 92	14, 21, 33, 43	0
1	D	243/243 (100%)	0.07	3 (1%) 76 82	13, 21, 38, 57	0
1	E	243/243 (100%)	0.03	1 (0%) 88 92	14, 21, 33, 42	0
1	F	243/243 (100%)	0.16	6 (2%) 58 66	13, 22, 41, 52	0
1	G	243/243 (100%)	0.05	3 (1%) 76 82	15, 21, 33, 41	0
1	H	243/243 (100%)	0.09	2 (0%) 82 87	14, 21, 33, 44	0
All	All	1944/1944 (100%)	0.07	19 (0%) 79 85	13, 21, 35, 57	0

The worst 5 of 19 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	143	TYR	3.2
1	C	143	TYR	3.1
1	F	190	ASP	2.9
1	F	191	TRP	2.7
1	E	143	TYR	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CL	C	301	1/1	0.97	0.05	29,29,29,29	0
2	CL	A	301	1/1	0.98	0.06	28,28,28,28	0
2	CL	E	301	1/1	0.98	0.06	26,26,26,26	0
2	CL	D	301	1/1	0.99	0.04	23,23,23,23	0
2	CL	B	301	1/1	0.99	0.03	22,22,22,22	0
2	CL	F	301	1/1	0.99	0.03	22,22,22,22	0
2	CL	G	301	1/1	0.99	0.05	24,24,24,24	0
2	CL	H	301	1/1	0.99	0.03	19,19,19,19	0

6.5 Other polymers [i](#)

There are no such residues in this entry.