



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 08:12 PM UTC

PDB ID : 8OXG / pdb_00008oxg
Title : Crystal structure of human methionine aminopeptidase-2 complexed with (3R,4S,5S,6R)-5-methoxy-4-[(2R,3R)-2-methyl-3-(3-methyl-2-buten-1-yl)-2-oxiran-1-yl]-1-oxaspiro[2.5]oct-6-yl N-(trans-4-aminocyclohexyl)carbamate
Authors : Moss, S.; Cornelius, P.
Deposited on : 2023-05-02
Resolution : 1.73 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.73 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1187 (1.74-1.74)
Clashscore	190562	1207 (1.74-1.74)
Ramachandran outliers	187476	1200 (1.74-1.74)
Sidechain outliers	187428	1200 (1.74-1.74)
RSRZ outliers	180081	1188 (1.74-1.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	384	4% 88% 7% • •
1	B	384	3% 86% 8% • •

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 12226 atoms, of which 5793 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Methionine aminopeptidase 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	368	Total	C	H	N	O	S	178	3	0
			5763	1824	2859	498	556	26			
1	B	369	Total	C	H	N	O	S	177	1	0
			5751	1820	2854	498	555	24			

There are 26 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	95	MET	-	initiating methionine	UNP P50579
A	96	HIS	-	expression tag	UNP P50579
A	97	HIS	-	expression tag	UNP P50579
A	98	HIS	-	expression tag	UNP P50579
A	99	HIS	-	expression tag	UNP P50579
A	100	HIS	-	expression tag	UNP P50579
A	101	HIS	-	expression tag	UNP P50579
A	102	GLU	-	expression tag	UNP P50579
A	103	ASN	-	expression tag	UNP P50579
A	104	LEU	-	expression tag	UNP P50579
A	105	TYR	-	expression tag	UNP P50579
A	106	PHE	-	expression tag	UNP P50579
A	107	GLN	-	expression tag	UNP P50579
B	95	MET	-	initiating methionine	UNP P50579
B	96	HIS	-	expression tag	UNP P50579
B	97	HIS	-	expression tag	UNP P50579
B	98	HIS	-	expression tag	UNP P50579
B	99	HIS	-	expression tag	UNP P50579
B	100	HIS	-	expression tag	UNP P50579
B	101	HIS	-	expression tag	UNP P50579
B	102	GLU	-	expression tag	UNP P50579
B	103	ASN	-	expression tag	UNP P50579
B	104	LEU	-	expression tag	UNP P50579
B	105	TYR	-	expression tag	UNP P50579
B	106	PHE	-	expression tag	UNP P50579

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	107	GLN	-	expression tag	UNP P50579

- Molecule 2 is [(1 {R},2 {S},3 {S},4 {R})-2-methoxy-4-methyl-3-[(2 {R},3 {S})-2-methyl-3-(3-methylbut-2-enyl)oxiran-2-yl]-4-oxidanyl-cyclohexyl] {N}-(4-azanylcyclohexyl)carbamate (CCD ID: W5N) (formula: C₂₃H₄₀N₂O₅) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	1	0
			70	23	40	2	5		
2	B	1	Total	C	H	N	O	1	0
			70	23	40	2	5		

- Molecule 3 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Mn	0	0
			2	2		
3	B	2	Total	Mn	0	0
			2	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	298	Total	O	0	0
			298	298		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	270	Total	O	0	0
			270	270		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Methionine aminopeptidase 2



• Molecule 1: Methionine aminopeptidase 2



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	119.05Å 101.13Å 83.74Å 90.00° 98.75° 90.00°	Depositor
Resolution (Å)	44.84 – 1.73 44.84 – 1.73	Depositor EDS
% Data completeness (in resolution range)	99.8 (44.84-1.73) 99.9 (44.84-1.73)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.68 (at 1.73Å)	Xtriage
Refinement program	REFMAC 5.8.0352, REFMAC 5.8.0352	Depositor
R, R_{free}	0.181 , 0.215 0.182 , 0.216	Depositor DCC
R_{free} test set	5131 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	27.7	Xtriage
Anisotropy	0.091	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 34.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	12226	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MN, W5N

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.99	5/2969 (0.2%)	1.30	4/4020 (0.1%)
1	B	1.04	6/2962 (0.2%)	1.29	4/4011 (0.1%)
All	All	1.02	11/5931 (0.2%)	1.29	8/8031 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	5
All	All	0	7

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	255	HIS	CE1-NE2	9.04	1.41	1.32
1	B	339	HIS	CE1-NE2	8.53	1.41	1.32
1	A	339	HIS	CE1-NE2	7.92	1.40	1.32
1	B	375	HIS	CE1-NE2	7.32	1.39	1.32
1	B	231	HIS	CE1-NE2	6.68	1.39	1.32
1	A	460	HIS	CE1-NE2	6.64	1.39	1.32
1	A	255	HIS	CE1-NE2	6.42	1.39	1.32
1	A	176	HIS	CE1-NE2	6.11	1.38	1.32
1	B	176	HIS	CE1-NE2	5.98	1.38	1.32
1	B	162	SER	CA-CB	-5.83	1.45	1.53
1	A	379	GLU	CD-OE2	5.42	1.35	1.25

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	326	ARG	NE-CZ-NH2	-7.95	112.05	119.20
1	B	282	ASP	CA-CB-CG	6.39	118.99	112.60
1	B	322	VAL	N-CA-CB	6.07	117.15	110.53
1	B	231	HIS	N-CA-C	5.87	120.03	112.86
1	B	262	ASP	CA-CB-CG	5.26	117.86	112.60
1	A	142	ASP	CA-CB-CG	5.20	117.80	112.60
1	A	150	THR	CA-CB-OG1	-5.15	101.87	109.60
1	A	270	ASN	CB-CA-C	5.03	116.62	109.38

There are no chirality outliers.

All (7) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	144	ARG	Sidechain
1	A	354	ARG	Sidechain
1	B	144	ARG	Sidechain
1	B	296	ARG	Sidechain
1	B	354	ARG	Sidechain
1	B	398	ARG	Sidechain
1	B	465	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2904	2859	2846	14	0
1	B	2897	2854	2843	16	0
2	A	30	40	0	0	0
2	B	30	40	0	2	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	298	0	0	7	0
4	B	270	0	0	3	0
All	All	6433	5793	5689	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:206:LYS:NZ	4:A:601:HOH:O	2.07	0.75
1:B:231:HIS:CE1	2:B:501:W5N:C13	2.73	0.70
1:A:144:ARG:NH2	4:A:606:HOH:O	2.28	0.65
1:A:210:GLU:OE1	4:A:602:HOH:O	2.16	0.61
1:B:391:HIS:O	1:B:391:HIS:ND1	2.33	0.55
1:B:145:THR:HG23	4:B:790:HOH:O	2.08	0.53
1:B:144:ARG:NH2	4:B:601:HOH:O	2.32	0.52
1:A:245:ASP:OD1	1:A:270:ASN:ND2	2.32	0.51
4:A:798:HOH:O	1:B:309:GLU:HG3	2.10	0.51
1:A:394:ILE:HB	1:A:400:LYS:HD2	1.93	0.51
1:A:139:PRO:HA	1:A:144:ARG:O	2.14	0.48
1:B:141:GLN:HE22	1:B:296:ARG:NH1	2.12	0.47
1:A:314:GLU:HG2	4:A:761:HOH:O	2.15	0.47
1:B:389:VAL:CG1	1:B:392:VAL:HG21	2.46	0.46
1:A:394:ILE:HB	1:A:400:LYS:CD	2.46	0.46
1:B:183:VAL:HG21	1:B:200:LEU:HD12	1.98	0.45
1:A:321:GLN:HA	1:A:321:GLN:OE1	2.17	0.44
1:B:398:ARG:HG3	4:B:796:HOH:O	2.17	0.44
1:B:400:LYS:HE2	1:B:400:LYS:HB2	1.78	0.44
1:B:389:VAL:HG21	1:B:439:GLY:O	2.19	0.43
1:B:450:ILE:O	1:B:451:LYS:C	2.62	0.43
1:A:111:VAL:HA	4:A:654:HOH:O	2.19	0.42
1:B:389:VAL:HG13	1:B:392:VAL:HG21	2.02	0.42
1:B:231:HIS:CD2	2:B:501:W5N:C13	2.84	0.42
1:A:164:GLU:HG2	1:B:371:LYS:HE2	2.00	0.41
1:A:436:CYS:HA	1:A:441:VAL:O	2.21	0.41
1:B:129:PHE:CD2	1:B:171:GLU:HB2	2.56	0.40
1:A:130:PRO:HD2	4:A:633:HOH:O	2.20	0.40
1:A:399:THR:HG21	1:A:434:ASN:ND2	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	369/384 (96%)	360 (98%)	9 (2%)	0	100	100
1	B	368/384 (96%)	354 (96%)	14 (4%)	0	100	100
All	All	737/768 (96%)	714 (97%)	23 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	319/331 (96%)	315 (99%)	4 (1%)	61	43
1	B	318/331 (96%)	308 (97%)	10 (3%)	35	12
All	All	637/662 (96%)	623 (98%)	14 (2%)	45	22

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	145	THR
1	A	289	LYS
1	A	389	VAL
1	A	391	HIS
1	B	111	VAL
1	B	153	GLU
1	B	164	GLU
1	B	207	LEU
1	B	209	LYS
1	B	318	LYS
1	B	371	LYS
1	B	387	PHE
1	B	443	PRO
1	B	451	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	434	ASN
1	B	141	GLN
1	B	211	ASN
1	B	214	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	W5N	A	501	1	32,32,32	0.94	1 (3%)	38,48,48	3.64	14 (36%)
2	W5N	B	501	1	32,32,32	0.89	1 (3%)	38,48,48	2.66	5 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	W5N	A	501	1	-	3/18/59/59	0/3/3/3
2	W5N	B	501	1	-	2/18/59/59	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	W5N	C4-C3	2.82	1.57	1.53
2	A	501	W5N	O2-C4	2.03	1.48	1.46

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	W5N	C5-C4-C6	-14.11	89.06	121.56
2	A	501	W5N	C5-C4-C6	-13.73	89.93	121.56
2	A	501	W5N	C18-N1-C17	8.59	135.45	122.23
2	A	501	W5N	O4-C17-N1	-7.39	100.83	110.26
2	A	501	W5N	C22-C21-C20	6.67	117.07	110.29
2	A	501	W5N	C23-C18-C19	5.27	119.90	110.80
2	A	501	W5N	O5-C17-N1	4.45	132.16	124.86
2	A	501	W5N	C23-C18-N1	4.39	119.35	110.57
2	A	501	W5N	C19-C20-C21	3.66	115.73	111.53
2	B	501	W5N	O3-C12-C13	-3.39	100.73	108.08
2	A	501	W5N	C19-C18-N1	3.34	117.25	110.57
2	A	501	W5N	C22-C23-C18	3.18	114.98	111.49
2	B	501	W5N	O2-C4-C6	-3.13	57.41	59.39
2	B	501	W5N	O2-C4-C5	2.89	118.19	114.12
2	A	501	W5N	O2-C4-C5	2.68	117.90	114.12
2	A	501	W5N	C23-C22-C21	2.53	114.43	111.53
2	B	501	W5N	C13-C12-C14	2.21	113.95	111.02
2	A	501	W5N	O2-C6-C4	2.21	61.58	59.81
2	A	501	W5N	O2-C4-C6	-2.03	58.11	59.39

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	501	W5N	C4-C6-C7-C8
2	B	501	W5N	C4-C6-C7-C8
2	B	501	W5N	O2-C6-C7-C8
2	A	501	W5N	C16-C2-O1-C1
2	A	501	W5N	C7-C8-C9-C11


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	501	W5N	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	368/384 (95%)	-0.11	16 (4%)	40 49	11, 26, 50, 105	10 (2%)
1	B	369/384 (96%)	-0.03	13 (3%)	47 56	15, 28, 51, 101	8 (2%)
All	All	737/768 (95%)	-0.07	29 (3%)	43 52	11, 27, 51, 105	18 (2%)

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	389	VAL	5.6
1	A	389	VAL	4.4
1	A	145	THR	4.3
1	A	390	GLY	3.8
1	A	184[A]	MET	3.4
1	A	148	TRP	3.2
1	B	392	VAL	3.1
1	A	391	HIS	3.1
1	A	146	ALA	3.1
1	A	125	PRO	3.0
1	B	391	HIS	2.9
1	A	388	ASP	2.8
1	B	111	VAL	2.7
1	B	379	GLU	2.6
1	B	393	PRO	2.6
1	B	390	GLY	2.6
1	A	156	LYS	2.6
1	B	387	PHE	2.6
1	A	111	VAL	2.5
1	B	156	LYS	2.5
1	A	147	ALA	2.4
1	A	387	PHE	2.3
1	B	375	HIS	2.2
1	A	144	ARG	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	110	LYS	2.2
1	B	350	GLY	2.1
1	A	430[A]	MET	2.1
1	B	141	GLN	2.1
1	A	127	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	W5N	A	501	30/30	0.96	0.06	16,21,30,31	1
2	W5N	B	501	30/30	0.96	0.06	18,27,35,41	1
3	MN	B	503	1/1	0.99	0.03	21,21,21,21	0
3	MN	A	503	1/1	1.00	0.03	19,19,19,19	0
3	MN	B	502	1/1	1.00	0.01	19,19,19,19	0
3	MN	A	502	1/1	1.00	0.01	18,18,18,18	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around W5N A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around W5N B 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.