



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 11:03 AM UTC

PDB ID : 2OYQ / pdb_00002oyq
Title : Crystal structure of RB69 gp43 in complex with DNA with 5-NIMP opposite an abasic site analog
Authors : Zahn, K.E.; Belrhali, H.; Wallace, S.S.; Doublié, S.
Deposited on : 2007-02-22
Resolution : 2.86 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1407 (2.88-2.84)
Clashscore	190562	1446 (2.88-2.84)
Ramachandran outliers	187476	1406 (2.88-2.84)
Sidechain outliers	187428	1407 (2.88-2.84)
RSRZ outliers	180081	1408 (2.88-2.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	21	
1	G	21	
1	I	21	
1	K	21	
2	F	15	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	H	15	7% 40% 53% 7%
2	J	15	47% 47% 7%
2	L	15	20% 67% 13%
3	A	903	% 73% 20% • 6%
3	B	903	2% 61% 21% • 16%
3	C	903	2% 76% 20% • •
3	D	903	10% 73% 15% • 11%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 28889 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called Template DNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	E	16	Total	C	N	O	P	0	0	0
			314	150	56	93	15			
1	G	11	Total	C	N	O	P	0	0	0
			223	106	44	63	10			
1	I	20	Total	C	N	O	P	0	0	0
			395	188	72	116	19			
1	K	12	Total	C	N	O	P	0	0	0
			244	116	49	68	11			

- Molecule 2 is a DNA chain called Primer DNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	F	15	Total	C	N	O	P	0	0	0
			310	150	57	89	14			
2	H	14	Total	C	N	O	P	0	0	0
			287	137	55	82	13			
2	J	15	Total	C	N	O	P	0	0	0
			310	150	57	89	14			
2	L	13	Total	C	N	O	P	0	0	0
			265	127	50	76	12			

- Molecule 3 is a protein called DNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	848	Total	C	N	O	S	0	0	0
			6877	4423	1143	1280	31			
3	B	756	Total	C	N	O	S	0	0	0
			6045	3883	1001	1134	27			
3	C	891	Total	C	N	O	S	0	0	0
			7150	4583	1182	1352	33			
3	D	807	Total	C	N	O	S	0	0	0
			5688	3586	953	1125	24			

Chain	Residue	Modelled	Actual	Comment	Reference
A	222	ALA	ASP	engineered mutation	UNP Q38087
A	327	ALA	ASP	engineered mutation	UNP Q38087
B	222	ALA	ASP	engineered mutation	UNP Q38087
B	327	ALA	ASP	engineered mutation	UNP Q38087
C	222	ALA	ASP	engineered mutation	UNP Q38087
C	327	ALA	ASP	engineered mutation	UNP Q38087
D	222	ALA	ASP	engineered mutation	UNP Q38087
D	327	ALA	ASP	engineered mutation	UNP Q38087

- | Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|-----|-------|----------|-----------------|---------|---------|
| 4 | I | 1 | Total Mg
1 1 | 0 | 0 |
| 4 | K | 1 | Total Mg
1 1 | 0 | 0 |

- # N5P

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	I	1	Total 32	C 13	N 2	O 14	P 3	0	0
5	A	1	Total 32	C 13	N 2	O 14	P 3	0	0



WORLD WIDE
PDB
PROTEIN DATA BANK

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			32	13	2	14	3		
5	C	1	Total	C	N	O	P	0	0
			32	13	2	14	3		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	E	5	Total	O	0	0
			5	5		
6	F	8	Total	O	0	0
			8	8		
6	G	2	Total	O	0	0
			2	2		
6	H	3	Total	O	0	0
			3	3		
6	I	22	Total	O	0	0
			22	22		
6	J	19	Total	O	0	0
			19	19		
6	K	3	Total	O	0	0
			3	3		
6	L	3	Total	O	0	0
			3	3		
6	A	235	Total	O	0	0
			235	235		
6	B	125	Total	O	0	0
			125	125		
6	C	185	Total	O	0	0
			185	185		
6	D	41	Total	O	0	0
			41	41		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Template DNA




- Molecule 1: Template DNA



- Molecule 1: Template DNA



- Molecule 1: Template DNA



- Molecule 2: Primer DNA



- Molecule 2: Primer DNA







• Molecule 2: Primer DNA



• Molecule 2: Primer DNA




• Molecule 3: DNA polymerase





• Molecule 3: DNA polymerase





• Molecule 3: DNA polymerase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	132.59Å 123.48Å 164.33Å 90.00° 96.22° 90.00°	Depositor
Resolution (Å)	30.00 – 2.86 30.00 – 2.86	Depositor EDS
% Data completeness (in resolution range)	91.2 (30.00-2.86) 91.1 (30.00-2.86)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.85 (at 2.86Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.228 , 0.294 0.227 , 0.287	Depositor DCC
R_{free} test set	23125 reflections (9.68%)	wwPDB-VP
Wilson B-factor (Å ²)	62.5	Xtriage
Anisotropy	0.210	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 41.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	28889	wwPDB-VP
Average B, all atoms (Å ²)	78.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: N5P, MG, N5I, 3DR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	E	0.19	0/342	0.53	0/525
1	G	0.28	0/250	0.76	0/384
1	I	0.32	0/429	0.98	0/657
1	K	0.25	0/274	0.79	0/421
2	F	0.26	0/322	0.86	0/496
2	H	0.28	0/322	0.87	0/496
2	J	0.27	0/322	0.88	0/496
2	L	0.27	0/297	0.85	0/457
3	A	0.51	0/7049	0.84	1/9536 (0.0%)
3	B	0.49	0/6193	0.83	0/8393
3	C	0.49	0/7324	0.82	2/9912 (0.0%)
3	D	0.47	0/5812	0.84	4/7955 (0.1%)
All	All	0.48	0/28936	0.83	7/39728 (0.0%)

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	737	THR	CA-C-N	5.55	125.56	119.90
3	D	737	THR	C-N-CA	5.55	125.56	119.90
3	D	125	GLU	CA-C-N	5.53	126.75	119.84
3	D	125	GLU	C-N-CA	5.53	126.75	119.84
3	C	449	ARG	CA-C-N	5.36	125.43	119.32

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	314	0	177	19	0
1	G	223	0	124	2	0
1	I	395	0	222	4	0
1	K	244	0	135	3	0
2	F	310	0	171	6	0
2	H	287	0	159	11	0
2	J	310	0	171	13	0
2	L	265	0	148	6	0
3	A	6877	0	6731	123	0
3	B	6045	0	5799	130	0
3	C	7150	0	6884	124	0
3	D	5688	0	4707	68	0
4	I	1	0	0	0	0
4	K	1	0	0	0	0
5	A	64	0	26	1	0
5	C	32	0	13	0	0
5	I	32	0	13	0	0
6	A	235	0	0	4	0
6	B	125	0	0	7	0
6	C	185	0	0	3	0
6	D	41	0	0	1	0
6	E	5	0	0	1	0
6	F	8	0	0	1	0
6	G	2	0	0	0	0
6	H	3	0	0	1	0
6	I	22	0	0	1	0
6	J	19	0	0	0	0
6	K	3	0	0	0	0
6	L	3	0	0	0	0
All	All	28889	0	25480	494	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 494 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:13:DA:H2''	1:E:14:DC:H5''	1.29	1.07
3:B:442:TYR:HB3	3:B:592:MET:HE2	1.36	1.06
1:E:9:DT:H71	6:E:210:HOH:O	1.59	1.00
3:A:356:GLN:H	3:A:356:GLN:HE21	1.02	0.98
3:B:597:ILE:HG21	3:B:683:MET:HE1	1.48	0.95

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	844/903 (94%)	794 (94%)	45 (5%)	5 (1%)	21	38
3	B	750/903 (83%)	697 (93%)	49 (6%)	4 (0%)	24	42
3	C	885/903 (98%)	817 (92%)	61 (7%)	7 (1%)	16	31
3	D	797/903 (88%)	701 (88%)	83 (10%)	13 (2%)	7	17
All	All	3276/3612 (91%)	3009 (92%)	238 (7%)	29 (1%)	14	28

5 of 29 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	B	622	THR
3	B	736	SER
3	C	622	THR
3	D	622	THR
3	A	549	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	738/800 (92%)	702 (95%)	36 (5%)	22	45
3	B	634/800 (79%)	598 (94%)	36 (6%)	18	39
3	C	755/800 (94%)	706 (94%)	49 (6%)	15	32
3	D	483/800 (60%)	454 (94%)	29 (6%)	17	36
All	All	2610/3200 (82%)	2460 (94%)	150 (6%)	18	39

5 of 150 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	815	ILE
3	D	633	ILE
3	C	897	LEU
3	D	192	ASP
3	B	181	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 69 such sidechains are listed below:

Mol	Chain	Res	Type
3	D	371	ASN
3	D	402	ASN
3	D	742	GLN
3	B	440	HIS
3	B	371	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	3DR	I	6	1	8,11,12	0.47	0	7,14,17	0.81	0
2	N5I	J	115	2	22,25,26	0.84	0	29,36,39	0.99	1 (3%)
2	N5I	F	115	2	22,25,26	0.87	0	29,36,39	0.94	2 (6%)
1	3DR	E	6	1	8,8,12	0.43	0	9,10,17	0.81	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	3DR	I	6	1	-	2/3/15/16	0/1/1/1
2	N5I	J	115	2	-	0/9/25/26	0/3/3/3
2	N5I	F	115	2	-	0/9/25/26	0/3/3/3
1	3DR	E	6	1	-	2/2/12/16	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	F	115	N5I	CE2-NE1-CD1	2.21	109.49	107.77
2	F	115	N5I	CD2-CE3-CZ3	-2.07	118.06	120.47
2	J	115	N5I	CD2-CE3-CZ3	-2.03	118.12	120.47

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	E	6	3DR	C3'-C4'-C5'-O5'
1	E	6	3DR	O4'-C4'-C5'-O5'
1	I	6	3DR	O4'-C4'-C5'-O5'
1	I	6	3DR	C3'-C4'-C5'-O5'

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	J	115	N5I	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	N5P	A	905	4	33,34,34	1.00	3 (9%)	47,53,53	1.00	2 (4%)
5	N5P	A	904	-	33,34,34	0.97	2 (6%)	47,53,53	0.95	2 (4%)
5	N5P	I	704	4	33,34,34	1.03	3 (9%)	47,53,53	0.93	2 (4%)
5	N5P	C	904	-	33,34,34	1.01	3 (9%)	47,53,53	0.94	1 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	N5P	A	905	4	-	4/24/38/38	0/3/3/3
5	N5P	A	904	-	-	4/24/38/38	0/3/3/3
5	N5P	I	704	4	-	4/24/38/38	0/3/3/3
5	N5P	C	904	-	-	1/24/38/38	0/3/3/3

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	904	N5P	PB-O3B	2.55	1.62	1.59
5	I	704	N5P	PA-O3A	2.45	1.62	1.59

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	I	704	N5P	PB-O3A	2.42	1.62	1.59
5	C	904	N5P	PB-O3A	2.33	1.62	1.59
5	A	905	N5P	PA-O3A	2.33	1.62	1.59

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	I	704	N5P	CE2-NE1-CD1	2.24	109.52	107.77
5	A	904	N5P	CE2-NE1-CD1	2.22	109.51	107.77
5	A	905	N5P	CD2-CE3-CZ3	-2.19	117.93	120.47
5	A	905	N5P	CE2-NE1-CD1	2.18	109.47	107.77
5	I	704	N5P	CD2-CE3-CZ3	-2.12	118.02	120.47

There are no chirality outliers.

5 of 13 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	904	N5P	O4'-C4'-C5'-O5'
5	A	904	N5P	C3'-C4'-C5'-O5'
5	I	704	N5P	PA-O3A-PB-O1B
5	A	905	N5P	PA-O3A-PB-O1B
5	A	905	N5P	PB-O3A-PA-O1A


There are no ring outliers.

1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	905	N5P	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand N5P A 905



Ligand N5P A 904





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	E	15/21 (71%)	0.25	0 100 100	58, 78, 115, 119	0
1	G	11/21 (52%)	1.48	3 (27%) 1 1	62, 115, 126, 132	0
1	I	19/21 (90%)	-0.19	1 (5%) 32 24	43, 56, 138, 144	0
1	K	12/21 (57%)	-0.04	0 100 100	42, 94, 107, 115	0
2	F	14/15 (93%)	0.22	0 100 100	73, 106, 124, 126	0
2	H	14/15 (93%)	1.29	1 (7%) 22 16	102, 129, 134, 137	0
2	J	14/15 (93%)	-0.35	0 100 100	44, 68, 85, 88	0
2	L	13/15 (86%)	0.55	0 100 100	85, 108, 113, 113	0
3	A	848/903 (93%)	-0.11	8 (0%) 81 76	36, 53, 79, 96	0
3	B	756/903 (83%)	0.22	19 (2%) 58 49	44, 68, 105, 116	0
3	C	891/903 (98%)	0.17	22 (2%) 58 49	36, 63, 110, 135	0
3	D	807/903 (89%)	0.94	86 (10%) 11 8	105, 121, 145, 147	0
All	All	3414/3756 (90%)	0.30	140 (4%) 41 32	36, 69, 138, 147	0

The worst 5 of 140 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	D	6	LEU	5.0
3	D	782	VAL	4.6
3	C	252	VAL	4.2
3	D	393	GLY	4.1
3	A	558	ASN	4.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	3DR	E	6	8/12	0.43	0.16	123,124,124,124	0
2	N5I	F	115	23/24	0.76	0.16	127,128,128,128	0
2	N5I	J	115	23/24	0.88	0.12	83,89,89,89	0
1	3DR	I	6	11/12	0.90	0.11	104,108,115,115	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MG	K	801	1/1	0.51	0.27	67,67,67,67	1
5	N5P	C	904	32/32	0.78	0.16	79,81,83,83	32
4	MG	I	802	1/1	0.85	0.29	82,82,82,82	1
5	N5P	I	704	32/32	0.86	0.18	78,80,84,84	32
5	N5P	A	904	32/32	0.88	0.13	74,76,88,88	32
5	N5P	A	905	32/32	0.91	0.13	61,65,75,76	32




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around N5P C 904:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around N5P I 704:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around N5P A 904:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around N5P A 905:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.