



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 05:33 PM UTC

PDB ID : 3OY3 / pdb_00003oy3
Title : Crystal structure of ABL T315I mutant kinase domain bound with a DFG-out inhibitor AP24589
Authors : Zhou, T.; Commodore, L.; Huang, W.S.; Wang, Y.; Thomas, M.; Keats, J.; Xu, Q.; Rivera, V.; Shakespeare, W.C.; Clackson, T.; Dalgarno, D.C.; Zhu, X.
Deposited on : 2010-09-22
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	284	6% 85% 12% .
1	B	284	8% 76% 20% .

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5073 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tyrosine-protein kinase ABL1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	284	Total	C	N	O	S	0	0	0
			2300	1477	372	434	17			
1	B	284	Total	C	N	O	S	0	0	0
			2300	1477	372	434	17			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	228	GLY	-	expression tag	UNP P00520
A	315	ILE	THR	engineered mutation	UNP P00520
B	228	GLY	-	expression tag	UNP P00520
B	315	ILE	THR	engineered mutation	UNP P00520

- Molecule 2 is 5-[(5-{[4-{[4-(2-hydroxyethyl)piperazin-1-yl]methyl}-3-(trifluoromethyl)phenyl]carbonyl}-2-methylphenyl)ethynyl]-1-methyl-1H-imidazole-2-carboxamide (CCD ID: XY3) (formula: C₂₉H₃₁F₃N₆O₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	0	0
			41	29	3	6	3		
2	B	1	Total	C	F	N	O	0	0
			41	29	3	6	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	238	Total	O	0	0
			238	238		
3	B	153	Total	O	0	0
			153	153		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Tyrosine-protein kinase ABL1



• Molecule 1: Tyrosine-protein kinase ABL1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	66.47Å 59.99Å 89.44Å 90.00° 97.97° 90.00°	Depositor
Resolution (Å)	50.00 – 1.95 50.00 – 1.95	Depositor EDS
% Data completeness (in resolution range)	90.6 (50.00-1.95) 90.6 (50.00-1.95)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.59 (at 1.93Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.220 , 0.266 0.210 , 0.254	Depositor DCC
R_{free} test set	2532 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	18.7	Xtriage
Anisotropy	0.740	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 48.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5073	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 79.55 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.4421e-07. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: XY3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.78	0/2358	0.91	3/3190 (0.1%)
1	B	0.65	0/2358	0.85	2/3190 (0.1%)
All	All	0.71	0/4716	0.88	5/6380 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	273	LEU	N-CA-C	-6.53	104.00	112.68
1	A	438	SER	CA-C-N	5.73	126.05	119.92
1	A	438	SER	C-N-CA	5.73	126.05	119.92
1	B	229	SER	CA-C-N	5.21	125.51	119.47
1	B	229	SER	C-N-CA	5.21	125.51	119.47

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2300	0	2239	38	0
1	B	2300	0	2239	49	0
2	A	41	0	25	1	0
2	B	41	0	25	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	238	0	0	3	0
3	B	153	0	0	1	0
All	All	5073	0	4528	86	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (86) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:388:MET:HE2	1:B:392:THR:HG21	1.30	1.06
1:B:494:GLU:O	1:B:498:GLN:HG2	1.74	0.86
1:A:275:GLU:HB2	1:A:276:ASP:CA	2.07	0.84
1:A:275:GLU:HB2	1:A:276:ASP:CG	2.04	0.83
1:B:273:LEU:HD11	1:B:279:GLU:HB2	1.60	0.82
1:A:415:LYS:HE2	3:A:68:HOH:O	1.79	0.80
1:A:275:GLU:HB2	1:A:276:ASP:HA	1.66	0.77
1:A:242:ILE:HD11	1:A:314:ILE:HD13	1.68	0.76
1:B:277:THR:HG22	1:B:280:VAL:HG21	1.69	0.74
1:B:280:VAL:HG13	1:B:311:PHE:HE1	1.52	0.74
1:B:273:LEU:O	1:B:274:LYS:HB2	1.88	0.73
1:B:339:VAL:O	1:B:343:MET:HG3	1.89	0.72
1:A:275:GLU:HB2	1:A:276:ASP:CB	2.20	0.71
1:B:471:LEU:HD22	1:B:493:PHE:CE2	2.26	0.70
1:B:471:LEU:HD22	1:B:493:PHE:HE2	1.56	0.70
1:B:307:ARG:C	1:B:308:GLU:HG3	2.17	0.70
1:B:465:PRO:HG2	1:B:468:VAL:CG2	2.22	0.69
1:B:360:ILE:HD13	1:B:417:SER:HA	1.74	0.69
1:B:252:GLN:OE1	1:B:389:THR:HG22	1.94	0.68
1:B:255:GLU:HB3	1:B:272:THR:HG22	1.77	0.67
1:A:273:LEU:HD11	1:A:279:GLU:HB2	1.76	0.66
1:A:274:LYS:HA	1:A:310:PRO:HB3	1.79	0.64
1:A:280:VAL:HG13	1:A:311:PHE:HE1	1.62	0.64
1:B:277:THR:HG22	1:B:280:VAL:CG2	2.29	0.61
1:A:298:LEU:HD23	1:A:379:VAL:HB	1.82	0.61
1:A:384:LEU:HB3	1:A:388:MET:HG3	1.83	0.61
1:A:275:GLU:CB	1:A:276:ASP:HA	2.31	0.61
1:B:388:MET:CE	1:B:392:THR:HG21	2.20	0.60
1:A:273:LEU:HD11	1:A:279:GLU:CB	2.34	0.58
1:B:277:THR:O	1:B:278:MET:HB2	2.03	0.57
1:B:242:ILE:HD11	1:B:314:ILE:HD13	1.87	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:252:GLN:HE22	1:B:389:THR:H	1.53	0.57
1:B:465:PRO:HG2	1:B:468:VAL:HG23	1.88	0.56
1:A:286:GLU:HG3	2:A:1:XY3:H14	1.88	0.55
1:B:403:ILE:HB	1:B:445:LEU:HD22	1.89	0.55
1:A:239:ARG:HG2	1:A:312:TYR:CE1	2.42	0.54
1:B:492:ALA:O	1:B:496:MET:HG3	2.08	0.54
1:B:273:LEU:O	1:B:274:LYS:CB	2.55	0.53
1:A:307:ARG:C	1:A:308:GLU:HG3	2.34	0.53
1:A:277:THR:O	1:A:278:MET:HB2	2.09	0.53
1:B:239:ARG:HG3	1:B:312:TYR:CE1	2.44	0.52
1:B:360:ILE:CD1	1:B:417:SER:HA	2.39	0.52
1:B:465:PRO:HG2	1:B:468:VAL:HG21	1.91	0.52
1:A:385:SER:HA	1:A:393:TYR:CD2	2.45	0.52
1:A:445:LEU:HD12	1:A:445:LEU:N	2.24	0.51
1:A:272:THR:HG22	1:A:312:TYR:CD2	2.47	0.50
1:B:295:HIS:CG	1:B:296:PRO:HD2	2.46	0.50
1:A:264:TYR:CZ	1:B:509:GLU:HG2	2.47	0.49
1:B:275:GLU:O	1:B:277:THR:N	2.46	0.49
1:B:275:GLU:HG3	1:B:309:PRO:HB2	1.94	0.48
1:A:242:ILE:CD1	1:A:314:ILE:HD13	2.41	0.48
1:A:445:LEU:H	1:A:445:LEU:CD1	2.26	0.48
1:A:274:LYS:HB2	3:A:209:HOH:O	2.12	0.48
1:B:290:MET:HE2	1:B:301:LEU:HD13	1.95	0.48
1:A:283:PHE:CE2	1:A:311:PHE:HB3	2.49	0.47
1:B:391:ASP:OD1	1:B:391:ASP:N	2.48	0.47
1:B:254:GLY:HA3	1:B:387:LEU:HD13	1.97	0.46
1:B:254:GLY:CA	1:B:387:LEU:HD13	2.46	0.46
1:B:361:HIS:O	1:B:362:ARG:HB2	2.14	0.46
1:B:468:VAL:HG22	1:B:496:MET:HE1	1.96	0.46
1:B:271:LYS:NZ	1:B:286:GLU:OE1	2.47	0.46
1:B:246:HIS:H	1:B:246:HIS:CD2	2.34	0.46
1:B:392:THR:O	1:B:402:PRO:HA	2.16	0.46
1:A:363:ASP:OD2	1:A:393:TYR:OH	2.33	0.45
1:A:386:ARG:H	1:A:386:ARG:HG2	1.38	0.45
1:B:243:THR:OG1	1:B:262:LYS:HD2	2.16	0.45
1:A:256:VAL:O	1:A:256:VAL:HG23	2.18	0.44
1:B:286:GLU:HG3	2:B:1:XY3:H14	2.00	0.44
1:A:445:LEU:HD12	1:A:445:LEU:H	1.83	0.44
1:B:263:LYS:HE2	1:B:264:TYR:CZ	2.52	0.44
1:A:340:LEU:HA	1:A:343:MET:HE2	2.00	0.43
1:A:434:THR:HB	1:A:437:MET:HG3	1.99	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:292:GLU:HB3	3:B:128:HOH:O	2.18	0.43
1:A:274:LYS:O	1:A:275:GLU:C	2.60	0.43
1:B:506:VAL:O	1:B:510:LEU:HG	2.19	0.42
1:A:242:ILE:CD1	1:A:314:ILE:CD1	2.96	0.42
1:A:275:GLU:HA	1:A:309:PRO:HB3	2.01	0.42
1:B:237:MET:SD	1:B:314:ILE:CD1	3.08	0.42
1:A:273:LEU:O	1:A:274:LYS:HB3	2.20	0.41
1:A:246:HIS:CD2	1:A:246:HIS:H	2.37	0.41
1:A:386:ARG:NH2	3:A:170:HOH:O	2.53	0.41
1:B:229:SER:HA	1:B:230:PRO:HD3	1.90	0.41
1:B:237:MET:SD	1:B:314:ILE:HD13	2.61	0.41
1:B:384:LEU:C	1:B:386:ARG:H	2.28	0.41
1:B:322:ASN:HA	1:B:369:CYS:O	2.20	0.40
1:A:445:LEU:N	1:A:445:LEU:CD1	2.82	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	282/284 (99%)	267 (95%)	12 (4%)	3 (1%)	11	4
1	B	282/284 (99%)	269 (95%)	10 (4%)	3 (1%)	11	4
All	All	564/568 (99%)	536 (95%)	22 (4%)	6 (1%)	11	4

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	276	ASP
1	B	277	THR
1	A	277	THR
1	A	278	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	274	LYS
1	A	274	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	247/248 (100%)	237 (96%)	10 (4%)	28	17
1	B	247/248 (100%)	235 (95%)	12 (5%)	22	11
All	All	494/496 (100%)	472 (96%)	22 (4%)	24	14

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	231	ASN
1	A	275	GLU
1	A	276	ASP
1	A	278	MET
1	A	308	GLU
1	A	362	ARG
1	A	386	ARG
1	A	388	MET
1	A	499	GLU
1	A	505	GLU
1	B	231	ASN
1	B	276	ASP
1	B	277	THR
1	B	278	MET
1	B	308	GLU
1	B	356	LYS
1	B	357	LYS
1	B	386	ARG
1	B	391	ASP
1	B	392	THR
1	B	471	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	473	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	246	HIS
1	A	396	HIS
1	A	490	HIS
1	A	491	GLN
1	B	246	HIS
1	B	368	ASN
1	B	396	HIS
1	B	447	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	XY3	B	1	-	43,44,44	3.05	17 (39%)	57,63,63	1.71	6 (10%)
2	XY3	A	1	-	43,44,44	3.06	18 (41%)	57,63,63	1.54	3 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	XY3	B	1	-	-	3/28/40/40	0/4/4/4
2	XY3	A	1	-	-	3/28/40/40	0/4/4/4

All (35) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	XY3	C25-C27	-10.10	1.12	1.51
2	A	1	XY3	C25-C27	-9.48	1.15	1.51
2	B	1	XY3	C81-N5	-7.56	1.11	1.33
2	A	1	XY3	C81-N5	-7.52	1.11	1.33
2	B	1	XY3	C84-N82	6.82	1.46	1.37
2	A	1	XY3	C84-N82	6.45	1.46	1.37
2	B	1	XY3	O2-C81	5.67	1.37	1.24
2	A	1	XY3	O2-C81	5.65	1.37	1.24
2	B	1	XY3	C16-C17	4.94	1.46	1.40
2	A	1	XY3	C26-N82	-4.73	1.38	1.47
2	B	1	XY3	C26-N82	-4.63	1.38	1.47
2	A	1	XY3	C8-C9	4.24	1.45	1.39
2	B	1	XY3	C8-C7	3.95	1.45	1.38
2	A	1	XY3	C18-C13	3.89	1.45	1.39
2	B	1	XY3	C18-C13	3.70	1.45	1.39
2	A	1	XY3	C15-C16	3.51	1.45	1.39
2	A	1	XY3	C8-C7	3.33	1.44	1.38
2	A	1	XY3	C16-C17	3.31	1.44	1.40
2	B	1	XY3	C10-C9	3.25	1.44	1.39
2	A	1	XY3	C18-C17	3.24	1.44	1.39
2	B	1	XY3	C15-C16	3.20	1.44	1.39
2	A	1	XY3	C14-C15	3.16	1.43	1.38
2	B	1	XY3	C8-C9	3.10	1.44	1.39
2	A	1	XY3	C3-C2	-3.01	1.38	1.42
2	A	1	XY3	O3-C27	2.67	1.55	1.42
2	A	1	XY3	C5-C6	2.51	1.44	1.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1	XY3	C10-C9	2.45	1.43	1.39
2	B	1	XY3	C14-C15	2.36	1.42	1.38
2	B	1	XY3	C19-C17	2.31	1.55	1.50
2	B	1	XY3	C10-C5	2.31	1.43	1.39
2	A	1	XY3	C20-C16	2.31	1.55	1.51
2	B	1	XY3	C18-C17	2.25	1.42	1.39
2	B	1	XY3	O3-C27	2.19	1.53	1.42
2	B	1	XY3	C5-C6	2.04	1.43	1.40
2	A	1	XY3	C20-N3	-2.01	1.43	1.47

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	XY3	C2-N82-C84	-9.11	102.78	106.37
2	A	1	XY3	C2-N82-C84	-7.93	103.24	106.37
2	B	1	XY3	C1-C2-C3	-4.29	123.67	130.61
2	A	1	XY3	C1-C2-C3	-3.98	124.18	130.61
2	B	1	XY3	C81-C84-N82	-3.82	119.87	123.65
2	A	1	XY3	C81-C84-N82	-3.43	120.26	123.65
2	B	1	XY3	C19-C17-C16	3.01	123.95	121.06
2	B	1	XY3	C2-C1-N1	-2.41	109.77	112.35
2	B	1	XY3	F1-C19-C17	-2.12	108.90	112.65

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1	XY3	C3-C4-C5-C6
2	B	1	XY3	C3-C4-C5-C10
2	B	1	XY3	C27-C25-N4-C23
2	A	1	XY3	C27-C25-N4-C23
2	B	1	XY3	C3-C4-C5-C6
2	A	1	XY3	C3-C4-C5-C10

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1	XY3	1	0
2	A	1	XY3	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	284/284 (100%)	0.13	17 (5%)	27 32	8, 19, 46, 70	0
1	B	284/284 (100%)	0.60	22 (7%)	19 22	17, 29, 50, 69	0
All	All	568/568 (100%)	0.37	39 (6%)	23 26	8, 25, 48, 70	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	228	GLY	6.0
1	A	277	THR	5.9
1	A	510	LEU	5.5
1	B	389	THR	4.8
1	B	278	MET	4.6
1	B	277	THR	4.4
1	B	275	GLU	4.2
1	A	278	MET	4.2
1	A	276	ASP	4.2
1	A	511	GLY	4.1
1	A	275	GLU	3.5
1	B	510	LEU	3.4
1	B	276	ASP	3.3
1	B	386	ARG	3.1
1	B	273	LEU	3.0
1	B	397	ALA	2.9
1	B	280	VAL	2.9
1	B	511	GLY	2.8
1	B	279	GLU	2.8
1	B	237	MET	2.7
1	B	228	GLY	2.5
1	A	386	ARG	2.5
1	B	442	GLY	2.4
1	B	443	ILE	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	358	ASN	2.3
1	A	508	LYS	2.3
1	B	394	THR	2.3
1	A	507	GLU	2.2
1	A	503	SER	2.2
1	B	399	ALA	2.1
1	A	398	GLY	2.1
1	A	506	VAL	2.1
1	B	395	ALA	2.1
1	A	279	GLU	2.0
1	A	273	LEU	2.0
1	A	387	LEU	2.0
1	B	241	ASP	2.0
1	A	274	LYS	2.0
1	B	400	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	XY3	B	1	41/41	0.94	0.07	15,22,30,37	0
2	XY3	A	1	41/41	0.96	0.06	8,13,19,28	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around XY3 B 1:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around XY3 A 1:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.