



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2026 – 08:58 PM UTC

PDB ID : 3OYR / pdb_00003oyr
Title : Crystal structure of polyprenyl synthase from *Caulobacter crescentus* CB15 complexed with calcium and isoprenyl diphosphate
Authors : Patskovsky, Y.; Toro, R.; Rutter, M.; Sauder, J.M.; Burley, S.K.; Poulter, C.D.; Gerlt, J.A.; Almo, S.C.; New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : 2010-09-23
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	345	9% 73% 10% 15%
1	B	345	% 87% 7% 5%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4996 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Trans-isoprenyl diphosphate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	292	Total	C	N	O	S	0	8	0
			2228	1410	393	417	8			
1	B	328	Total	C	N	O	S	0	8	0
			2518	1585	452	473	8			

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-9	MET	-	expression tag	UNP Q9A6I1
A	-8	ALA	-	expression tag	UNP Q9A6I1
A	-7	HIS	-	expression tag	UNP Q9A6I1
A	-6	HIS	-	expression tag	UNP Q9A6I1
A	-5	HIS	-	expression tag	UNP Q9A6I1
A	-4	HIS	-	expression tag	UNP Q9A6I1
A	-3	HIS	-	expression tag	UNP Q9A6I1
A	-2	HIS	-	expression tag	UNP Q9A6I1
A	-1	SER	-	expression tag	UNP Q9A6I1
B	-9	MET	-	expression tag	UNP Q9A6I1
B	-8	ALA	-	expression tag	UNP Q9A6I1
B	-7	HIS	-	expression tag	UNP Q9A6I1
B	-6	HIS	-	expression tag	UNP Q9A6I1
B	-5	HIS	-	expression tag	UNP Q9A6I1
B	-4	HIS	-	expression tag	UNP Q9A6I1
B	-3	HIS	-	expression tag	UNP Q9A6I1
B	-2	HIS	-	expression tag	UNP Q9A6I1
B	-1	SER	-	expression tag	UNP Q9A6I1

- Molecule 2 is PYROPHOSPHATE 2- (CCD ID: POP) (formula: $\text{H}_2\text{O}_7\text{P}_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			9	7	2		

- Molecule 3 is DIMETHYLALLYL DIPHOSPHATE (CCD ID: DMA) (formula: $C_5H_{12}O_7P_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	O	P	0	0
			14	5	7	2		
3	B	1	Total	C	O	P	0	0
			14	5	7	2		

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total 2	Ca 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	59	Total 59	O 59	0	0
5	B	152	Total 152	O 152	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Trans-isoprenyl diphosphate synthase



- Molecule 1: Trans-isoprenyl diphosphate synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	72.89Å 72.44Å 125.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 2.00 40.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	90.0 (40.00-2.00) 90.0 (40.00-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.66 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.197 , 0.247 0.197 , 0.244	Depositor DCC
R_{free} test set	1283 reflections (2.81%)	wwPDB-VP
Wilson B-factor (Å ²)	34.1	Xtriage
Anisotropy	0.496	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 51.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.018 for k,h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4996	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, POP, DMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	0/2279	0.98	3/3081 (0.1%)
1	B	0.67	0/2576	0.94	1/3478 (0.0%)
All	All	0.66	0/4855	0.96	4/6559 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	261	ARG	N-CA-C	6.10	120.19	112.87
1	B	275	GLN	N-CA-C	5.93	118.20	109.24
1	A	259	GLY	CA-C-N	5.78	127.06	119.84
1	A	259	GLY	C-N-CA	5.78	127.06	119.84

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2228	0	2257	21	0
1	B	2518	0	2560	15	0
2	A	9	0	0	0	0
3	B	28	0	18	1	0
4	B	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	59	0	0	0	0
5	B	152	0	0	0	0
All	All	4996	0	4835	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:19[A]:ARG:HH11	1:A:19[A]:ARG:CG	1.65	1.08
1:A:19[A]:ARG:HH11	1:A:19[A]:ARG:HG3	0.90	1.06
1:A:19[A]:ARG:HG3	1:A:19[A]:ARG:NH1	1.68	1.00
1:A:140:MET:HE2	1:B:135:VAL:HG22	1.63	0.79
1:A:135:VAL:HG11	1:B:144:GLU:HB2	1.68	0.76
1:A:140:MET:CE	1:B:135:VAL:HG22	2.23	0.69
1:A:33:THR:HA	1:A:36:MET:HE3	1.75	0.68
1:B:140:MET:HA	1:B:140:MET:CE	2.27	0.65
1:B:184:GLU:OE2	1:B:208:ARG:NH2	2.29	0.64
1:B:140:MET:HA	1:B:140:MET:HE2	1.83	0.60
1:B:253:LEU:O	1:B:257:ARG:HG2	2.06	0.56
1:A:141:LYS:HB3	1:A:195[B]:VAL:HG11	1.89	0.53
1:A:261:ARG:HA	1:A:264:GLU:HB2	1.92	0.51
1:A:62[B]:MET:HE3	1:A:62[B]:MET:HA	1.93	0.50
1:A:19[A]:ARG:CG	1:A:19[A]:ARG:NH1	2.39	0.49
1:A:42:ILE:HG21	1:B:162:MET:HE2	1.94	0.49
1:B:87:ILE:O	1:B:91:THR:HG23	2.14	0.48
1:B:92:LEU:HD23	3:B:339:DMA:H52	1.95	0.48
1:A:60:PRO:HG3	1:A:85:GLU:HB2	1.96	0.47
1:A:158:VAL:HG11	1:B:43:ILE:HG13	2.00	0.44
1:A:129:ARG:HD3	1:B:151:ARG:HG3	2.00	0.43
1:A:140:MET:HE3	1:A:143:LEU:HB2	2.00	0.43
1:A:20:LEU:HD11	1:A:318:ASP:HB3	2.00	0.43
1:B:214:LEU:HD12	1:B:327:ALA:HA	2.00	0.42
1:A:140:MET:HE1	1:B:143:LEU:HD13	2.03	0.41
1:B:238:ALA:O	1:B:243:ARG:NH2	2.54	0.41
1:A:171:GLN:HE21	1:A:253:LEU:HD21	1.86	0.41
1:A:174:TYR:OH	1:A:223:ASP:OD2	2.38	0.40
1:A:201:VAL:O	1:A:205:GLU:HB2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	292/345 (85%)	282 (97%)	9 (3%)	1 (0%)	36	35
1	B	334/345 (97%)	330 (99%)	4 (1%)	0	100	100
All	All	626/690 (91%)	612 (98%)	13 (2%)	1 (0%)	43	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	260	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	219/253 (87%)	198 (90%)	21 (10%)	8	5
1	B	248/253 (98%)	233 (94%)	15 (6%)	17	14
All	All	467/506 (92%)	431 (92%)	36 (8%)	13	8

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	SER
1	A	19[A]	ARG
1	A	19[B]	ARG
1	A	42	ILE
1	A	46	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	48	GLU
1	A	62[A]	MET
1	A	62[B]	MET
1	A	75	ASP
1	A	97	VAL
1	A	113	ILE
1	A	118	GLN
1	A	129	ARG
1	A	133	LEU
1	A	143	LEU
1	A	144	GLU
1	A	164	SER
1	A	167	LEU
1	A	226	ASP
1	A	281	ARG
1	A	293	LEU
1	B	119	SER
1	B	140	MET
1	B	143	LEU
1	B	167	LEU
1	B	190	SER
1	B	196	SER
1	B	200	ASP
1	B	212[A]	LEU
1	B	212[B]	LEU
1	B	214	LEU
1	B	233[A]	THR
1	B	233[B]	THR
1	B	257	ARG
1	B	261	ARG
1	B	267	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	74	ASN
1	A	111	HIS
1	A	118	GLN
1	A	138	ASN
1	A	160	GLN
1	A	165	HIS
1	A	171	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	76	HIS
1	B	171	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	DMA	B	339	4	12,13,13	0.95	1 (8%)	15,19,19	1.50	2 (13%)
3	DMA	B	336	-	12,13,13	0.98	1 (8%)	15,19,19	1.54	5 (33%)
2	POP	A	336	-	6,8,8	0.72	0	12,13,13	1.03	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	DMA	B	339	4	-	1/13/13/13	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	DMA	B	336	-	-	0/13/13/13	-
2	POP	A	336	-	-	0/6/6/6	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	336	DMA	PA-O3A	2.83	1.62	1.59
3	B	339	DMA	PA-O3A	2.45	1.62	1.59

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	336	DMA	C1-C2-C3	-2.81	122.22	126.76
3	B	339	DMA	C1-C2-C3	-2.59	122.57	126.76
3	B	336	DMA	C5-C3-C4	2.53	120.40	114.59
3	B	339	DMA	O3A-PB-O1B	-2.52	97.80	111.04
3	B	336	DMA	O2B-PB-O3A	-2.41	96.56	104.64
3	B	336	DMA	O3B-PB-O2B	2.17	115.94	107.80
3	B	336	DMA	C5-C3-C2	-2.02	116.61	122.66

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	339	DMA	C2-C1-O1-PA

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	339	DMA	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	292/345 (84%)	0.49	31 (10%) 11 10	15, 45, 122, 139	8 (2%)
1	B	328/345 (95%)	-0.11	2 (0%) 85 85	13, 37, 85, 112	8 (2%)
All	All	620/690 (89%)	0.17	33 (5%) 32 31	13, 40, 104, 139	16 (2%)

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	167	LEU	5.6
1	A	263	ALA	4.7
1	A	114	TRP	4.0
1	A	98	VAL	3.9
1	A	276	THR	3.6
1	A	165	HIS	3.6
1	A	265	PHE	3.5
1	A	283	ALA	3.4
1	A	259	GLY	3.4
1	A	108	VAL	3.3
1	A	258	SER	3.2
1	A	266	TRP	3.0
1	A	267	GLU	3.0
1	A	227	TYR	3.0
1	A	292	ALA	2.9
1	A	53	ALA	2.8
1	A	109	ALA	2.6
1	A	260	PRO	2.6
1	A	166	ASP	2.4
1	A	247	ALA	2.4
1	B	256	ALA	2.4
1	A	169	LEU	2.3
1	A	294	ASP	2.3
1	A	52	ALA	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	269	ALA	2.3
1	A	256	ALA	2.2
1	A	280	PHE	2.1
1	A	72	ALA	2.1
1	B	263	ALA	2.1
1	A	295	ALA	2.1
1	A	224	ALA	2.1
1	A	255	ILE	2.0
1	A	288	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	POP	A	336	9/9	0.76	0.10	107,112,114,116	0
3	DMA	B	336	14/14	0.98	0.06	24,34,40,44	0
3	DMA	B	339	14/14	0.98	0.06	32,37,47,55	0
4	CA	B	337	1/1	0.99	0.02	33,33,33,33	0
4	CA	B	338	1/1	0.99	0.04	39,39,39,39	0

6.5 Other polymers [i](#)

There are no such residues in this entry.