



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 10:20 AM UTC

PDB ID : 4OYY / pdb_00004oyy
Title : Humicola insolens cutinase
Authors : Dauter, Z.D.; Brzozowski, A.M.; Turkenburg, J.P.; Wilson, K.S.
Deposited on : 2014-02-13
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	194	90% 8% .
1	B	194	87% 12% .
1	C	194	87% 11% ..
1	D	194	89% 9% ..
1	E	194	86% 9% ...

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	194	86% 13% ..
1	G	194	89% 9% ...
1	H	194	89% 9% ..
1	I	194	% 89% 10% ..
1	J	194	87% 10% ...
1	K	194	% 87% 10% ...
1	L	194	% 89% 9% ..

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 16975 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cutinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	191	Total	C	N	O	S	0	0	0
			1402	874	257	266	5			
1	B	191	Total	C	N	O	S	0	0	0
			1397	872	254	266	5			
1	C	193	Total	C	N	O	S	0	0	0
			1419	885	260	269	5			
1	D	191	Total	C	N	O	S	0	0	0
			1402	874	257	266	5			
1	E	191	Total	C	N	O	S	0	0	0
			1396	871	254	266	5			
1	F	193	Total	C	N	O	S	0	0	0
			1419	885	260	269	5			
1	G	193	Total	C	N	O	S	0	0	0
			1416	884	260	267	5			
1	H	191	Total	C	N	O	S	0	0	0
			1402	874	257	266	5			
1	I	193	Total	C	N	O	S	0	0	0
			1410	881	256	268	5			
1	J	191	Total	C	N	O	S	0	0	0
			1402	874	257	266	5			
1	K	191	Total	C	N	O	S	0	0	0
			1402	874	257	266	5			
1	L	192	Total	C	N	O	S	0	0	0
			1410	880	258	267	5			

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	10	Total	O	0	0
			10	10		
2	B	4	Total	O	0	0
			4	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	C	7	Total 7	O 7	0	0
2	D	8	Total 8	O 8	0	0
2	E	3	Total 3	O 3	0	0
2	F	8	Total 8	O 8	0	0
2	G	15	Total 15	O 15	0	0
2	H	8	Total 8	O 8	0	0
2	I	8	Total 8	O 8	0	0
2	J	10	Total 10	O 10	0	0
2	K	9	Total 9	O 9	0	0
2	L	8	Total 8	O 8	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cutinase

Chain A: 




- Molecule 1: Cutinase

Chain B: 



- Molecule 1: Cutinase

Chain C: 



- Molecule 1: Cutinase

Chain D: 



- Molecule 1: Cutinase

Chain E: 



- Molecule 1: Cutinase

Chain F: 



- Molecule 1: Cutinase

Chain G: 89% 9% ...



- Molecule 1: Cutinase

Chain H: 89% 9% ..



- Molecule 1: Cutinase

Chain I: 89% 10% ..




- Molecule 1: Cutinase

Chain J: 87% 10% ...



- Molecule 1: Cutinase

Chain K: 87% 10% ...



- Molecule 1: Cutinase

Chain L: 89% 9% ..



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	125.55Å 127.00Å 134.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.76 – 3.00 29.76 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.4 (29.76-3.00) 99.4 (29.76-3.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.76 (at 3.00Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.174 , 0.196 0.181 , 0.201	Depositor DCC
R_{free} test set	1338 reflections (3.06%)	wwPDB-VP
Wilson B-factor (Å ²)	34.1	Xtriage
Anisotropy	0.332	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 36.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.000 for k,h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	16975	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 48.72 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.2586e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.85	0/1426	0.92	2/1936 (0.1%)
1	B	0.86	0/1421	0.95	3/1930 (0.2%)
1	C	0.84	1/1443 (0.1%)	1.00	5/1959 (0.3%)
1	D	0.85	1/1426 (0.1%)	0.94	2/1936 (0.1%)
1	E	0.83	3/1420 (0.2%)	1.01	5/1929 (0.3%)
1	F	0.82	0/1443	0.99	3/1959 (0.2%)
1	G	0.87	1/1440 (0.1%)	0.97	2/1955 (0.1%)
1	H	0.84	0/1426	0.94	1/1936 (0.1%)
1	I	0.82	0/1434	0.96	2/1948 (0.1%)
1	J	0.86	1/1426 (0.1%)	1.01	6/1936 (0.3%)
1	K	0.84	1/1426 (0.1%)	0.96	4/1936 (0.2%)
1	L	0.87	2/1434 (0.1%)	1.02	5/1947 (0.3%)
All	All	0.85	10/17165 (0.1%)	0.97	40/23307 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	2

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	J	29	THR	CA-CB	6.65	1.62	1.54
1	L	72	PRO	CA-C	6.54	1.62	1.52
1	E	168	ILE	N-CA	6.28	1.53	1.46
1	K	168	ILE	N-CA	5.39	1.52	1.46
1	D	169	ILE	CA-CB	5.37	1.60	1.53

The worst 5 of 40 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	192	ILE	CA-CB-CG1	8.81	125.37	110.40
1	E	4	ALA	N-CA-CB	8.45	123.07	110.40
1	B	34	MET	CG-SD-CE	-8.14	82.98	100.90
1	F	158	VAL	N-CA-C	-7.62	99.75	109.58
1	E	126	VAL	N-CA-CB	7.29	119.77	111.90

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	1	GLN	Peptide
1	C	3	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1402	0	1394	5	0
1	B	1397	0	1385	6	0
1	C	1419	0	1416	8	0
1	D	1402	0	1394	6	0
1	E	1396	0	1383	13	0
1	F	1419	0	1416	11	0
1	G	1416	0	1414	12	0
1	H	1402	0	1394	6	0
1	I	1410	0	1401	9	0
1	J	1402	0	1394	9	0
1	K	1402	0	1394	11	0
1	L	1410	0	1405	7	0
2	A	10	0	0	0	0
2	B	4	0	0	1	0
2	C	7	0	0	0	0
2	D	8	0	0	0	0
2	E	3	0	0	0	0
2	F	8	0	0	0	0
2	G	15	0	0	0	0
2	H	8	0	0	1	0
2	I	8	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	J	10	0	0	0	0
2	K	9	0	0	1	0
2	L	8	0	0	1	0
All	All	16975	0	16790	97	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 97 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:68:THR:HG23	1:E:106:GLN:OE1	1.63	0.98
1:C:94:CYS:HB3	1:C:97:THR:HG23	1.48	0.96
1:G:1:GLN:OE1	1:G:1:GLN:N	2.03	0.90
1:E:68:THR:CG2	1:E:106:GLN:OE1	2.22	0.88
1:L:72:PRO:HG2	2:L:203:HOH:O	1.75	0.86

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	189/194 (97%)	181 (96%)	7 (4%)	1 (0%)	24	60
1	B	189/194 (97%)	180 (95%)	8 (4%)	1 (0%)	24	60
1	C	191/194 (98%)	181 (95%)	9 (5%)	1 (0%)	24	60
1	D	189/194 (97%)	181 (96%)	7 (4%)	1 (0%)	24	60
1	E	189/194 (97%)	180 (95%)	7 (4%)	2 (1%)	11	43
1	F	191/194 (98%)	182 (95%)	8 (4%)	1 (0%)	24	60
1	G	191/194 (98%)	183 (96%)	7 (4%)	1 (0%)	24	60

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	H	189/194 (97%)	181 (96%)	7 (4%)	1 (0%)	24 60
1	I	191/194 (98%)	182 (95%)	8 (4%)	1 (0%)	24 60
1	J	189/194 (97%)	181 (96%)	7 (4%)	1 (0%)	24 60
1	K	189/194 (97%)	180 (95%)	8 (4%)	1 (0%)	24 60
1	L	190/194 (98%)	183 (96%)	6 (3%)	1 (0%)	24 60
All	All	2277/2328 (98%)	2175 (96%)	89 (4%)	13 (1%)	21 56

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	4	ALA
1	E	120	GLY
1	F	120	GLY
1	I	120	GLY
1	J	120	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	141/143 (99%)	137 (97%)	4 (3%)	38 70
1	B	140/143 (98%)	132 (94%)	8 (6%)	18 52
1	C	143/143 (100%)	135 (94%)	8 (6%)	19 52
1	D	141/143 (99%)	137 (97%)	4 (3%)	38 70
1	E	140/143 (98%)	133 (95%)	7 (5%)	22 56
1	F	143/143 (100%)	137 (96%)	6 (4%)	26 61
1	G	142/143 (99%)	136 (96%)	6 (4%)	26 61
1	H	141/143 (99%)	133 (94%)	8 (6%)	18 52
1	I	141/143 (99%)	135 (96%)	6 (4%)	26 60
1	J	141/143 (99%)	135 (96%)	6 (4%)	26 60
1	K	141/143 (99%)	133 (94%)	8 (6%)	18 52

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	L	142/143 (99%)	137 (96%)	5 (4%)	32 65
All	All	1696/1716 (99%)	1620 (96%)	76 (4%)	24 59

5 of 76 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	J	13	SER
1	L	2	LEU
1	J	29	THR
1	K	129	VAL
1	L	179	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	I	106	GLN
1	K	91	ASN
1	L	91	ASN
1	L	49	HIS
1	J	106	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å²)	Q<0.9	
1	A	191/194 (98%)	-0.53	0	100	100	10, 20, 41, 55	0
1	B	191/194 (98%)	-0.40	0	100	100	17, 28, 42, 54	0
1	C	193/194 (99%)	-0.43	0	100	100	10, 22, 47, 68	0
1	D	191/194 (98%)	-0.44	0	100	100	12, 26, 45, 63	0
1	E	191/194 (98%)	-0.18	0	100	100	16, 34, 56, 65	0
1	F	193/194 (99%)	-0.44	0	100	100	12, 25, 48, 74	0
1	G	193/194 (99%)	-0.59	0	100	100	9, 20, 36, 56	0
1	H	191/194 (98%)	-0.55	0	100	100	12, 22, 37, 64	0
1	I	193/194 (99%)	-0.40	2 (1%)	79	59	11, 27, 51, 75	0
1	J	191/194 (98%)	-0.40	0	100	100	12, 28, 51, 83	0
1	K	191/194 (98%)	-0.49	1 (0%)	87	72	11, 24, 40, 68	0
1	L	192/194 (98%)	-0.37	1 (0%)	87	72	14, 29, 48, 69	0
All	All	2301/2328 (98%)	-0.44	4 (0%)	91	83	9, 26, 48, 83	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	K	167	LEU	2.9
1	I	1	GLN	2.7
1	L	72	PRO	2.2
1	I	121	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.