



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 1, 2026 – 11:43 PM UTC

PDB ID : 1OZ3 / pdb_00001oz3
Title : Crystal Structure of 3-MBT repeats of lethal (3) malignant Brain Tumor (Native-I) at 1.85 angstrom
Authors : Wang, W.K.; Tereshko, V.; Boccuni, P.; MacGrogan, D.; Nimer, S.D.; Patel, D.J.
Deposited on : 2003-04-07
Resolution : 1.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	331	
1	B	331	
1	C	331	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	A	1005	-	-	X	-
3	MES	C	1011	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 8232 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lethal(3)malignant brain tumor-like protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	313	Total	C	N	O	S	Se	0	0	0
			2545	1638	432	463	7	5			
1	B	313	Total	C	N	O	S	Se	0	0	0
			2545	1638	432	463	7	5			
1	C	313	Total	C	N	O	S	Se	0	0	0
			2545	1638	432	463	7	5			

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: $C_6H_{13}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	C	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	C	1	Total	C	N	O	S	0	0
			12	6	1	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	172	Total	O	0	0
			172	172		
4	B	189	Total	O	0	0
			189	189		
4	C	114	Total	O	0	0
			114	114		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Lethal(3)malignant brain tumor-like protein



- Molecule 1: Lethal(3)malignant brain tumor-like protein



- Molecule 1: Lethal(3)malignant brain tumor-like protein





4 Data and refinement statistics

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	105.25Å 105.25Å 90.49Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.84 – 1.85 19.84 – 1.85	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.84-1.85) 99.9 (19.84-1.85)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.58 (at 1.85Å)	Xtriage
Refinement program	REFMAC 5.1.13	Depositor
R, R_{free}	0.211 , 0.245 0.217 , 0.248	Depositor DCC
R_{free} test set	4822 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	24.7	Xtriage
Anisotropy	0.074	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 30.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.014 for -h,-k,l 0.026 for h,-h-k,-l 0.018 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8232	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MES, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.98	3/2639 (0.1%)	1.08	6/3599 (0.2%)
1	B	0.97	1/2639 (0.0%)	1.06	9/3599 (0.3%)
1	C	0.95	0/2639	1.12	7/3599 (0.2%)
All	All	0.97	4/7917 (0.1%)	1.09	22/10797 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	341	PRO	CA-C	6.81	1.55	1.51
1	A	357	MSE	SE-CE	6.06	2.13	1.95
1	B	458	VAL	CB-CG2	-5.36	1.34	1.52
1	A	213	GLU	CA-C	-5.21	1.46	1.52

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	267	ARG	NE-CZ-NH2	-9.58	110.58	119.20
1	C	341	PRO	N-CA-C	8.92	121.58	110.70
1	A	309	TYR	N-CA-C	8.00	120.25	110.41
1	B	267	ARG	NE-CZ-NH2	-7.56	112.39	119.20
1	A	414	GLN	CA-C-N	-7.19	109.53	123.13

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	321	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2545	0	2380	41	0
1	B	2545	0	2380	29	0
1	C	2545	0	2380	59	0
2	A	10	0	0	2	0
2	B	25	0	0	1	0
2	C	15	0	0	1	0
3	A	24	0	25	1	0
3	B	24	0	25	4	0
3	C	24	0	25	11	0
4	A	172	0	0	4	0
4	B	189	0	0	5	0
4	C	114	0	0	6	0
All	All	8232	0	7215	129	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 129 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:275:TYR:CD2	3:C:1011:MES:H81	2.08	0.88
1:A:301:HIS:HE1	1:A:369:ASP:OD1	1.63	0.82
1:C:301:HIS:HE1	1:C:369:ASP:OD1	1.62	0.81
1:C:347:VAL:HG12	4:C:1091:HOH:O	1.80	0.80
1:C:408:LYS:HG3	1:C:409:PRO:HD2	1.62	0.80

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	311/331 (94%)	304 (98%)	7 (2%)	0	100	100
1	B	311/331 (94%)	307 (99%)	4 (1%)	0	100	100
1	C	311/331 (94%)	303 (97%)	7 (2%)	1 (0%)	36	25
All	All	933/993 (94%)	914 (98%)	18 (2%)	1 (0%)	48	35

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	341	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	277/286 (97%)	270 (98%)	7 (2%)	42	27
1	B	277/286 (97%)	265 (96%)	12 (4%)	26	11
1	C	277/286 (97%)	258 (93%)	19 (7%)	14	3
All	All	831/858 (97%)	793 (95%)	38 (5%)	24	9

5 of 38 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	351	LEU
1	C	430	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	354	VAL
1	C	385	THR
1	C	514	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 26 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	358	ASN
1	B	414	GLN
1	C	406	GLN
1	B	406	GLN
1	B	488	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	B	1010	-	4,4,4	0.35	0	6,6,6	0.54	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	B	1003	-	4,4,4	0.32	0	6,6,6	1.02	0
3	MES	C	1012	-	12,12,12	1.23	1 (8%)	15,16,16	1.32	1 (6%)
2	SO4	B	1001	-	4,4,4	0.23	0	6,6,6	0.80	0
2	SO4	C	1004	-	4,4,4	0.37	0	6,6,6	0.84	0
3	MES	B	1021	-	12,12,12	0.95	1 (8%)	15,16,16	2.56	5 (33%)
2	SO4	A	1005	-	4,4,4	0.44	0	6,6,6	0.80	0
3	MES	A	1031	-	12,12,12	1.11	1 (8%)	15,16,16	2.10	4 (26%)
3	MES	C	1011	-	12,12,12	2.76	3 (25%)	15,16,16	2.25	5 (33%)
2	SO4	B	1007	-	4,4,4	0.47	0	6,6,6	0.98	1 (16%)
2	SO4	B	1009	-	4,4,4	0.27	0	6,6,6	1.35	2 (33%)
3	MES	A	1032	-	12,12,12	1.08	1 (8%)	15,16,16	1.70	2 (13%)
3	MES	B	1022	-	12,12,12	0.85	0	15,16,16	1.68	4 (26%)
2	SO4	C	1006	-	4,4,4	0.39	0	6,6,6	0.54	0
2	SO4	C	1008	-	4,4,4	0.60	0	6,6,6	0.80	0
2	SO4	A	1002	-	4,4,4	0.36	0	6,6,6	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MES	C	1012	-	-	2/6/14/14	0/1/1/1
3	MES	C	1011	-	-	5/6/14/14	0/1/1/1
3	MES	B	1021	-	-	1/6/14/14	0/1/1/1
3	MES	A	1031	-	-	4/6/14/14	0/1/1/1
3	MES	A	1032	-	-	1/6/14/14	0/1/1/1
3	MES	B	1022	-	-	5/6/14/14	0/1/1/1

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	1011	MES	C8-S	7.43	1.88	1.77
3	C	1011	MES	C7-N4	4.16	1.56	1.47
3	C	1012	MES	C8-S	3.84	1.83	1.77
3	C	1011	MES	C7-C8	3.46	1.61	1.52
3	A	1032	MES	C8-S	3.07	1.81	1.77

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1021	MES	O1-C6-C5	-6.30	98.20	111.77
3	A	1031	MES	O1-C2-C3	-5.17	100.63	111.77
3	B	1021	MES	C5-N4-C3	4.43	118.39	108.84
3	C	1011	MES	C8-C7-N4	4.41	129.04	112.36
3	C	1011	MES	C7-N4-C5	4.10	122.16	111.24

There are no chirality outliers.

5 of 18 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1031	MES	N4-C7-C8-S
3	A	1031	MES	C7-C8-S-O1S
3	A	1031	MES	C7-C8-S-O3S
3	A	1032	MES	C8-C7-N4-C3
3	B	1021	MES	N4-C7-C8-S

There are no ring outliers.

7 monomers are involved in 20 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	1012	MES	3	0
3	B	1021	MES	4	0
2	A	1005	SO4	2	0
3	A	1031	MES	1	0
3	C	1011	MES	8	0
2	B	1009	SO4	1	0
2	C	1006	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	308/331 (93%)	0.37	20 (6%) 25 27	17, 26, 39, 47	0
1	B	308/331 (93%)	0.10	11 (3%) 46 50	13, 23, 34, 42	0
1	C	308/331 (93%)	0.60	35 (11%) 10 9	17, 28, 44, 50	0
All	All	924/993 (93%)	0.36	66 (7%) 22 24	13, 25, 39, 50	0

The worst 5 of 66 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	310	LYS	5.1
1	C	417	PRO	5.0
1	A	312	GLU	4.8
1	C	443	VAL	4.6
1	A	343	LEU	4.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MES	B	1021	12/12	0.61	0.19	40,42,50,51	0
3	MES	A	1031	12/12	0.67	0.17	37,39,49,49	0
2	SO4	C	1008	5/5	0.70	0.22	36,40,45,46	0
3	MES	C	1011	12/12	0.73	0.19	36,41,48,48	0
2	SO4	A	1005	5/5	0.74	0.23	47,47,49,52	0
2	SO4	B	1007	5/5	0.74	0.19	42,43,46,49	0
2	SO4	C	1006	5/5	0.78	0.27	49,51,54,54	0
3	MES	C	1012	12/12	0.79	0.14	33,37,41,42	0
2	SO4	B	1010	5/5	0.82	0.20	37,41,41,43	0
2	SO4	B	1009	5/5	0.87	0.14	33,33,37,38	0
2	SO4	C	1004	5/5	0.90	0.18	40,42,43,46	0
3	MES	B	1022	12/12	0.90	0.10	20,25,37,39	0
3	MES	A	1032	12/12	0.92	0.09	24,29,35,36	0
2	SO4	B	1003	5/5	0.93	0.14	37,40,43,43	0
2	SO4	B	1001	5/5	0.95	0.09	30,31,33,34	0
2	SO4	A	1002	5/5	0.95	0.08	37,41,41,43	0

6.5 Other polymers [i](#)

There are no such residues in this entry.