



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 27, 2026 – 05:28 PM EDT

PDB ID : 2OZ4 / pdb_00002oz4
Title : Structural Plasticity in IgSF Domain 4 of ICAM-1 Mediates Cell Surface Dimerization
Authors : Chen, X.; Kim, T.D.; Carman, C.V.; Mi, L.; Song, G.; Springer, T.A.
Deposited on : 2007-02-23
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	265	9% 92% 8%
2	L	214	8% 91% 9%
3	H	214	2% 89% 8% .
4	B	2	50% 50%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
8	TRS	L	2003	-	X	-	-

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 5601 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Intercellular adhesion molecule 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	265	Total	C	N	O	S	0	1	0
			2022	1258	355	400	9			

- Molecule 2 is a protein called FAB FRAGMENT LIGHT CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	214	Total	C	N	O	S	0	0	0
			1659	1031	280	342	6			

- Molecule 3 is a protein called FAB FRAGMENT, HEAVY CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	H	208	Total	C	N	O	S	0	0	0
			1564	993	252	312	7			

- Molecule 4 is an oligosaccharide called alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	B	2	Total	C	N	O	0	0	0
			24	14	1	9			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	0
			5	4	1		
6	L	1	Total	O	S	0	0
			5	4	1		

- Molecule 7 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	L	2	Total	Zn	0	0
			2	2		

- Molecule 8 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: C₄H₁₂NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	L	1	Total	C	N	O	0	0
			8	4	1	3		
8	L	1	Total	C	N	O	0	0
			8	4	1	3		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	127	Total	O	0	0
			127	127		
9	L	86	Total	O	0	0
			86	86		
9	H	63	Total	O	0	0
			63	63		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Intercellular adhesion molecule 1



- Molecule 2: FAB FRAGMENT LIGHT CHAIN



- Molecule 3: FAB FRAGMENT, HEAVY CHAIN



- Molecule 4: alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	185.40Å 69.34Å 88.17Å 90.00° 112.76° 90.00°	Depositor
Resolution (Å)	28.50 – 2.70 28.50 – 2.70	Depositor EDS
% Data completeness (in resolution range)	100.0 (28.50-2.70) 90.6 (28.50-2.70)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.18 (at 2.72Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.204 , 0.253 0.204 , 0.250	Depositor DCC
R_{free} test set	2559 reflections (9.86%)	wwPDB-VP
Wilson B-factor (Å ²)	77.0	Xtriage
Anisotropy	0.067	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 63.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	5601	wwPDB-VP
Average B, all atoms (Å ²)	78.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, NAG, TRS, ZN, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	0/2062	0.56	0/2821
2	L	0.39	0/1696	0.57	0/2302
3	H	0.38	0/1604	0.60	0/2190
All	All	0.39	0/5362	0.58	0/7313

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2022	0	2006	10	0
2	L	1659	0	1581	15	0
3	H	1564	0	1535	10	0
4	B	24	0	22	0	0
5	A	28	0	26	0	0
6	A	5	0	0	0	0
6	L	5	0	0	0	0
7	L	2	0	0	0	0
8	L	16	0	24	1	0
9	A	127	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
9	H	63	0	0	0	0
9	L	86	0	0	0	0
All	All	5601	0	5194	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:H:12:VAL:HG21	3:H:86:LEU:HD13	1.76	0.67
2:L:37:GLN:HB2	2:L:47:LEU:HD11	1.79	0.63
2:L:11:LEU:HD21	2:L:19:VAL:HG22	1.82	0.60
3:H:126:ALA:HB2	3:H:211:ILE:CG2	2.31	0.60
3:H:4:LEU:HD22	3:H:24:THR:HG22	1.82	0.60
2:L:19:VAL:HG12	2:L:75:ILE:HB	1.84	0.59
1:A:412:LEU:HD12	1:A:413:PRO:HD2	1.87	0.55
2:L:2:ILE:HD13	2:L:29:ILE:CG2	2.38	0.53
2:L:161:ASN:HB3	2:L:175:MET:HE3	1.94	0.50
2:L:150:ILE:CD1	2:L:181:LEU:HD11	2.43	0.49
3:H:139:LEU:HD12	3:H:194:VAL:HG11	1.93	0.49
1:A:215:LEU:HD12	1:A:218:VAL:HG22	1.94	0.48
1:A:225:LEU:HG	1:A:232:LEU:HD12	1.95	0.48
2:L:160:LEU:HD11	3:H:172:GLN:HG3	1.95	0.47
1:A:405:LEU:HD23	1:A:411:PRO:HA	1.97	0.47
1:A:234:PRO:HA	1:A:247:ALA:HB2	1.97	0.46
3:H:126:ALA:HB2	3:H:211:ILE:HG22	1.96	0.46
1:A:313:VAL:HG21	1:A:326:ARG:HA	1.98	0.46
1:A:369:PRO:O	1:A:439:VAL:HG21	2.16	0.45
2:L:188:ARG:CG	2:L:188:ARG:O	2.64	0.45
2:L:2:ILE:HD13	2:L:29:ILE:HG21	1.98	0.44
2:L:11:LEU:HD21	2:L:19:VAL:CG2	2.47	0.44
3:H:39:GLN:O	3:H:92:ALA:HB1	2.18	0.44
2:L:136:LEU:HD12	2:L:136:LEU:N	2.33	0.43
3:H:11:LEU:HD22	3:H:148:PRO:HG3	2.01	0.43
2:L:165:ASP:HA	8:L:2004:TRS:N	2.34	0.42
3:H:65:LYS:O	3:H:66:ASP:HB2	2.20	0.41
2:L:150:ILE:HD11	2:L:179:LEU:HD21	2.03	0.41
1:A:413:PRO:HG3	1:A:418:VAL:HG11	2.03	0.41
1:A:194:LEU:HD13	1:A:263:CYS:HB2	2.02	0.41
2:L:50:TYR:O	2:L:51:ALA:HB3	2.20	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:280:ILE:N	1:A:280:ILE:HD12	2.36	0.40
3:H:2:VAL:HG11	3:H:103:TYR:CG	2.56	0.40
2:L:140:TYR:CG	2:L:141:PRO:HA	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	264/265 (100%)	257 (97%)	6 (2%)	1 (0%)	30	54
2	L	212/214 (99%)	202 (95%)	10 (5%)	0	100	100
3	H	204/214 (95%)	197 (97%)	7 (3%)	0	100	100
All	All	680/693 (98%)	656 (96%)	23 (3%)	1 (0%)	48	73

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	321	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/230 (100%)	230 (100%)	1 (0%)	84	93

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	L	192/192 (100%)	192 (100%)	0	100	100
3	H	181/184 (98%)	180 (99%)	1 (1%)	78	91
All	All	604/606 (100%)	602 (100%)	2 (0%)	86	94

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	295	VAL
3	H	72	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	270	GLN
1	A	353	GLN
1	A	359	GLN
2	L	37	GLN
2	L	38	GLN
2	L	137	ASN
3	H	39	GLN
3	H	165	HIS
3	H	197	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	B	1	1,4	14,14,15	0.52	0	17,19,21	1.01	1 (5%)
4	FUC	B	2	4	10,10,11	0.63	0	14,14,16	0.61	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	B	1	1,4	-	0/6/23/26	0/1/1/1
4	FUC	B	2	4	-	-	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1	NAG	C1-O5-C5	3.40	116.75	112.19


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	A	1269	1	14,14,15	0.66	0	17,19,21	0.84	1 (5%)
6	SO4	A	2000	-	4,4,4	0.23	0	6,6,6	0.08	0
8	TRS	L	2004	-	7,7,7	0.32	0	9,9,9	0.28	0
8	TRS	L	2003	-	7,7,7	0.72	0	9,9,9	4.95	6 (66%)
5	NAG	A	1358	1	14,14,15	0.57	0	17,19,21	0.69	0
6	SO4	L	2000	-	4,4,4	0.24	0	6,6,6	0.14	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	TRS	L	2003	-	-	7/9/9/9	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	A	1269	1	-	0/6/23/26	0/1/1/1
5	NAG	A	1358	1	-	2/6/23/26	0/1/1/1
8	TRS	L	2004	-	-	4/9/9/9	-

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	L	2003	TRS	C3-C-N	-7.92	87.96	108.17
8	L	2003	TRS	C1-C-N	-7.87	88.11	108.17
8	L	2003	TRS	C2-C-N	-7.65	88.66	108.17
8	L	2003	TRS	C3-C-C2	4.07	121.48	110.66
8	L	2003	TRS	C3-C-C1	3.29	119.41	110.66
8	L	2003	TRS	C2-C-C1	3.07	118.83	110.66
5	A	1269	NAG	C4-C3-C2	2.27	114.35	111.02

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	L	2003	TRS	C3-C-C2-O2
8	L	2003	TRS	C2-C-C3-O3
8	L	2004	TRS	C3-C-C1-O1
8	L	2004	TRS	N-C-C1-O1
5	A	1358	NAG	O5-C5-C6-O6
8	L	2003	TRS	C3-C-C1-O1
8	L	2003	TRS	C1-C-C2-O2
8	L	2004	TRS	C2-C-C1-O1
8	L	2003	TRS	C2-C-C1-O1
8	L	2003	TRS	C1-C-C3-O3
8	L	2004	TRS	C2-C-C3-O3
5	A	1358	NAG	C4-C5-C6-O6
8	L	2003	TRS	N-C-C3-O3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	L	2004	TRS	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	265/265 (100%)	0.85	24 (9%)	15 12	37, 78, 83, 86	6 (2%)
2	L	214/214 (100%)	0.67	18 (8%)	17 14	62, 78, 82, 100	6 (2%)
3	H	208/214 (97%)	0.41	4 (1%)	66 63	60, 78, 82, 94	5 (2%)
All	All	687/693 (99%)	0.66	46 (6%)	24 21	37, 78, 83, 100	17 (2%)

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	361[A]	ARG	4.2
2	L	214	CYS	4.1
1	A	286	PRO	4.0
1	A	322	PRO	3.9
3	H	134	ASN	3.5
3	H	214	ARG	3.3
1	A	324	GLY	3.2
3	H	212	VAL	3.2
2	L	165	ASP	3.2
2	L	76	ASN	3.1
1	A	320	ALA	2.9
2	L	173	TYR	2.8
2	L	142	LYS	2.8
2	L	140	TYR	2.8
1	A	295	VAL	2.7
1	A	323	LEU	2.7
2	L	209	PHE	2.7
1	A	366	LEU	2.6
1	A	399	LEU	2.5
2	L	143	ASP	2.5
2	L	94	TRP	2.5
1	A	305	CYS	2.4
3	H	196	CYS	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	L	181	LEU	2.4
1	A	344	CYS	2.4
2	L	164	THR	2.3
1	A	283	PHE	2.3
1	A	390	PRO	2.3
1	A	450	PRO	2.3
1	A	363	LEU	2.3
1	A	415	GLY	2.2
2	L	199	LYS	2.2
1	A	285	ALA	2.2
1	A	374	ARG	2.2
2	L	213	GLU	2.2
2	L	187	GLU	2.1
1	A	357	LYS	2.1
1	A	193	GLN	2.1
1	A	338	ASN	2.1
2	L	141	PRO	2.1
1	A	435	THR	2.1
1	A	365	VAL	2.1
1	A	393	GLN	2.0
2	L	4	LEU	2.0
2	L	166	GLN	2.0
2	L	188	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	B	1	14/15	-	-	80,82,83,83	0
4	FUC	B	2	10/11	-	-	78,79,80,80	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	A	1269	14/15	0.26	0.17	92,95,97,97	0
8	TRS	L	2004	8/8	0.70	0.15	90,91,92,93	0
5	NAG	A	1358	14/15	0.75	0.13	86,89,90,90	0
8	TRS	L	2003	8/8	0.76	0.13	101,101,101,102	0
6	SO4	A	2000	5/5	0.88	0.13	111,111,111,111	5
6	SO4	L	2000	5/5	0.94	0.14	86,86,86,86	0
7	ZN	L	2001	1/1	0.98	0.12	79,79,79,79	0
7	ZN	L	2002	1/1	0.99	0.08	99,99,99,99	1

6.5 Other polymers [i](#)

There are no such residues in this entry.