



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 08:57 AM EDT

PDB ID : 2OZU / pdb_00002ozu
Title : Crystal structure of human MYST histone acetyltransferase 3 in complex with acetylcoenzyme A
Authors : Min, J.; Wu, H.; Dombrowski, L.; Bernstein, G.; Dong, A.; Loppnau, P.; Weigelt, J.; Sundstrom, M.; Arrowsmith, C.H.; Edwards, A.M.; Bochkarev, A.; Plotnikov, A.N.; Structural Genomics Consortium (SGC)
Deposited on : 2007-02-27
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	284	7% 70% 18% 11%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 2339 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone acetyltransferase MYST3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	254	Total	C	N	O	S	0	2	0
			2139	1387	364	369	19			


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	604	ALY	LYS	modified residue	UNP Q92794

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is ACETYL COENZYME *A (CCD ID: ACO) (formula: C₂₃H₃₈N₇O₁₇P₃S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	S	
			51	23	7	17	3	1	
								0	0

- Molecule 4 is ACETAMIDE (CCD ID: ACM) (formula: C₂H₅NO).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O		
			4	2	1	1	0	0
4	A	1	Total	C	N	O		
			4	2	1	1	0	0
4	A	1	Total	C	N	O		
			4	2	1	1	0	0
4	A	1	Total	C	N	O		
			4	2	1	1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	132	Total	O		
			132	132	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Histone acetyltransferase MYST3



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	58.85Å 104.05Å 109.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	75.38 – 2.30 75.38 – 2.30	Depositor EDS
% Data completeness (in resolution range)	98.1 (75.38-2.30) 98.2 (75.38-2.30)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.61 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.191 , 0.245 0.182 , 0.238	Depositor DCC
R_{free} test set	756 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	31.3	Xtriage
Anisotropy	0.328	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 54.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.008 for -h,-l,-k	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2339	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, ALY, ACM, ACO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.65	0/2183	0.87	0/2941

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2139	0	2136	33	0
2	A	1	0	0	0	0
3	A	51	0	34	0	0
4	A	16	0	13	1	0
5	A	132	0	0	2	1
All	All	2339	0	2183	33	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:534:LEU:HD12	1:A:536:LYS:O	1.84	0.77
1:A:534:LEU:HD13	1:A:535:PRO:HD2	1.76	0.66
1:A:734:LEU:HD13	1:A:736:MET:HE2	1.77	0.64
1:A:720:THR:HG23	1:A:722:ILE:H	1.63	0.64
1:A:541:GLU:OE1	1:A:627:HIS:HD2	1.84	0.60
1:A:704:TYR:HB2	1:A:757:MET:HE3	1.88	0.55
1:A:569:GLU:OE1	1:A:572:ARG:HD3	2.08	0.54
1:A:750:GLU:O	1:A:754:GLN:HG2	2.09	0.52
1:A:760:LEU:O	1:A:761:GLN:HB2	2.10	0.50
1:A:622:ASP:OD1	1:A:627:HIS:HE1	1.95	0.50
1:A:734:LEU:HD13	1:A:736:MET:CE	2.40	0.49
1:A:734:LEU:CD1	1:A:736:MET:HE2	2.43	0.49
1:A:720:THR:CG2	1:A:722:ILE:H	2.24	0.48
1:A:720:THR:HG23	1:A:722:ILE:HG13	1.96	0.48
1:A:577:SER:HB2	1:A:619:THR:OG1	2.14	0.47
1:A:635:GLU:CD	4:A:902:ACM:HN2	2.25	0.45
1:A:716:LEU:O	1:A:720:THR:HB	2.16	0.45
1:A:696:LYS:HE3	5:A:60:HOH:O	2.16	0.45
1:A:641:LYS:HE2	1:A:675:GLN:OE1	2.17	0.44
1:A:571:TYR:HB3	1:A:578:VAL:HB	1.99	0.44
1:A:692[B]:MET:HE1	1:A:733[B]:HIS:CB	2.47	0.43
1:A:642:TYR:HA	1:A:678:SER:O	2.18	0.43
1:A:582:ASP:HA	1:A:614:LEU:HD23	1.99	0.43
1:A:591:GLN:HG2	1:A:608:TYR:HA	2.00	0.43
1:A:692[B]:MET:O	1:A:696:LYS:HG3	2.18	0.43
1:A:538:TYR:HB3	1:A:553:LEU:HD22	2.01	0.42
1:A:606:LEU:HD12	1:A:607:TYR:H	1.83	0.42
1:A:692[B]:MET:HE1	1:A:733[B]:HIS:HB2	2.00	0.42
1:A:645:SER:HB3	5:A:68:HOH:O	2.20	0.42
1:A:632:PHE:HA	1:A:646:CYS:O	2.20	0.41
1:A:696:LYS:O	1:A:700:LEU:HG	2.20	0.41
1:A:508:CYS:HA	1:A:509:PRO:HA	1.98	0.41
1:A:692[A]:MET:O	1:A:696:LYS:HG3	2.20	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:56:HOH:O	5:A:56:HOH:O[2_555]	2.02	0.18

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	247/284 (87%)	238 (96%)	9 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	239/265 (90%)	227 (95%)	12 (5%)	22	33

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	510	SER
1	A	532	SER
1	A	534	LEU
1	A	559	LYS
1	A	602	ASP
1	A	640	GLN
1	A	645	SER
1	A	650	LEU
1	A	720	THR
1	A	735	ARG
1	A	774	LEU
1	A	775	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	584	ASN
1	A	627	HIS
1	A	640	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	ALY	A	604	1	10,11,12	0.54	0	7,12,14	0.84	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	ALY	A	604	1	-	1/9/10/12	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	604	ALY	O-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ACO	A	850	-	51,53,53	1.23	5 (9%)	73,79,79	1.64	14 (19%)
4	ACM	A	904	1	3,3,3	0.65	0	3,3,3	0.36	0
4	ACM	A	901	1	3,3,3	1.12	0	3,3,3	0.65	0
4	ACM	A	902	1	3,3,3	0.22	0	3,3,3	0.57	0
4	ACM	A	903	1	3,3,3	0.48	0	3,3,3	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ACO	A	850	-	-	5/51/67/67	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	850	ACO	C5A-C4A	4.77	1.47	1.39
3	A	850	ACO	C5A-C6A	2.89	1.49	1.41
3	A	850	ACO	P1A-O3A	2.34	1.62	1.59
3	A	850	ACO	C8A-N7A	2.33	1.36	1.31
3	A	850	ACO	P2A-O3A	2.01	1.61	1.59

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	850	ACO	C5A-C4A-N3A	-4.89	119.98	126.72
3	A	850	ACO	N3A-C4A-N9A	4.60	134.99	127.17
3	A	850	ACO	C4A-N9A-C8A	4.18	110.12	105.74
3	A	850	ACO	N3A-C2A-N1A	-3.75	122.91	128.58
3	A	850	ACO	C2A-N3A-C4A	3.62	120.66	111.83
3	A	850	ACO	C4A-C5A-N7A	-3.14	106.99	110.58
3	A	850	ACO	C5A-N7A-C8A	2.80	107.85	103.45
3	A	850	ACO	N9A-C8A-N7A	-2.76	110.02	113.94
3	A	850	ACO	C6A-C5A-N7A	2.35	136.62	132.09
3	A	850	ACO	O-C-CH3	-2.34	114.07	123.12
3	A	850	ACO	CH3-C-S1P	2.29	128.59	114.13
3	A	850	ACO	C2A-N1A-C6A	2.26	122.45	118.73
3	A	850	ACO	O5A-P2A-O4A	2.17	122.54	112.44
3	A	850	ACO	O4B-C1B-N9A	2.12	112.17	108.09

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	850	ACO	C3P-C2P-S1P-C
3	A	850	ACO	O-C-S1P-C2P
3	A	850	ACO	CH3-C-S1P-C2P
3	A	850	ACO	P2A-O3A-P1A-O1A
3	A	850	ACO	P2A-O3A-P1A-O2A


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	902	ACM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	253/284 (89%)	0.28	21 (8%) 17 19	6, 18, 41, 46	2 (0%)

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	531	TYR	4.5
1	A	760	LEU	4.3
1	A	762	LEU	4.3
1	A	753	ILE	4.0
1	A	534	LEU	3.6
1	A	747	ILE	3.4
1	A	756	HIS	3.4
1	A	757	MET	3.4
1	A	758	ALA	2.9
1	A	761	GLN	2.8
1	A	535	PRO	2.8
1	A	529	GLN	2.8
1	A	608	TYR	2.6
1	A	752	LEU	2.5
1	A	748	ARG	2.5
1	A	603	HIS	2.4
1	A	528	PRO	2.4
1	A	606	LEU	2.2
1	A	533	ARG	2.1
1	A	552	ILE	2.1
1	A	637	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	ALY	A	604	12/13	0.94	0.07	13,15,18,18	0

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	ACM	A	903	4/4	0.79	0.19	35,38,38,38	0
4	ACM	A	902	4/4	0.80	0.17	37,40,40,40	0
4	ACM	A	901	4/4	0.88	0.14	27,28,28,29	0
4	ACM	A	904	4/4	0.94	0.11	18,18,18,18	0
3	ACO	A	850	51/51	0.96	0.07	12,18,26,31	0
2	ZN	A	800	1/1	0.99	0.07	23,23,23,23	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.