



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 04:36 PM UTC

PDB ID : 3OZE / pdb_00003oze
Title : Crystal Structure of human 5'-deoxy-5'-methyladenosine phosphorylase
Authors : Ho, M.; Guan, R.; Almo, S.C.; Schramm, V.L.
Deposited on : 2010-09-24
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	283	81%11%• 7%
1	B	283	2% 83%10%• 5%
1	C	283	% 79%12%• 8%
1	D	283	% 80%12%• 7%
1	E	283	% 79%13%• 7%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	283	81% 12% • 6%

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 12542 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called S-methyl-5'-thioadenosine phosphorylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	263	Total	C	N	O	S	0	0	0
			2031	1286	350	378	17			
1	B	268	Total	C	N	O	S	0	0	0
			2072	1309	358	388	17			
1	C	261	Total	C	N	O	S	0	0	0
			2016	1275	348	376	17			
1	D	263	Total	C	N	O	S	0	0	0
			2031	1286	350	378	17			
1	E	264	Total	C	N	O	S	0	0	0
			2043	1292	353	381	17			
1	F	265	Total	C	N	O	S	0	0	0
			2050	1296	355	382	17			

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).







Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	C	1	Total	O	P	0	0
			5	4	1		
2	D	1	Total	O	P	0	0
			5	4	1		
2	E	1	Total	O	P	0	0
			5	4	1		
2	F	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is water.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	71	Total	O	0	0
			71	71		
3	B	35	Total	O	0	0
			35	35		
3	C	38	Total	O	0	0
			38	38		
3	D	45	Total	O	0	0
			45	45		
3	E	41	Total	O	0	0
			41	41		
3	F	39	Total	O	0	0
			39	39		

- Molecule 1: S-methyl-5'-thioadenosine phosphorylase







● Molecule 1: S-methyl-5'-thioadenosine phosphorylase



● Molecule 1: S-methyl-5'-thioadenosine phosphorylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	79.52Å 81.81Å 130.28Å 90.00° 91.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.00 20.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.0 (20.00-2.00) 99.0 (20.00-2.00)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.44 (at 2.01Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.216 , 0.262 0.226 , 0.272	Depositor DCC
R_{free} test set	5598 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	40.2	Xtriage
Anisotropy	0.146	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 45.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.000 for -k,-h,-l 0.000 for k,h,-l 0.010 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	12542	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.04	2/2070 (0.1%)	1.05	1/2802 (0.0%)
1	B	0.85	1/2111 (0.0%)	1.02	6/2857 (0.2%)
1	C	0.85	0/2054	0.97	0/2779
1	D	0.91	1/2070 (0.0%)	1.01	2/2802 (0.1%)
1	E	0.83	1/2083 (0.0%)	0.96	5/2819 (0.2%)
1	F	0.80	0/2088	0.95	1/2825 (0.0%)
All	All	0.88	5/12476 (0.0%)	0.99	15/16884 (0.1%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	202	VAL	CA-CB	6.65	1.62	1.54
1	E	255	ILE	CA-C	6.36	1.58	1.52
1	A	109	ILE	CA-C	6.32	1.58	1.52
1	D	255	ILE	CA-CB	5.88	1.57	1.54
1	B	233	VAL	CA-CB	-5.39	1.47	1.54

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	142	GLU	CA-C-N	6.93	126.54	119.19
1	E	142	GLU	C-N-CA	6.93	126.54	119.19
1	B	255	ILE	CA-C-N	-6.63	112.16	119.19
1	B	255	ILE	C-N-CA	-6.63	112.16	119.19
1	D	40	LYS	CA-C-N	6.09	126.06	120.03

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2031	0	2063	14	0
1	B	2072	0	2100	20	1
1	C	2016	0	2045	24	0
1	D	2031	0	2063	25	0
1	E	2043	0	2067	18	0
1	F	2050	0	2082	20	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
2	D	5	0	0	0	0
2	E	5	0	0	0	0
2	F	5	0	0	0	0
3	A	71	0	0	0	0
3	B	35	0	0	0	0
3	C	38	0	0	0	0
3	D	45	0	0	1	0
3	E	41	0	0	0	1
3	F	39	0	0	0	0
All	All	12542	0	12420	109	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 109 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:217:MET:HE3	1:D:247:ALA:HB2	1.29	1.15
1:E:14:ILE:HD11	1:E:57:LEU:HD13	1.42	1.00
1:A:68:MET:HE2	1:C:126:ASP:HA	1.52	0.91
1:D:217:MET:CE	1:D:247:ALA:HB2	2.05	0.86
1:A:30:THR:HG22	1:A:47:LEU:HB2	1.59	0.85

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:234:ASP:OD2	3:E:323:HOH:O[2_546]	2.10	0.10

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	259/283 (92%)	254 (98%)	5 (2%)	0	100	100
1	B	264/283 (93%)	253 (96%)	11 (4%)	0	100	100
1	C	257/283 (91%)	248 (96%)	9 (4%)	0	100	100
1	D	259/283 (92%)	255 (98%)	4 (2%)	0	100	100
1	E	260/283 (92%)	254 (98%)	6 (2%)	0	100	100
1	F	261/283 (92%)	253 (97%)	7 (3%)	1 (0%)	30	27
All	All	1560/1698 (92%)	1517 (97%)	42 (3%)	1 (0%)	48	46

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	29	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	225/242 (93%)	210 (93%)	15 (7%)	15	11
1	B	230/242 (95%)	219 (95%)	11 (5%)	23	21

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	223/242 (92%)	211 (95%)	12 (5%)	20	17
1	D	225/242 (93%)	210 (93%)	15 (7%)	15	11
1	E	226/242 (93%)	211 (93%)	15 (7%)	15	12
1	F	227/242 (94%)	210 (92%)	17 (8%)	12	9
All	All	1356/1452 (93%)	1271 (94%)	85 (6%)	16	13

5 of 85 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	47	LEU
1	F	47	LEU
1	E	128	SER
1	E	240	LEU
1	F	130	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	103	GLN
1	F	269	ASN
1	C	269	ASN
1	D	257	GLN
1	D	269	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	D	284	-	4,4,4	1.05	0	6,6,6	0.91	0
2	PO4	E	284	-	4,4,4	0.66	0	6,6,6	0.91	0
2	PO4	A	284	-	4,4,4	0.73	0	6,6,6	1.20	1 (16%)
2	PO4	C	284	-	4,4,4	0.82	0	6,6,6	1.04	0
2	PO4	F	284	-	4,4,4	1.03	0	6,6,6	1.37	1 (16%)
2	PO4	B	284	-	4,4,4	0.89	0	6,6,6	0.76	0

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	F	284	PO4	O3-P-O2	2.37	115.28	107.91
2	A	284	PO4	O2-P-O1	2.07	118.26	110.95

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	263/283 (92%)	-0.08	1 (0%) 88 88	30, 42, 59, 68	0
1	B	268/283 (94%)	0.17	6 (2%) 62 61	27, 51, 78, 86	0
1	C	261/283 (92%)	0.17	2 (0%) 82 82	35, 52, 71, 78	0
1	D	263/283 (92%)	0.04	2 (0%) 82 82	31, 46, 65, 75	0
1	E	264/283 (93%)	0.21	4 (1%) 72 71	34, 53, 67, 79	0
1	F	265/283 (93%)	0.20	1 (0%) 88 88	35, 54, 76, 83	0
All	All	1584/1698 (93%)	0.12	16 (1%) 79 79	27, 50, 71, 86	0

The worst 5 of 16 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	276	PHE	4.1
1	E	237	LEU	2.9
1	B	234	ASP	2.9
1	A	239	THR	2.7
1	B	235	ARG	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PO4	E	284	5/5	0.94	0.07	38,39,41,42	0
2	PO4	F	284	5/5	0.96	0.05	39,41,42,43	0
2	PO4	D	284	5/5	0.97	0.06	37,40,42,46	0
2	PO4	B	284	5/5	0.97	0.04	38,41,41,43	0
2	PO4	C	284	5/5	0.97	0.04	42,46,48,49	0
2	PO4	A	284	5/5	0.98	0.05	32,32,35,37	0

6.5 Other polymers [i](#)

There are no such residues in this entry.