



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 09:23 PM EDT

PDB ID : 3OZU / pdb_00003ozu
Title : The Crystal Structure of flavohemoglobin from *R. eutrophus* in complex with miconazole
Authors : El Hammi, E.; Warkentin, E.; Demmer, U.; Ermler, U.; Baciou, L.
Deposited on : 2010-09-27
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	403	10% 74% 23% .

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3666 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Flavohemoprotein.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	403	Total	C	N	O	S	0	6	0
			3204	2044	552	593	15			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 4 is 1-[(2R)-2-[(2,4-dichlorobenzyl)oxy]-2-(2,4-dichlorophenyl)ethyl]-1H-imidazole (CCD ID: X89) (formula: C₁₈H₁₄Cl₄N₂O).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	Cl	N	O	0	0
			25	18	4	2	1		

- Molecule 5 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	P	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	336	Total	O	0	0
			336	336		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Flavohemoprotein



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	121.34Å 105.28Å 46.45Å 90.00° 101.90° 90.00°	Depositor
Resolution (Å)	30.00 – 2.00 30.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.2 (30.00-2.00) 98.1 (30.00-2.00)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.56 (at 2.01Å)	Xtriage
Refinement program	REFMAC 5.6.0046	Depositor
R, R_{free}	0.185 , 0.233 0.199 , 0.239	Depositor DCC
R_{free} test set	1931 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	25.3	Xtriage
Anisotropy	0.056	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 49.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3666	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, X89, HEM, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.29	10/3294 (0.3%)	1.21	9/4472 (0.2%)

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	323	ASN	N-CA	6.85	1.55	1.46
1	A	205	ILE	C-O	-6.80	1.17	1.24
1	A	89	GLY	N-CA	6.47	1.54	1.45
1	A	16	VAL	CA-CB	5.67	1.60	1.54
1	A	162	LYS	N-CA	-5.46	1.40	1.46
1	A	395	VAL	C-O	5.27	1.29	1.24
1	A	253	ALA	CA-CB	5.26	1.62	1.53
1	A	106	ILE	C-O	5.17	1.29	1.24
1	A	25[A]	ILE	CA-CB	5.13	1.60	1.54
1	A	25[B]	ILE	CA-CB	5.13	1.60	1.54

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	232	PRO	CA-C-N	6.93	126.97	119.90
1	A	232	PRO	C-N-CA	6.93	126.97	119.90
1	A	91	LYS	CD-CE-NZ	-6.51	91.07	111.90
1	A	50	GLN	N-CA-C	6.42	118.81	111.11
1	A	14	ALA	CA-C-N	-5.54	113.19	119.28
1	A	14	ALA	C-N-CA	-5.54	113.19	119.28
1	A	127	GLY	N-CA-C	-5.44	106.20	112.73
1	A	201	GLY	N-CA-C	-5.28	106.95	115.08
1	A	51	GLY	N-CA-C	-5.05	106.64	112.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3204	0	3173	69	0
2	A	43	0	30	1	0
3	A	53	0	31	2	0
4	A	25	0	14	3	0
5	A	5	0	0	1	0
6	A	336	0	0	11	0
All	All	3666	0	3248	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:356:LYS:HD2	6:A:729:HOH:O	1.49	1.10
1:A:25[A]:ILE:HD12	4:A:411:X89:CL2A	1.93	1.06
1:A:25[A]:ILE:CD1	4:A:411:X89:CL2A	2.56	0.91
2:A:404:HEM:HMB1	5:A:406:PO4:O3	1.84	0.78
1:A:341:ARG:HG2	1:A:342:ASP:OD1	1.86	0.75
1:A:21:GLY:H	1:A:54[A]:GLN:HE22	1.39	0.70
1:A:79:LYS:CB	1:A:136:MET:HE1	2.22	0.69
1:A:18:ALA:O	1:A:54[A]:GLN:NE2	2.26	0.69
1:A:44:ASN:ND2	1:A:46:ALA:H	1.91	0.68
1:A:25[A]:ILE:HD11	1:A:57:LEU:HD22	1.74	0.68
1:A:217[A]:ARG:NH1	6:A:545:HOH:O	2.29	0.65
1:A:47:HIS:HB3	6:A:668:HOH:O	1.97	0.64
1:A:110:LEU:HB3	1:A:114:ALA:HB2	1.81	0.61
1:A:44:ASN:HD22	1:A:44:ASN:C	2.08	0.61
1:A:79:LYS:CA	1:A:136:MET:HE1	2.31	0.60
1:A:47:HIS:CB	6:A:668:HOH:O	2.49	0.60
1:A:364:ASP:OD1	1:A:390[B]:ARG:HD3	2.02	0.60
1:A:33:PHE:CZ	1:A:45[A]:MET:HE1	2.37	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:356:LYS:HD3	1:A:357:SER:H	1.69	0.57
1:A:79:LYS:HB3	1:A:136:MET:HE1	1.86	0.57
1:A:79:LYS:HA	1:A:136:MET:HE1	1.85	0.56
1:A:44:ASN:HD22	1:A:46:ALA:H	1.53	0.54
1:A:348:LEU:O	1:A:349:VAL:C	2.51	0.53
1:A:25[B]:ILE:HD13	1:A:53:GLN:HB3	1.92	0.52
1:A:21:GLY:N	1:A:54[A]:GLN:HE22	2.06	0.51
1:A:171:SER:OG	1:A:220:ARG:NH2	2.41	0.50
1:A:269:ILE:HD11	1:A:294:ARG:CZ	2.41	0.50
1:A:110:LEU:CB	1:A:114:ALA:HB2	2.41	0.50
1:A:138:SER:O	1:A:142:GLU:HG3	2.11	0.49
1:A:377:GLN:NE2	1:A:377:GLN:HA	2.29	0.48
1:A:331:ASP:HA	1:A:347:GLY:H	1.78	0.48
1:A:359:LEU:CD1	1:A:386:ILE:HD11	2.43	0.48
1:A:304:ASN:HA	1:A:330:TYR:CD2	2.49	0.48
1:A:13:THR:HG21	1:A:118:ILE:HG12	1.97	0.47
1:A:184:ASN:HD22	1:A:214:PRO:HB2	1.79	0.47
1:A:32:MET:HE1	1:A:43:PHE:CE2	2.50	0.47
1:A:359:LEU:HD13	1:A:386:ILE:HD11	1.96	0.47
1:A:48:GLN:HB3	6:A:589:HOH:O	2.16	0.46
1:A:223:VAL:O	1:A:241:HIS:HE1	1.98	0.46
1:A:355:GLU:HG3	6:A:615:HOH:O	2.16	0.46
1:A:275:GLY:HA2	1:A:403:GLU:O	2.15	0.46
1:A:78:LEU:HD13	1:A:132:VAL:HG11	1.98	0.45
1:A:48:GLN:CB	6:A:589:HOH:O	2.65	0.44
1:A:262:ASP:HA	6:A:590:HOH:O	2.16	0.44
1:A:303:ARG:HE	1:A:332:GLN:HG2	1.82	0.44
1:A:232:PRO:O	3:A:405:FAD:H8A	2.18	0.44
1:A:21:GLY:O	1:A:25[A]:ILE:HG12	2.17	0.44
1:A:390[B]:ARG:NH2	6:A:731:HOH:O	2.40	0.44
1:A:184:ASN:ND2	1:A:214:PRO:HB2	2.32	0.44
1:A:5:LYS:HB3	1:A:5:LYS:HE3	1.24	0.44
1:A:74:LEU:O	1:A:77:VAL:HG22	2.18	0.43
1:A:307:VAL:HG12	1:A:403:GLU:HG3	1.99	0.43
1:A:27:CYS:SG	1:A:108:GLU:OE2	2.76	0.43
1:A:259:PHE:CE2	1:A:284:MET:HE1	2.53	0.43
1:A:271:LEU:HD23	1:A:366:TYR:HB2	2.00	0.43
1:A:91:LYS:NZ	1:A:93:GLU:OE2	2.52	0.43
1:A:145:ALA:HB2	1:A:152:LYS:HG3	2.01	0.43
1:A:334:LEU:O	1:A:335:PRO:C	2.62	0.43
1:A:217[A]:ARG:CZ	6:A:545:HOH:O	2.65	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:106:ILE:O	1:A:110:LEU:HB2	2.19	0.42
1:A:79:LYS:HA	1:A:136:MET:CE	2.50	0.42
1:A:272:ILE:O	1:A:367:ILE:HA	2.20	0.42
1:A:278:LEU:HD21	1:A:313:ARG:HB2	2.02	0.41
1:A:231:GLN:HE21	1:A:231:GLN:HB3	1.68	0.41
1:A:232:PRO:HG2	3:A:405:FAD:N7A	2.36	0.41
1:A:331:ASP:HA	1:A:347:GLY:CA	2.51	0.41
1:A:47:HIS:HB2	6:A:668:HOH:O	2.18	0.41
1:A:2:LEU:HD12	1:A:2:LEU:N	2.36	0.41
1:A:102:LEU:HD21	4:A:411:X89:C1B	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	407/403 (101%)	395 (97%)	12 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	340/334 (102%)	316 (93%)	24 (7%)	13	10

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	LYS
1	A	44	ASN
1	A	49	GLU
1	A	50	GLN
1	A	69	GLU
1	A	84	LYS
1	A	91	LYS
1	A	143	ARG
1	A	161	GLU
1	A	163	ARG
1	A	165	GLU
1	A	198	PRO
1	A	211	SER
1	A	217[A]	ARG
1	A	217[B]	ARG
1	A	278	LEU
1	A	285	LEU
1	A	316	GLU
1	A	319	LYS
1	A	341	ARG
1	A	342	ASP
1	A	352	LYS
1	A	356	LYS
1	A	360	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	4	GLN
1	A	44	ASN
1	A	48	GLN
1	A	52	GLN
1	A	55	GLN
1	A	184	ASN
1	A	231	GLN
1	A	241	HIS
1	A	245	ASN
1	A	290	GLN
1	A	353	GLN
1	A	377	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	A	404	4,1	50,50,50	2.25	13 (26%)	67,82,82	1.95	19 (28%)
3	FAD	A	405	-	58,58,58	1.61	10 (17%)	85,89,89	1.74	14 (16%)
5	PO4	A	406	-	4,4,4	1.44	0	6,6,6	1.63	2 (33%)
4	X89	A	411	2	27,27,27	1.35	4 (14%)	37,37,37	1.88	10 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	404	4,1	-	1/14/54/54	-
3	FAD	A	405	-	-	3/34/50/50	0/6/6/6
4	X89	A	411	2	-	1/13/13/13	0/3/3/3

All (27) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	404	HEM	C3D-C2D	7.24	1.52	1.36
3	A	405	FAD	PA-O3P	7.06	1.67	1.59
2	A	404	HEM	FE-NC	6.35	2.16	1.95
2	A	404	HEM	FE-NA	5.47	2.13	1.95
2	A	404	HEM	FE-ND	4.61	2.09	1.94
2	A	404	HEM	CMD-C2D	3.74	1.58	1.50
2	A	404	HEM	FE-NB	3.66	2.06	1.94
2	A	404	HEM	CAB-C3B	3.48	1.56	1.47
4	A	411	X89	C2B-CL2B	3.43	1.81	1.73
3	A	405	FAD	C2A-N3A	3.03	1.39	1.33
3	A	405	FAD	C2A-N1A	3.00	1.39	1.33
3	A	405	FAD	C1'-C2'	2.94	1.56	1.52
3	A	405	FAD	C4X-N5	2.86	1.36	1.30
4	A	411	X89	C1-N1	2.64	1.53	1.46
2	A	404	HEM	CMC-C2C	2.64	1.56	1.50
2	A	404	HEM	C1A-NA	-2.63	1.34	1.39
2	A	404	HEM	CAD-C3D	2.58	1.58	1.51
4	A	411	X89	C4A-CL4A	2.55	1.80	1.74
3	A	405	FAD	C1'-N10	2.35	1.53	1.47
2	A	404	HEM	C1B-NB	-2.30	1.36	1.40
2	A	404	HEM	C4A-NA	-2.28	1.35	1.39
3	A	405	FAD	C5X-N5	2.24	1.43	1.39
3	A	405	FAD	C8A-N7A	2.19	1.35	1.31
4	A	411	X89	C6A-C1A	-2.09	1.37	1.39
2	A	404	HEM	CBA-CGA	2.09	1.55	1.50
3	A	405	FAD	C10-N1	2.08	1.37	1.33
3	A	405	FAD	C5'-C4'	2.06	1.54	1.51

All (45) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	405	FAD	N3A-C2A-N1A	-7.28	117.56	128.58
2	A	404	HEM	C4D-ND-C1D	5.50	111.72	105.21
4	A	411	X89	O-CBA-C1A	-4.94	106.09	112.05
3	A	405	FAD	C5A-C4A-N3A	-4.67	120.28	126.72
2	A	404	HEM	C3B-C2B-C1B	4.60	109.86	106.41
4	A	411	X89	C3B-C2B-C1B	-4.44	118.54	122.41
2	A	404	HEM	CMB-C2B-C1B	-4.27	118.37	125.03
3	A	405	FAD	C2A-N3A-C4A	4.23	122.15	111.83
3	A	405	FAD	C4X-C10-N10	4.15	122.43	116.48
2	A	404	HEM	C4A-NA-C1A	3.94	112.24	105.82
2	A	404	HEM	CBA-CAA-C2A	-3.82	101.97	112.53
2	A	404	HEM	C4C-NC-C1C	3.81	112.03	105.82

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	404	HEM	C2A-C1A-NA	-3.80	105.94	110.15
3	A	405	FAD	N9A-C8A-N7A	-3.57	108.87	113.94
4	A	411	X89	C2A-C1A-CBA	-3.55	118.91	122.15
2	A	404	HEM	C1B-NB-C4B	3.34	109.16	105.21
3	A	405	FAD	N3A-C4A-N9A	3.29	132.77	127.17
3	A	405	FAD	C5A-N7A-C8A	3.17	108.43	103.45
4	A	411	X89	C6A-C1A-C2A	3.15	119.98	116.82
3	A	405	FAD	C5A-C6A-N6A	-3.02	115.80	123.29
4	A	411	X89	O-CBA-C1	2.92	110.93	106.24
2	A	404	HEM	C3A-C4A-NA	-2.79	105.66	110.14
3	A	405	FAD	C10-C4X-N5	-2.66	119.38	124.81
2	A	404	HEM	O2D-CGD-CBD	2.61	122.23	114.00
5	A	406	PO4	O4-P-O3	-2.54	100.01	107.91
4	A	411	X89	CBB-C1B-C2B	-2.52	117.52	121.27
2	A	404	HEM	C4C-C3C-C2C	2.48	108.96	106.81
4	A	411	X89	C6B-C1B-C2B	2.45	120.77	116.91
4	A	411	X89	C3B-C4B-CL4B	-2.44	116.11	119.17
2	A	404	HEM	CHD-C4C-C3C	2.39	129.23	125.21
5	A	406	PO4	O3-P-O2	2.39	115.33	107.91
3	A	405	FAD	C2B-C1B-N9A	2.35	119.15	113.30
3	A	405	FAD	C9A-C5X-N5	-2.35	119.96	122.45
3	A	405	FAD	C4-C4X-C10	2.35	120.95	116.93
4	A	411	X89	C2A-C3A-C4A	-2.33	116.08	118.72
3	A	405	FAD	C6-C5X-C9A	2.27	122.16	119.05
2	A	404	HEM	C2C-C1C-NC	-2.27	105.44	109.64
2	A	404	HEM	CHB-C1B-NB	2.24	127.14	124.37
2	A	404	HEM	O2A-CGA-CBA	2.18	120.87	114.00
2	A	404	HEM	O2D-CGD-O1D	-2.15	117.79	123.33
2	A	404	HEM	C2B-C1B-NB	-2.11	107.41	109.84
3	A	405	FAD	O2B-C2B-C1B	2.11	117.36	110.10
2	A	404	HEM	C3B-C4B-NB	-2.05	108.00	109.47
4	A	411	X89	C1A-C2A-CL2A	-2.03	118.34	120.41
2	A	404	HEM	CHA-C4D-ND	2.01	126.85	124.37

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	405	FAD	C2B-C1B-N9A-C8A
3	A	405	FAD	C2B-C1B-N9A-C4A
4	A	411	X89	N1-C1-CBA-C1A
2	A	404	HEM	C3D-CAD-CBD-CGD

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	A	405	FAD	PA-O3P-P-O2P


There are no ring outliers.


4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	404	HEM	1	0
3	A	405	FAD	2	0
5	A	406	PO4	1	0
4	A	411	X89	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	403/403 (100%)	0.41	40 (9%)	13 11	11, 29, 61, 88	6 (1%)

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	338	VAL	5.7
1	A	335	PRO	4.8
1	A	228	GLY	4.3
1	A	345	TYR	4.3
1	A	47	HIS	4.3
1	A	230	PRO	4.3
1	A	334	LEU	3.9
1	A	333	PRO	3.6
1	A	306	ALA	3.3
1	A	340	GLY	3.3
1	A	303	ARG	3.2
1	A	232	PRO	3.2
1	A	49	GLU	3.0
1	A	336	GLU	3.0
1	A	229	GLY	2.9
1	A	318	ALA	2.8
1	A	279	THR	2.7
1	A	308	HIS	2.7
1	A	330	TYR	2.6
1	A	114	ALA	2.6
1	A	52	GLN	2.6
1	A	320	THR	2.5
1	A	50	GLN	2.5
1	A	227	GLY	2.4
1	A	356	LYS	2.4
1	A	339	GLN	2.4
1	A	343	TYR	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	302	ALA	2.4
1	A	307	VAL	2.4
1	A	346	PRO	2.4
1	A	344	ASP	2.3
1	A	48	GLN	2.2
1	A	380	ALA	2.2
1	A	231	GLN	2.1
1	A	341	ARG	2.1
1	A	337	ASP	2.1
1	A	163	ARG	2.1
1	A	347	GLY	2.1
1	A	310	MET	2.0
1	A	352	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	PO4	A	406	5/5	0.93	0.14	46,47,52,57	0
4	X89	A	411	25/25	0.95	0.09	26,33,41,55	0
3	FAD	A	405	53/53	0.97	0.09	15,22,74,81	0
2	HEM	A	404	43/43	0.98	0.08	17,18,34,44	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around X89 A 411:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FAD A 405:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM A 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.