



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 05:05 PM UTC

PDB ID : 2P0U / pdb_00002p0u
Title : crystal structure of Marchantia polymorpha stilbenecarboxylate synthase 2 (STCS2)
Authors : Borel, F.; Schroeder, G.; Schroeder, J.; Ferrer, J.-L.
Deposited on : 2007-03-01
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	413	81% 15% ..
1	B	413	81% 15% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	801	-	X	-	-
3	GOL	B	802	-	X	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6667 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Stilbenecarboxylate synthase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	403	Total	C	N	O	S	0	0	0
			3055	1932	540	567	16			
1	B	399	Total	C	N	O	S	0	0	0
			3028	1916	536	560	16			

There are 46 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	cloning artifact	UNP Q5I6Y1
A	2	LYS	-	cloning artifact	UNP Q5I6Y1
A	3	HIS	-	expression tag	UNP Q5I6Y1
A	4	HIS	-	expression tag	UNP Q5I6Y1
A	5	HIS	-	expression tag	UNP Q5I6Y1
A	6	HIS	-	expression tag	UNP Q5I6Y1
A	7	HIS	-	expression tag	UNP Q5I6Y1
A	8	HIS	-	expression tag	UNP Q5I6Y1
A	9	HIS	-	expression tag	UNP Q5I6Y1
A	10	HIS	-	expression tag	UNP Q5I6Y1
A	11	GLY	-	cloning artifact	UNP Q5I6Y1
A	12	GLY	-	cloning artifact	UNP Q5I6Y1
A	13	LEU	-	cloning artifact	UNP Q5I6Y1
A	14	VAL	-	cloning artifact	UNP Q5I6Y1
A	15	PRO	-	cloning artifact	UNP Q5I6Y1
A	16	ARG	-	cloning artifact	UNP Q5I6Y1
A	17	GLY	-	cloning artifact	UNP Q5I6Y1
A	18	SER	-	cloning artifact	UNP Q5I6Y1
A	19	HIS	-	cloning artifact	UNP Q5I6Y1
A	20	GLY	-	cloning artifact	UNP Q5I6Y1
A	21	GLY	-	cloning artifact	UNP Q5I6Y1
A	22	SER	-	cloning artifact	UNP Q5I6Y1
A	182	CSD	CYS	modified residue	UNP Q5I6Y1
B	1	MET	-	cloning artifact	UNP Q5I6Y1
B	2	LYS	-	cloning artifact	UNP Q5I6Y1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	3	HIS	-	expression tag	UNP Q5I6Y1
B	4	HIS	-	expression tag	UNP Q5I6Y1
B	5	HIS	-	expression tag	UNP Q5I6Y1
B	6	HIS	-	expression tag	UNP Q5I6Y1
B	7	HIS	-	expression tag	UNP Q5I6Y1
B	8	HIS	-	expression tag	UNP Q5I6Y1
B	9	HIS	-	expression tag	UNP Q5I6Y1
B	10	HIS	-	expression tag	UNP Q5I6Y1
B	11	GLY	-	cloning artifact	UNP Q5I6Y1
B	12	GLY	-	cloning artifact	UNP Q5I6Y1
B	13	LEU	-	cloning artifact	UNP Q5I6Y1
B	14	VAL	-	cloning artifact	UNP Q5I6Y1
B	15	PRO	-	cloning artifact	UNP Q5I6Y1
B	16	ARG	-	cloning artifact	UNP Q5I6Y1
B	17	GLY	-	cloning artifact	UNP Q5I6Y1
B	18	SER	-	cloning artifact	UNP Q5I6Y1
B	19	HIS	-	cloning artifact	UNP Q5I6Y1
B	20	GLY	-	cloning artifact	UNP Q5I6Y1
B	21	GLY	-	cloning artifact	UNP Q5I6Y1
B	22	SER	-	cloning artifact	UNP Q5I6Y1
B	182	CSD	CYS	modified residue	UNP Q5I6Y1

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is NICKEL (II) ION (CCD ID: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Ni	0	0
			1	1		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	287	Total	O	0	0
			287	287		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	264	Total 264	O 264	0	0

- Molecule 1: Stilbenecarboxylate synthase 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	129.27Å 133.20Å 53.49Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.99 – 1.90 41.99 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.0 (41.99-1.90) 99.1 (41.99-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	8.40	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.88 (at 1.89Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.188 , 0.217 0.185 , 0.213	Depositor DCC
R_{free} test set	3651 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	20.3	Xtriage
Anisotropy	0.527	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 41.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.019 for k,h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6667	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NI, CSD, SO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.35	0/3110	0.91	8/4209 (0.2%)
1	B	0.34	0/3083	0.91	11/4174 (0.3%)
All	All	0.34	0/6193	0.91	19/8383 (0.2%)

There are no bond length outliers.

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	394	GLY	N-CA-C	-8.27	104.92	115.42
1	A	394	GLY	N-CA-C	-8.16	103.55	115.30
1	A	89	TYR	N-CA-C	-6.75	103.39	111.69
1	B	89	TYR	N-CA-C	-6.74	103.74	112.23
1	B	29	ALA	N-CA-C	6.19	118.70	110.53

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3055	0	3069	56	0
1	B	3028	0	3021	43	0
2	A	10	0	0	0	0
2	B	10	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	6	0	4	0	0
3	B	6	0	4	0	0
4	B	1	0	0	0	0
5	A	287	0	0	2	0
5	B	264	0	0	1	0
All	All	6667	0	6098	95	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 95 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:117:VAL:HG22	1:A:154:ASN:HD21	0.99	1.12
1:A:117:VAL:HG22	1:A:154:ASN:ND2	1.77	0.98
1:A:113:GLN:O	1:A:117:VAL:HG23	1.71	0.91
1:A:252:ARG:HB3	1:A:252:ARG:NH1	1.95	0.81
1:B:246:PRO:HG2	1:B:253:PRO:HG3	1.60	0.80

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	400/413 (97%)	392 (98%)	8 (2%)	0	100	100
1	B	394/413 (95%)	382 (97%)	11 (3%)	1 (0%)	36	29
All	All	794/826 (96%)	774 (98%)	19 (2%)	1 (0%)	48	40

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	23	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	320/330 (97%)	316 (99%)	4 (1%)	61	61
1	B	315/330 (96%)	312 (99%)	3 (1%)	68	70
All	All	635/660 (96%)	628 (99%)	7 (1%)	65	67

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	395	LEU
1	B	247	ARG
1	B	395	LEU
1	B	252	ARG
1	A	328	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	180	GLN
1	B	92	GLN
1	A	223	HIS
1	A	180	GLN
1	B	7	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CSD	B	182	1	4,7,8	2.12	1 (25%)	1,8,10	6.09	1 (100%)
1	CSD	A	182	1	4,7,8	1.92	1 (25%)	1,8,10	5.62	1 (100%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSD	B	182	1	-	1/2/6/8	-
1	CSD	A	182	1	-	1/2/6/8	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	182	CSD	OD1-SG	-4.06	1.44	1.47
1	A	182	CSD	OD1-SG	-3.72	1.44	1.47

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	182	CSD	OD1-SG-CB	6.09	116.82	105.60
1	A	182	CSD	OD1-SG-CB	5.62	115.95	105.60

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	182	CSD	CA-CB-SG-OD1
1	B	182	CSD	CA-CB-SG-OD1

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	801	-	5,5,5	4.77	5 (100%)	5,5,5	6.06	3 (60%)
2	SO4	A	702	-	4,4,4	0.37	0	6,6,6	0.08	0
2	SO4	B	703	-	4,4,4	0.38	0	6,6,6	0.13	0
2	SO4	B	704	-	4,4,4	0.37	0	6,6,6	0.09	0
3	GOL	B	802	-	5,5,5	4.78	5 (100%)	5,5,5	6.07	3 (60%)
2	SO4	A	701	-	4,4,4	0.38	0	6,6,6	0.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	802	-	-	3/4/4/4	-
3	GOL	A	801	-	-	3/4/4/4	-

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	802	GOL	C3-C2	-8.03	1.21	1.51
3	A	801	GOL	C3-C2	-8.00	1.21	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	802	GOL	O1-C1	4.54	1.61	1.42
3	A	801	GOL	O1-C1	4.53	1.61	1.42
3	B	802	GOL	O3-C3	3.44	1.56	1.42

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	802	GOL	O3-C3-C2	11.10	160.35	110.38
3	A	801	GOL	O3-C3-C2	11.09	160.32	110.38
3	B	802	GOL	O2-C2-C3	6.93	137.89	109.18
3	A	801	GOL	O2-C2-C3	6.92	137.84	109.18
3	B	802	GOL	O1-C1-C2	3.44	125.87	110.38

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	801	GOL	C1-C2-C3-O3
3	A	801	GOL	O1-C1-C2-C3
3	B	802	GOL	O1-C1-C2-C3
3	A	801	GOL	O2-C2-C3-O3
3	B	802	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	402/413 (97%)	-0.10	6 (1%) 72 75	12, 20, 33, 48	0
1	B	398/413 (96%)	0.01	16 (4%) 42 45	11, 21, 36, 64	0
All	All	800/826 (96%)	-0.05	22 (2%) 55 59	11, 21, 34, 64	0

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	394	GLY	7.1
1	A	395	LEU	6.4
1	B	395	LEU	5.6
1	B	23	SER	5.5
1	B	394	GLY	5.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSD	A	182	8/9	0.94	0.09	22,25,31,32	0
1	CSD	B	182	8/9	0.95	0.09	21,23,29,30	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	B	703	5/5	0.56	0.26	67,67,69,70	0
2	SO4	A	702	5/5	0.64	0.23	70,72,73,74	0
2	SO4	B	704	5/5	0.64	0.21	80,80,81,82	0
3	GOL	B	802	6/6	0.72	0.20	50,53,54,54	0
3	GOL	A	801	6/6	0.78	0.18	52,56,56,57	0
2	SO4	A	701	5/5	0.78	0.19	83,83,83,84	0
4	NI	B	601	1/1	0.99	0.04	28,28,28,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.