



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 09:29 PM UTC

PDB ID : 7P01 / pdb_00007p01
Title : Structure of the maltase BaAG2 from Blastobotrys adeninivorans in complex with acarbose
Authors : Ernits, K.; Visnapuu, T.; Persson, K.
Deposited on : 2021-06-29
Resolution : 2.12 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	8290 (2.14-2.10)
Clashscore	190562	8817 (2.14-2.10)
Ramachandran outliers	187476	8738 (2.14-2.10)
Sidechain outliers	187428	8739 (2.14-2.10)
RSRZ outliers	180081	8294 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	585	91% 6% .
1	B	585	2% 84% 13% ..
2	D	3	33% 67%
2	E	3	100%
2	F	3	67% 33%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 19262 atoms, of which 9085 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called BaAG2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	572	Total	C	H	N	O	S	0	1	0
			9143	2978	4466	792	895	12			
1	B	571	Total	C	H	N	O	S	0	0	0
			9131	2973	4463	790	893	12			

- Molecule 2 is an oligosaccharide called 4,6-dideoxy-4-[(1S,4R,5S,6S)-4,5,6-trihydroxy-3-(hydroxymethyl)cyclohex-2-en-1-yl]amino}-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	D	3	Total	C	H	N	O		0	0	0
			87	25	43	1	18				
2	E	3	Total	C	H	N	O		0	0	0
			87	25	43	1	18				
2	F	3	Total	C	H	N	O		0	0	0
			87	25	43	1	18				

- Molecule 3 is 1-methylpyrrolidin-2-one (CCD ID: MB3) (formula: C₅H₉NO).




- | Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|-----|-------|----------|-----------------|---------|---------|
| 4 | A | 1 | Total Ca
1 1 | 0 | 0 |
| 4 | B | 1 | Total Ca
1 1 | 0 | 0 |

- | Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|-----|-------|----------|--------------------|---------|---------|
| 5 | A | 443 | Total O
443 443 | 0 | 0 |
| 5 | B | 234 | Total O
234 234 | 0 | 0 |


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: BaAG2



• Molecule 1: BaAG2



• Molecule 2: 4,6-dideoxy-4-[(1S,4R,5S,6S)-4,5,6-trihydroxy-3-(hydroxymethyl)cyclohex-2-en-1-yl]amino}-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



• Molecule 2: 4,6-dideoxy-4-[(1S,4R,5S,6S)-4,5,6-trihydroxy-3-(hydroxymethyl)cyclohex-2-en-1-yl]amino}-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



GLC1
GLC2
AC13

- Molecule 2: 4,6-dideoxy-4-{[(1S,4R,5S,6S)-4,5,6-trihydroxy-3-(hydroxymethyl)cyclohex-2-en-1-yl]amino}-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain F:

67%

33%

GLC1
GLC2
AC13

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	68.03Å 78.08Å 121.93Å 90.00° 94.10° 90.00°	Depositor
Resolution (Å)	46.32 – 2.12 46.32 – 2.12	Depositor EDS
% Data completeness (in resolution range)	98.9 (46.32-2.12) 99.9 (46.32-2.12)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.23	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.37 (at 2.12Å)	Xtriage
Refinement program	PHENIX 1.17.1_3660, PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.184 , 0.195 0.186 , 0.198	Depositor DCC
R_{free} test set	2027 reflections (2.80%)	wwPDB-VP
Wilson B-factor (Å ²)	33.3	Xtriage
Anisotropy	0.170	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 39.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	19262	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MB3, GLC, CA, AC1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.29	21/4812 (0.4%)	1.31	7/6529 (0.1%)
1	B	1.15	8/4798 (0.2%)	1.34	14/6510 (0.2%)
All	All	1.22	29/9610 (0.3%)	1.33	21/13039 (0.2%)

The worst 5 of 29 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	451[A]	ALA	N-CA	6.59	1.54	1.46
1	A	451[B]	ALA	N-CA	6.59	1.54	1.46
1	A	471	PRO	C-O	-6.55	1.16	1.24
1	A	284	LEU	C-O	-6.36	1.16	1.24
1	B	435	LEU	C-O	-6.36	1.16	1.24

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	330	GLN	CA-C-N	6.87	130.40	120.38
1	B	330	GLN	C-N-CA	6.87	130.40	120.38
1	B	272	VAL	CA-C-N	6.82	129.75	122.38
1	B	272	VAL	C-N-CA	6.82	129.75	122.38
1	A	331	ARG	CB-CA-C	-6.74	99.22	110.68

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4677	4466	4465	4	1
1	B	4668	4463	4462	32	1
2	D	44	43	30	1	0
2	E	44	43	29	0	0
2	F	44	43	30	2	0
3	A	14	18	18	0	0
3	B	7	9	9	1	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	443	0	0	0	0
5	B	234	0	0	0	0
All	All	10177	9085	9043	38	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 38 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:256:LEU:HD23	1:B:259:MET:HE3	1.69	0.73
1:B:256:LEU:HD23	1:B:259:MET:CE	2.19	0.72
1:B:146:GLU:OE2	1:B:146:GLU:HA	1.94	0.68
1:B:182:GLU:N	1:B:182:GLU:OE1	2.32	0.63
1:B:255:ILE:HG22	1:B:259:MET:HE2	1.80	0.63

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:150:ARG:HE	1:B:149:ASN:HD22[2_454]	1.18	0.42

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	571/585 (98%)	556 (97%)	15 (3%)	0	100	100
1	B	569/585 (97%)	541 (95%)	27 (5%)	1 (0%)	43	44
All	All	1140/1170 (97%)	1097 (96%)	42 (4%)	1 (0%)	48	49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	12	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	508/520 (98%)	500 (98%)	8 (2%)	55	63
1	B	507/520 (98%)	478 (94%)	29 (6%)	18	16
All	All	1015/1040 (98%)	978 (96%)	37 (4%)	31	33

5 of 37 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	433	LYS
1	B	551	SER
1	B	438	VAL
1	B	470	TYR
1	B	97	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	548	GLN
1	B	24	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	237	GLN
1	B	149	ASN
1	A	491	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

9 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLC	D	1	2	12,12,12	1.03	0	17,17,17	1.09	1 (5%)
2	GLC	D	2	2	11,11,12	1.27	2 (18%)	15,15,17	1.65	4 (26%)
2	AC1	D	3	2	21,22,23	1.45	3 (14%)	22,32,34	1.73	5 (22%)
2	GLC	E	1	2	12,12,12	1.11	0	17,17,17	2.33	8 (47%)
2	GLC	E	2	2	11,11,12	1.26	2 (18%)	15,15,17	1.00	0
2	AC1	E	3	2	21,22,23	6.44	8 (38%)	22,32,34	1.46	5 (22%)
2	GLC	F	1	2	12,12,12	0.73	0	17,17,17	1.36	3 (17%)
2	GLC	F	2	2	11,11,12	1.04	1 (9%)	15,15,17	1.32	2 (13%)
2	AC1	F	3	2	21,22,23	1.33	1 (4%)	22,32,34	1.61	5 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	D	1	2	-	0/2/22/22	0/1/1/1
2	GLC	D	2	2	-	0/2/19/22	0/1/1/1
2	AC1	D	3	2	-	1/6/43/46	0/2/2/2
2	GLC	E	1	2	-	1/2/22/22	0/1/1/1
2	GLC	E	2	2	-	2/2/19/22	0/1/1/1
2	AC1	E	3	2	-	4/6/43/46	0/2/2/2
2	GLC	F	1	2	-	0/2/22/22	0/1/1/1
2	GLC	F	2	2	-	0/2/19/22	0/1/1/1
2	AC1	F	3	2	-	1/6/43/46	0/2/2/2

The worst 5 of 17 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	3	AC1	C7B-C5B	22.00	1.63	1.32
2	E	3	AC1	C4A-C5B	-11.10	1.42	1.51
2	E	3	AC1	C2B-C1B	9.40	1.65	1.53
2	E	3	AC1	C3B-C4A	8.23	1.64	1.53
2	E	3	AC1	C1B-C7B	-6.52	1.41	1.50

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	1	GLC	O5-C1-C2	5.26	119.55	110.30
2	D	3	AC1	C2B-C3B-C4A	-4.77	102.67	110.22
2	E	1	GLC	O5-C5-C4	4.26	117.38	109.70
2	F	3	AC1	O6B-C6B-C5B	-3.26	102.48	112.74
2	D	3	AC1	O2B-C2B-C1B	3.12	115.22	109.08

There are no chirality outliers.

5 of 9 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	D	3	AC1	C7B-C1B-N4A-C4
2	E	3	AC1	C7B-C5B-C6B-O6B
2	F	3	AC1	C7B-C1B-N4A-C4
2	E	2	GLC	C4-C5-C6-O6
2	E	1	GLC	C4-C5-C6-O6


There are no ring outliers.


3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	1	GLC	2	0
2	D	1	GLC	1	0
2	D	2	GLC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	MB3	A	902	-	7,7,7	2.32	4 (57%)	9,9,9	1.31	1 (11%)
3	MB3	A	901	-	7,7,7	2.67	4 (57%)	9,9,9	1.25	1 (11%)
3	MB3	B	701	-	7,7,7	2.34	4 (57%)	9,9,9	2.06	3 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MB3	A	902	-	-	-	0/1/1/1
3	MB3	A	901	-	-	-	0/1/1/1
3	MB3	B	701	-	-	-	0/1/1/1

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	901	MB3	OAB-CAF	-4.48	1.13	1.23
3	A	902	MB3	CAC-CAE	3.86	1.65	1.51
3	A	902	MB3	CAF-NAG	3.49	1.51	1.31
3	B	701	MB3	OAB-CAF	-3.42	1.15	1.23
3	A	901	MB3	CAC-CAE	3.35	1.63	1.51

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	701	MB3	CAE-CAC-CAD	-3.47	95.43	105.94
3	B	701	MB3	CAC-CAE-NAG	-3.43	90.94	102.99
3	B	701	MB3	CAC-CAD-CAF	-2.92	95.73	104.41
3	A	902	MB3	CAE-CAC-CAD	-2.61	98.04	105.94
3	A	901	MB3	CAE-CAC-CAD	-2.19	99.32	105.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	701	MB3	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	572/585 (97%)	-0.39	2 (0%) 90 91	25, 38, 63, 120	0
1	B	571/585 (97%)	0.06	9 (1%) 70 73	32, 54, 94, 141	0
All	All	1143/1170 (97%)	-0.16	11 (0%) 79 82	25, 45, 82, 141	0

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	310	ILE	4.6
1	A	581	HIS	3.7
1	B	148	GLY	3.2
1	B	230	ALA	2.9
1	B	236	GLY	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLC	E	1	12/12	0.77	0.12	29,35,44,47	0
2	AC1	D	3	21/22	0.91	0.09	25,34,43,45	0
2	GLC	E	2	11/12	0.92	0.08	31,34,40,43	0
2	AC1	E	3	21/22	0.93	0.08	36,57,77,78	0
2	GLC	D	1	12/12	0.94	0.07	43,48,58,60	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GLC	D	2	11/12	0.95	0.05	36,41,49,50	0
2	GLC	F	1	12/12	-	-	61,68,82,83	0
2	GLC	F	2	11/12	-	-	48,55,65,68	0
2	AC1	F	3	21/22	-	-	40,45,52,54	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain F:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MB3	A	902	7/7	0.64	0.20	71,86,86,86	0
3	MB3	A	901	7/7	0.87	0.13	44,55,60,60	0
3	MB3	B	701	7/7	0.88	0.14	59,72,73,73	0
4	CA	B	702	1/1	0.99	0.03	44,44,44,44	0
4	CA	A	903	1/1	1.00	0.02	31,31,31,31	0

6.5 Other polymers [i](#)

There are no such residues in this entry.