



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 01:07 PM UTC

PDB ID : 3P1U / pdb_00003p1u
Title : Crystal structure of a SusD homolog (BDI_0600) from Parabacteroides distasonis ATCC 8503 AT 2.05 Å resolution
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2010-09-30
Resolution : 2.05 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2260 (2.04-2.04)
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)
RSRZ outliers	180081	2260 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	529	18% 88% 9% .
1	B	529	8% 90% 7% .

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 8729 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SusD homolog.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	515	Total	C	N	O	S	Se	0	7	0
			4067	2588	686	782	4	7			
1	B	515	Total	C	N	O	S	Se	0	5	0
			4065	2583	681	790	4	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP A6L9L2
B	0	GLY	-	expression tag	UNP A6L9L2

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	298	Total 298	O 298	0	0
3	B	294	Total 294	O 294	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: SusD homolog



• Molecule 1: SusD homolog



4 Data and refinement statistics

Property	Value	Source
Space group	P 6	Depositor
Cell constants a, b, c, α , β , γ	130.24Å 130.24Å 114.82Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	28.70 – 2.05 28.70 – 2.05	Depositor EDS
% Data completeness (in resolution range)	(Not available) (28.70-2.05) 100.0 (28.70-2.05)	Depositor EDS
R_{merge}	0.25	Depositor
R_{sym}	0.25	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.96 (at 2.04Å)	Xtriage
Refinement program	BUSTER-TNT BUSTER 2.8.0, BUSTER 2.8.0	Depositor
R, R_{free}	0.243 , 0.265 0.272 , 0.298	Depositor DCC
R_{free} test set	3503 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	16.4	Xtriage
Anisotropy	0.364	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 43.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.042 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	8729	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.89	0/4174	0.72	0/5651
1	B	0.87	0/4168	0.71	0/5646
All	All	0.88	0/8342	0.71	0/11297

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4067	0	3943	29	0
1	B	4065	0	3925	22	0
2	B	5	0	0	0	0
3	A	298	0	0	2	0
3	B	294	0	0	0	0
All	All	8729	0	7868	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:41:ASN:HD22	1:A:44[B]:ARG:HH12	1.30	0.76
1:A:533:GLY:HA2	3:A:676:HOH:O	1.87	0.73
1:B:64:LYS:HB3	1:B:65:PRO:CD	2.22	0.70
1:A:64:LYS:HB3	1:A:65:PRO:CD	2.23	0.68
1:A:105:ALA:O	1:A:109:ILE:HG12	2.03	0.58
1:A:207:LYS:HB3	1:A:241:ALA:HB1	1.88	0.55
1:B:256:LYS:HB2	1:B:263:GLN:HB3	1.89	0.55
1:B:207:LYS:HB3	1:B:241:ALA:HB1	1.87	0.55
1:A:256:LYS:HB2	1:A:263:GLN:HB3	1.88	0.54
1:B:295:ALA:HA	1:B:316[A]:LYS:HD3	1.90	0.53
1:B:64:LYS:HB3	1:B:65:PRO:HD3	1.90	0.53
1:A:64:LYS:HB3	1:A:65:PRO:HD3	1.92	0.52
1:A:41:ASN:HB3	1:A:110:VAL:HG13	1.92	0.51
1:B:211:ASN:HD21	1:B:239:SER:HB3	1.76	0.50
1:B:134:VAL:O	1:B:138:THR:HG23	2.13	0.49
1:A:131[B]:ILE:HD11	1:A:178:ALA:HB3	1.94	0.49
1:A:134:VAL:O	1:A:138:THR:HG23	2.13	0.48
1:A:211:ASN:HD21	1:A:239:SER:HB3	1.78	0.48
1:A:198:ILE:HD12	1:A:206:TRP:HH2	1.79	0.47
1:A:293:GLY:HA2	1:A:316:LYS:HD3	1.96	0.47
1:B:219:VAL:O	1:B:222:THR:HG23	2.14	0.47
1:B:505:ILE:HG13	1:B:536:VAL:HB	1.97	0.47
1:A:219:VAL:O	1:A:222:THR:HG23	2.14	0.47
1:A:505:ILE:HG13	1:A:536:VAL:HB	1.97	0.47
1:B:198:ILE:HD12	1:B:206:TRP:HH2	1.80	0.47
1:B:290:ASN:HA	1:B:316[A]:LYS:HB2	1.98	0.46
1:B:301:VAL:HG12	1:B:308:ALA:HA	1.97	0.46
1:A:64:LYS:HB3	1:A:65:PRO:HD2	1.99	0.44
1:B:64:LYS:HB3	1:B:65:PRO:HD2	1.98	0.44
1:B:408:ARG:HA	1:B:420:VAL:HG21	1.99	0.44
1:A:453:ILE:HG23	3:A:605:HOH:O	2.17	0.43
1:A:74:MSE:SE	1:A:465:ALA:HA	2.69	0.43
1:B:88:GLN:HG3	1:B:509:ARG:HG3	1.99	0.43
1:B:321:ALA:HB3	1:B:322:PRO:HD3	2.00	0.43
1:A:321:ALA:HB3	1:A:322:PRO:HD3	2.00	0.42
1:A:62:ARG:HG2	1:A:92:TRP:CE2	2.55	0.42
1:A:74:MSE:HE3	1:A:468:GLY:HA3	2.02	0.42
1:A:341:PRO:HG3	1:A:351:TYR:CD1	2.55	0.42
1:B:64:LYS:CB	1:B:65:PRO:CD	2.94	0.42
1:A:211:ASN:ND2	1:A:236:ALA:O	2.53	0.42
1:B:62:ARG:HG2	1:B:92:TRP:CE2	2.55	0.42
1:A:64:LYS:CB	1:A:65:PRO:CD	2.95	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:75:THR:HB	1:B:278:ILE:HG22	2.02	0.41
1:B:341:PRO:HG3	1:B:351:TYR:CD1	2.55	0.41
1:A:198:ILE:HD12	1:A:206:TRP:CH2	2.54	0.41
1:A:380:GLU:HG3	1:A:409:ALA:HB3	2.03	0.41
1:A:75:THR:HB	1:A:278:ILE:HG22	2.03	0.41
1:A:461:ILE:HA	1:A:462:PRO:HA	1.89	0.41
1:B:74:MSE:HE3	1:B:468:GLY:HA3	2.02	0.40
1:A:161:TYR:CD1	1:A:521:GLU:HB3	2.56	0.40
1:B:198:ILE:HD12	1:B:206:TRP:CH2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	520/529 (98%)	496 (95%)	21 (4%)	3 (1%)	21	13
1	B	518/529 (98%)	494 (95%)	21 (4%)	3 (1%)	21	13
All	All	1038/1058 (98%)	990 (95%)	42 (4%)	6 (1%)	21	13

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	64	LYS
1	A	487	ASP
1	B	64	LYS
1	B	487	ASP
1	A	54	ILE
1	B	54	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	422/435 (97%)	420 (100%)	2 (0%)	81	85
1	B	425/435 (98%)	423 (100%)	2 (0%)	81	85
All	All	847/870 (97%)	843 (100%)	4 (0%)	81	85

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	56	GLU
1	A	513	LYS
1	B	56	GLU
1	B	513	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	41	ASN
1	A	67	GLN
1	A	158	ASN
1	A	545	ASN
1	B	67	GLN
1	B	94	ASN
1	B	158	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	B	554	-	4,4,4	0.19	0	6,6,6	0.08	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	507/529 (95%)	1.34	94 (18%) 3 2	10, 18, 32, 51	7 (1%)
1	B	507/529 (95%)	0.83	42 (8%) 17 17	8, 16, 30, 48	5 (0%)
All	All	1014/1058 (95%)	1.08	136 (13%) 7 6	8, 17, 32, 51	12 (1%)

All (136) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	120[A]	ASP	7.2
1	A	118	ASP	5.9
1	B	116	GLU	5.1
1	A	117	LYS	4.9
1	A	121	ALA	4.8
1	A	122	TYR	4.8
1	A	155	VAL	4.8
1	B	157	ALA	4.6
1	A	119	LYS	4.3
1	A	41	ASN	4.2
1	B	152	TYR	4.1
1	B	155	VAL	4.1
1	A	185	SER	4.0
1	B	115	THR	3.8
1	A	157	ALA	3.7
1	A	343	ALA	3.7
1	A	532[A]	LYS	3.6
1	A	39	ASN	3.6
1	B	154	ILE	3.6
1	A	154	ILE	3.6
1	A	123	ALA	3.6
1	A	189	ILE	3.5
1	B	316[A]	LYS	3.5
1	A	110	VAL	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	399	ALA	3.4
1	A	211	ASN	3.4
1	B	176[A]	ASP	3.4
1	A	115	THR	3.3
1	A	114	LEU	3.3
1	B	109	ILE	3.3
1	A	187	GLN	3.2
1	A	236	ALA	3.1
1	B	114	LEU	3.1
1	B	111	ILE	3.1
1	B	122	TYR	3.0
1	A	44[A]	ARG	3.0
1	A	509	ARG	3.0
1	A	109	ILE	3.0
1	B	126	ILE	3.0
1	B	39	ASN	3.0
1	A	405	ALA	3.0
1	A	237	ILE	3.0
1	A	159	PRO	3.0
1	A	116	GLU	2.9
1	B	41	ASN	2.9
1	A	153	GLU	2.9
1	B	153	GLU	2.9
1	A	449	LEU	2.8
1	A	111	ILE	2.8
1	A	192	ASP	2.7
1	B	177	ALA	2.7
1	A	448	ALA	2.7
1	A	150	PRO	2.7
1	A	158	ASN	2.7
1	A	541	TRP	2.7
1	B	112	ARG	2.7
1	B	180	ASN	2.6
1	A	171	TYR	2.6
1	A	178	ALA	2.6
1	A	190	PHE	2.6
1	A	169	ASP	2.6
1	A	524[A]	LYS	2.6
1	A	152	TYR	2.6
1	A	160	PRO	2.6
1	A	106	SER	2.6
1	A	112	ARG	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	113	SER	2.6
1	B	408	ARG	2.6
1	A	198	ILE	2.6
1	B	156	GLU	2.6
1	B	349	GLY	2.6
1	B	128	PHE	2.5
1	A	203	VAL	2.5
1	A	453	ILE	2.5
1	B	184	ASP	2.5
1	B	439	ASP	2.5
1	A	197	LEU	2.5
1	A	516	LEU	2.5
1	A	53	LEU	2.5
1	A	70	PHE	2.5
1	A	350	ILE	2.5
1	A	452	ILE	2.5
1	A	245	SER	2.4
1	A	162	LYS	2.4
1	A	526	LYS	2.4
1	A	438	TYR	2.4
1	A	156	GLU	2.4
1	A	107	ILE	2.4
1	B	311[A]	SER	2.4
1	A	512	THR	2.4
1	A	373	PRO	2.4
1	A	186	ALA	2.3
1	A	54	ILE	2.3
1	A	309	VAL	2.3
1	A	502	ALA	2.3
1	B	516	LEU	2.3
1	A	182	PHE	2.3
1	B	158	ASN	2.3
1	A	354	LYS	2.3
1	B	247	LYS	2.3
1	A	299	GLY	2.2
1	A	433	GLY	2.2
1	B	121	ALA	2.2
1	A	42	GLU	2.2
1	A	105	ALA	2.2
1	A	129	ALA	2.2
1	A	233	ALA	2.2
1	A	475	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	282[A]	LYS	2.2
1	B	427	THR	2.2
1	A	173	ARG	2.2
1	B	443	GLY	2.1
1	A	151	ALA	2.1
1	A	144	PHE	2.1
1	B	363	ILE	2.1
1	A	500	LYS	2.1
1	B	119	LYS	2.1
1	A	476	ASN	2.1
1	B	485	TYR	2.1
1	B	299	GLY	2.1
1	A	206	TRP	2.1
1	B	484	VAL	2.1
1	A	45	LEU	2.1
1	B	245[A]	SER	2.1
1	A	183	ASN	2.1
1	A	43	PRO	2.1
1	A	104	ILE	2.1
1	A	230	ILE	2.1
1	A	377	PHE	2.1
1	A	241	ALA	2.1
1	B	185	SER	2.1
1	B	150	PRO	2.0
1	B	113[A]	SER	2.0
1	A	540	LEU	2.0
1	B	228	LYS	2.0
1	A	188	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SO4	B	554	5/5	0.91	0.18	35,39,40,40	0

6.5 Other polymers [i](#)

There are no such residues in this entry.