



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 11:20 AM UTC

PDB ID : 8P1P / pdb_00008p1p
Title : USP28 in complex with AZ1
Authors : Sauer, F.; Karal-Nair, R.; Kisker, C.
Deposited on : 2023-05-12
Resolution : 2.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1009 (2.76-2.76)
Clashscore	190562	1044 (2.76-2.76)
Ramachandran outliers	187476	1024 (2.76-2.76)
Sidechain outliers	187428	1024 (2.76-2.76)
RSRZ outliers	180081	1009 (2.76-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	494	8% 78% 12% 10%
1	B	494	10% 74% 12% 14%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7206 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ubiquitin carboxyl-terminal hydrolase 28.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	444	Total	C	N	O	S	0	3	0
			3654	2334	615	684	21			
1	B	425	Total	C	N	O	S	0	1	0
			3450	2213	573	644	20			


There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	148	GLY	-	expression tag	UNP Q96RU2
A	280	HIS	PRO	conflict	UNP Q96RU2
A	525	SER	-	linker	UNP Q96RU2
A	526	GLY	-	linker	UNP Q96RU2
A	527	SER	-	linker	UNP Q96RU2
A	528	GLY	-	linker	UNP Q96RU2
B	148	GLY	-	expression tag	UNP Q96RU2
B	280	HIS	PRO	conflict	UNP Q96RU2
B	525	SER	-	linker	UNP Q96RU2
B	526	GLY	-	linker	UNP Q96RU2
B	527	SER	-	linker	UNP Q96RU2
B	528	GLY	-	linker	UNP Q96RU2

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cl	0	0
			1	1		

- Molecule 3 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	O	S		0	0
			4	2	1	1			

- Molecule 4 is 2-[[5-bromanyl-2-[[4-fluoranyl-3-(trifluoromethyl)phenyl]methoxy]phenyl]methylamino]ethanol (CCD ID: WF0) (formula: $C_{17}H_{16}BrF_4NO_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total	Br	C	F	N	O	0	0
			25	1	17	4	1	2		
4	B	1	Total	Br	C	F	N	O	0	0
			25	1	17	4	1	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	33	Total 33	O 33	0	0
5	B	14	Total 14	O 14	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ubiquitin carboxyl-terminal hydrolase 28



- Molecule 1: Ubiquitin carboxyl-terminal hydrolase 28



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 2 21	Depositor
Cell constants a, b, c, α , β , γ	100.44Å 105.54Å 178.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.35 – 2.76 48.35 – 2.76	Depositor EDS
% Data completeness (in resolution range)	69.1 (48.35-2.76) 69.3 (48.35-2.76)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.25 (at 2.77Å)	Xtriage
Refinement program	PHENIX (1.19.2_4158: ???)	Depositor
R, R_{free}	0.213 , 0.241 0.209 , 0.236	Depositor DCC
R_{free} test set	1213 reflections (3.52%)	wwPDB-VP
Wilson B-factor (Å ²)	60.9	Xtriage
Anisotropy	0.192	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 62.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.023 for k,h,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	7206	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: WF0, DMS, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.13	0/3748	0.31	0/5075
1	B	0.13	0/3535	0.34	1/4792 (0.0%)
All	All	0.13	0/7283	0.32	1/9867 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	349	LYS	CB-CA-C	-5.44	109.83	117.23

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3654	0	3477	33	0
1	B	3450	0	3250	32	0
2	A	1	0	0	0	0
3	A	4	0	6	0	0
4	A	25	0	0	0	0
4	B	25	0	0	0	0
5	A	33	0	0	0	0
5	B	14	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	7206	0	6733	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:425:LYS:HG2	1:A:428:ARG:HH21	1.43	0.81
1:A:633:SER:HB3	1:A:642:ALA:HB2	1.72	0.70
1:B:295:THR:HG22	1:B:312:THR:HG22	1.72	0.70
1:B:427:GLU:HA	1:B:430:VAL:HG22	1.75	0.68
1:A:295:THR:HG22	1:A:312:THR:HG22	1.75	0.67
1:A:320:VAL:HG22	1:A:386:LEU:HD22	1.78	0.65
1:A:195:LEU:HG	1:A:199:VAL:HG21	1.82	0.61
1:A:212:ILE:O	1:A:216:GLN:HG2	2.03	0.59
1:B:212:ILE:O	1:B:216:GLN:HG2	2.03	0.59
1:B:633:SER:HB3	1:B:642:ALA:HB2	1.84	0.58
1:B:668:SER:HB3	1:B:671:GLU:HG2	1.86	0.57
1:B:334:MET:HE2	1:B:397:TYR:HB3	1.88	0.55
1:A:398:MET:HE2	1:A:582:VAL:HG21	1.88	0.55
1:B:289:VAL:HA	1:B:293:TYR:CD1	2.42	0.54
1:A:337:GLY:HA3	1:A:351:GLY:HA2	1.89	0.54
1:B:330:LEU:O	1:B:334:MET:HG2	2.07	0.54
1:B:161:PRO:HB2	1:B:621:VAL:HG11	1.89	0.54
1:B:275:VAL:HG21	1:B:286:ASN:HA	1.91	0.53
1:A:149:PRO:HG3	1:B:559:LEU:HD23	1.91	0.53
1:B:439:PHE:CG	1:B:440:PRO:HD2	2.44	0.52
1:A:533:VAL:HG13	1:A:537:GLU:HB3	1.92	0.52
1:B:286:ASN:HB3	1:B:289:VAL:HG23	1.91	0.52
1:A:590:LEU:HB2	1:A:644:CYS:HB3	1.91	0.51
1:A:330:LEU:O	1:A:334:MET:HG2	2.11	0.51
1:A:152:ASN:O	1:A:152:ASN:ND2	2.44	0.50
1:B:173:PHE:CZ	1:B:177:ILE:HG13	2.47	0.49
1:B:630:GLU:HG2	1:B:634:TYR:CZ	2.48	0.48
1:A:370:PHE:CZ	1:A:596:ALA:HB2	2.49	0.48
1:A:439:PHE:CG	1:A:440:PRO:HD2	2.48	0.48
1:A:630:GLU:HG2	1:A:634:TYR:CZ	2.47	0.48
1:A:156:ARG:HD2	1:A:160:TRP:O	2.14	0.48
1:A:286:ASN:HB3	1:A:289:VAL:HG23	1.96	0.47
1:B:180:LEU:HB3	1:B:186:PHE:CD2	2.49	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:440:PRO:HG2	1:A:443:ASP:HB2	1.96	0.47
1:A:321:ASN:HD21	1:A:382:ILE:HG21	1.79	0.47
1:B:404:LEU:HD11	1:B:408:LYS:HE3	1.98	0.46
1:B:408:LYS:HD3	1:B:573:MET:HG3	1.98	0.46
1:B:652:LEU:HD12	1:B:652:LEU:HA	1.79	0.46
1:B:258:GLU:OE1	1:B:258:GLU:N	2.49	0.45
1:B:638:ARG:HD2	1:B:638:ARG:HA	1.77	0.45
1:B:398:MET:HE2	1:B:582:VAL:HG21	1.99	0.45
1:A:190:VAL:HG22	1:A:215:MET:HE1	1.98	0.45
1:A:593:GLU:HG3	1:A:640:VAL:HG22	2.00	0.44
1:A:289:VAL:HA	1:A:293:TYR:CD1	2.53	0.44
1:A:431:LYS:O	1:A:438:ARG:HD3	2.18	0.43
1:A:154:TRP:HH2	1:B:561:THR:HG22	1.83	0.43
1:B:320:VAL:HG13	1:B:386:LEU:HB2	1.99	0.43
1:B:301:VAL:HA	1:B:306:PRO:HA	2.01	0.43
1:A:652:LEU:HD12	1:A:652:LEU:HA	1.83	0.43
1:A:274:ALA:O	1:A:278:ASN:ND2	2.49	0.42
1:B:242:LEU:HD22	1:B:266:TRP:CZ3	2.53	0.42
1:A:321:ASN:OD1	1:A:382:ILE:HB	2.20	0.42
1:B:297:LEU:HD21	1:B:308:CYS:SG	2.59	0.42
1:B:187:ARG:HD2	1:B:226:MET:HE3	2.02	0.42
1:A:177:ILE:HD13	1:A:177:ILE:HA	1.90	0.41
1:A:425:LYS:HG2	1:A:428:ARG:NH2	2.24	0.41
1:B:293:TYR:HA	1:B:313:PHE:O	2.20	0.41
1:B:321:ASN:OD1	1:B:382:ILE:HB	2.20	0.41
1:A:408:LYS:HD3	1:A:573:MET:HG3	2.03	0.41
1:A:231:LYS:HZ3	1:B:569:THR:HG21	1.86	0.41
1:B:637:LEU:HB2	1:B:640:VAL:HG13	2.03	0.41
1:A:593:GLU:HA	1:A:639:ASN:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	435/494 (88%)	420 (97%)	15 (3%)	0	100	100
1	B	410/494 (83%)	393 (96%)	16 (4%)	1 (0%)	43	64
All	All	845/988 (86%)	813 (96%)	31 (4%)	1 (0%)	48	71

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	303	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	393/449 (88%)	383 (98%)	10 (2%)	42	64
1	B	365/449 (81%)	357 (98%)	8 (2%)	45	67
All	All	758/898 (84%)	740 (98%)	18 (2%)	43	65

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	198	ASN
1	A	199	VAL
1	A	263	LEU
1	A	301	VAL
1	A	321	ASN
1	A	336	GLU
1	A	533	VAL
1	A	536	GLU
1	A	629	VAL
1	A	699	GLN
1	B	153	ASP
1	B	157	VAL
1	B	263	LEU
1	B	297	LEU
1	B	321	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	403	GLU
1	B	629	VAL
1	B	640	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	198	ASN
1	A	384	ASN
1	A	424	GLN
1	A	617	ASN
1	A	679	HIS
1	B	539	ASN
1	B	592	HIS
1	B	617	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	WF0	B	801	-	26,26,26	0.15	0	36,36,36	0.46	1 (2%)
3	DMS	A	802	-	3,3,3	0.68	0	3,3,3	0.60	0
4	WF0	A	803	-	26,26,26	0.12	0	36,36,36	0.45	1 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	WF0	B	801	-	-	4/16/16/16	0/2/2/2
4	WF0	A	803	-	-	5/16/16/16	0/2/2/2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	801	WF0	C16-C14-C15	2.20	121.93	116.60
4	A	803	WF0	C16-C14-C15	2.12	121.75	116.60

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	803	WF0	O-C-C1-N
4	A	803	WF0	C13-C14-C15-F2
4	A	803	WF0	C13-C14-C15-F3
4	B	801	WF0	C13-C14-C15-F1
4	B	801	WF0	C13-C14-C15-F2
4	A	803	WF0	C13-C14-C15-F1
4	B	801	WF0	C13-C14-C15-F3
4	B	801	WF0	O-C-C1-N
4	A	803	WF0	C3-C2-N-C1

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	444/494 (89%)	0.45	39 (8%) 15 15	23, 66, 142, 182	3 (0%)
1	B	425/494 (86%)	0.76	51 (12%) 9 7	26, 78, 152, 210	1 (0%)
All	All	869/988 (87%)	0.60	90 (10%) 11 11	23, 71, 145, 210	4 (0%)

All (90) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	266	TRP	6.6
1	B	280	HIS	5.7
1	A	246	PHE	5.6
1	B	243	LYS	5.6
1	B	151	PRO	5.3
1	B	195	LEU	4.6
1	A	255	ASP	4.5
1	B	531	ARG	4.3
1	B	254	GLN	4.3
1	B	372	PHE	4.3
1	B	303	GLU	4.1
1	B	196	PRO	4.0
1	A	256	VAL	4.0
1	B	695	TRP	3.9
1	B	654	TYR	3.8
1	A	380	GLU	3.7
1	B	206[A]	HIS	3.7
1	A	257	SER	3.6
1	B	157	VAL	3.6
1	B	455	LYS	3.5
1	A	169	ASN	3.5
1	B	348	VAL	3.5
1	A	349	LYS	3.4
1	B	159	GLY	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	665	ASP	3.4
1	B	169	ASN	3.4
1	B	351	GLY	3.2
1	A	302	ARG	3.2
1	B	673	LEU	3.2
1	B	277	VAL	3.2
1	A	667	MET	3.1
1	A	530	PRO	3.1
1	B	683	GLU	3.1
1	A	436	PRO	3.1
1	B	653	PRO	3.1
1	B	278	ASN	3.0
1	A	371	GLU	3.0
1	A	657	ALA	3.0
1	B	194	SER	3.0
1	B	154	TRP	3.0
1	B	668	SER	3.0
1	A	214	PHE	3.0
1	A	595	GLN	2.9
1	A	173	PHE	2.9
1	A	320	VAL	2.9
1	A	655	PHE	2.9
1	A	284	SER	2.8
1	A	200	LEU	2.8
1	A	303	GLU	2.8
1	A	532	THR	2.8
1	B	639	ASN	2.7
1	B	437	ALA	2.7
1	A	277	VAL	2.7
1	B	641	SER	2.6
1	A	278	ASN	2.5
1	B	258	GLU	2.5
1	B	370	PHE	2.5
1	B	350	TYR	2.5
1	B	173	PHE	2.4
1	A	318	LEU	2.4
1	A	382	ILE	2.4
1	A	281	ARG	2.4
1	A	437	ALA	2.4
1	A	350	TYR	2.3
1	B	257	SER	2.3
1	B	696	GLU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	435	GLY	2.3
1	B	540	PHE	2.3
1	B	686	TRP	2.3
1	A	434	SER	2.2
1	B	200	LEU	2.2
1	A	206	HIS	2.2
1	B	211	ASN	2.2
1	A	690	GLN	2.2
1	B	216	GLN	2.2
1	A	365	PHE	2.2
1	B	204	ARG	2.1
1	B	275	VAL	2.1
1	A	428	ARG	2.1
1	B	436	PRO	2.1
1	B	608	GLN	2.1
1	B	218	LEU	2.1
1	A	188[A]	ARG	2.1
1	A	263	LEU	2.1
1	B	607	ASN	2.1
1	B	224	LEU	2.1
1	B	532	THR	2.0
1	B	267	LEU	2.0
1	A	244	GLY	2.0
1	B	371	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CL	A	801	1/1	0.82	0.13	62,62,62,62	0
4	WF0	B	801	25/25	0.94	0.14	41,56,81,99	0
4	WF0	A	803	25/25	0.95	0.12	26,56,95,120	0
3	DMS	A	802	4/4	0.97	0.11	27,42,45,62	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around WF0 A 803:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.