



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 11:54 PM UTC

PDB ID : 6P24 / pdb_00006p24
Title : Escherichia coli tRNA synthetase
Authors : Kahne, D.; Baidin, V.; Owens, T.W.
Deposited on : 2019-05-20
Resolution : 2.12 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	8290 (2.14-2.10)
Clashscore	190562	8817 (2.14-2.10)
Ramachandran outliers	187476	8738 (2.14-2.10)
Sidechain outliers	187428	8739 (2.14-2.10)
RSRZ outliers	180081	8294 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	332	12% 63% 9% 27%
1	C	332	31% 80% 17% .
2	B	795	16% 90% 10%
2	D	795	9% 90% 9%

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 17272 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phenylalanine–tRNA ligase alpha subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	241	Total	C	N	O	S	0	1	0
			1925	1227	337	352	9			
1	C	323	Total	C	N	O	S	0	0	0
			2447	1542	441	455	9			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	expression tag	UNP P08312
A	-3	SER	-	expression tag	UNP P08312
A	-2	HIS	-	expression tag	UNP P08312
A	-1	MET	-	expression tag	UNP P08312
A	0	ALA	-	expression tag	UNP P08312
A	1	SER	-	expression tag	UNP P08312
C	-4	GLY	-	expression tag	UNP P08312
C	-3	SER	-	expression tag	UNP P08312
C	-2	HIS	-	expression tag	UNP P08312
C	-1	MET	-	expression tag	UNP P08312
C	0	ALA	-	expression tag	UNP P08312
C	1	SER	-	expression tag	UNP P08312

- Molecule 2 is a protein called Phenylalanine–tRNA ligase beta subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	793	Total	C	N	O	S	0	0	0
			6041	3797	1061	1156	27			
2	D	794	Total	C	N	O	S	0	0	0
			6088	3832	1076	1153	27			


- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	5	Total	Mg	0	0
			5	5		
3	B	4	Total	Mg	0	0
			4	4		
3	C	2	Total	Mg	0	0
			2	2		
3	D	2	Total	Mg	0	0
			2	2		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		

- Molecule 5 is HEXANE-1,6-DIOL (CCD ID: HEZ) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			8	6	2		
5	D	1	Total	C	O	0	0
			8	6	2		
5	D	1	Total	C	O	0	0
			8	6	2		
5	D	1	Total	C	O	0	0
			8	6	2		
5	D	1	Total	C	O	0	0
			8	6	2		

- Molecule 6 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			8	4	1	3		
6	A	1	Total	C	N	O	0	0
			8	4	1	3		
6	C	1	Total	C	N	O	0	0
			8	4	1	3		
6	D	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 7 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			7	4	3		
7	B	1	Total	C	O	0	0
			7	4	3		
7	B	1	Total	C	O	0	0
			7	4	3		
7	B	1	Total	C	O	0	0
			7	4	3		
7	B	1	Total	C	O	0	0
			7	4	3		
7	C	1	Total	C	O	0	0
			7	4	3		
7	C	1	Total	C	O	0	0
			7	4	3		
7	D	1	Total	C	O	0	0
			7	4	3		
7	D	1	Total	C	O	0	0
			7	4	3		

- Molecule 8 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			4	2	2		
8	B	1	Total	C	O	0	0
			4	2	2		
8	B	1	Total	C	O	0	0
			4	2	2		
8	B	1	Total	C	O	0	0
			4	2	2		
8	B	1	Total	C	O	0	0
			4	2	2		
8	C	1	Total	C	O	0	0
			4	2	2		
8	C	1	Total	C	O	0	0
			4	2	2		
8	C	1	Total	C	O	0	0
			4	2	2		
8	D	1	Total	C	O	0	0
			4	2	2		
8	D	1	Total	C	O	0	0
			4	2	2		
8	D	1	Total	C	O	0	0
			4	2	2		
8	D	1	Total	C	O	0	0
			4	2	2		
8	D	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	D	1	Total	C	O	0	0
			4	2	2		
8	D	1	Total	C	O	0	0
			4	2	2		
8	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 9 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: $C_6H_{14}O_4$).






Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	B	1	Total	C	O	0	0
			10	6	4		
9	B	1	Total	C	O	0	0
			10	6	4		

- Molecule 10 is water.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	56	Total	O	0	0
			56	56		
10	B	169	Total	O	0	0
			169	169		
10	C	46	Total	O	0	0
			46	46		
10	D	256	Total	O	0	0
			256	256		

- Molecule 1: Phenylalanine-tRNA ligase alpha subunit





• Molecule 2: Phenylalanine-tRNA ligase beta subunit



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	61.27Å 173.58Å 251.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	82.04 – 2.12 82.04 – 2.12	Depositor EDS
% Data completeness (in resolution range)	99.1 (82.04-2.12) 99.5 (82.04-2.12)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.09 (at 2.12Å)	Xtriage
Refinement program	PHENIX 1.17.1 _3660	Depositor
R, R_{free}	0.229 , 0.265 0.235 , 0.270	Depositor DCC
R_{free} test set	2000 reflections (1.30%)	wwPDB-VP
Wilson B-factor (Å ²)	50.9	Xtriage
Anisotropy	0.148	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 36.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	17272	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, EDO, PEG, HEZ, CL, PGE, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.11	0/1979	0.30	0/2684
1	C	0.10	0/2503	0.29	0/3402
2	B	0.09	0/6142	0.27	0/8350
2	D	0.10	0/6190	0.29	0/8408
All	All	0.10	0/16814	0.28	0/22844

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1925	0	1829	23	0
1	C	2447	0	2281	36	0
2	B	6041	0	6019	52	0
2	D	6088	0	6122	48	0
3	A	5	0	0	0	0
3	B	4	0	0	0	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
4	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	8	0	14	0	0
5	D	32	0	56	2	0
6	A	16	0	24	3	0
6	C	8	0	12	1	0
6	D	8	0	12	0	0
7	A	7	0	10	0	0
7	B	35	0	50	3	0
7	C	14	0	20	2	0
7	D	14	0	20	1	0
8	A	4	0	6	0	0
8	B	16	0	24	4	0
8	C	12	0	18	0	0
8	D	36	0	54	1	0
9	B	20	0	28	1	0
10	A	56	0	0	1	0
10	B	169	0	0	3	0
10	C	46	0	0	1	0
10	D	256	0	0	4	0
All	All	17272	0	16599	143	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 143 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:28:ALA:HB2	2:D:182:LEU:HD11	1.63	0.81
1:C:116:ILE:HD12	1:C:299:MET:HE1	1.64	0.77
2:B:40:GLY:O	2:B:148:ARG:NH2	2.21	0.72
2:B:604:HIS:HA	8:B:814:EDO:H21	1.70	0.71
2:B:497:ARG:HH21	7:C:404:PEG:H32	1.58	0.68

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	240/332 (72%)	230 (96%)	10 (4%)	0	100	100
1	C	321/332 (97%)	305 (95%)	15 (5%)	1 (0%)	36	36
2	B	791/795 (100%)	771 (98%)	19 (2%)	1 (0%)	48	49
2	D	792/795 (100%)	773 (98%)	19 (2%)	0	100	100
All	All	2144/2254 (95%)	2079 (97%)	63 (3%)	2 (0%)	48	49

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	732	VAL
1	C	199	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	202/279 (72%)	199 (98%)	3 (2%)	57	65
1	C	238/279 (85%)	237 (100%)	1 (0%)	84	89
2	B	642/663 (97%)	636 (99%)	6 (1%)	70	78
2	D	651/663 (98%)	645 (99%)	6 (1%)	70	78
All	All	1733/1884 (92%)	1717 (99%)	16 (1%)	70	78

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	732	VAL
2	D	713	VAL
2	B	773	GLU
2	D	568	ARG
2	B	751	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	420	HIS
2	D	488	HIS
2	D	503	ASN
2	B	565	ASN
1	C	61	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 52 ligands modelled in this entry, 14 are monoatomic - leaving 38 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	HEZ	D	805	-	7,7,7	0.31	0	6,6,6	0.93	0
8	EDO	C	408	-	3,3,3	0.45	0	2,2,2	0.38	0
9	PGE	B	813	-	9,9,9	0.32	0	8,8,8	0.28	0
5	HEZ	A	407	-	7,7,7	0.32	0	6,6,6	0.91	0
5	HEZ	D	804	-	7,7,7	0.34	0	6,6,6	0.92	0
7	PEG	B	810	-	6,6,6	0.71	0	5,5,5	0.59	0
8	EDO	C	403	-	3,3,3	0.42	0	2,2,2	0.34	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	TRS	A	411	-	7,7,7	0.31	0	9,9,9	0.26	0
8	EDO	C	406	3	3,3,3	0.44	0	2,2,2	0.37	0
7	PEG	B	805	-	6,6,6	0.70	0	5,5,5	0.55	0
8	EDO	D	811	-	3,3,3	0.44	0	2,2,2	0.32	0
6	TRS	D	807	-	7,7,7	0.32	0	9,9,9	0.28	0
7	PEG	C	404	-	6,6,6	0.72	0	5,5,5	0.55	0
8	EDO	D	808	-	3,3,3	0.41	0	2,2,2	0.46	0
8	EDO	D	817	-	3,3,3	0.43	0	2,2,2	0.39	0
6	TRS	C	407	-	7,7,7	0.31	0	9,9,9	0.31	0
5	HEZ	D	806	-	7,7,7	0.35	0	6,6,6	0.88	0
7	PEG	B	812	-	6,6,6	0.72	0	5,5,5	0.55	0
7	PEG	C	405	-	6,6,6	0.71	0	5,5,5	0.55	0
5	HEZ	D	803	-	7,7,7	0.31	0	6,6,6	0.90	0
6	TRS	A	408	-	7,7,7	0.29	0	9,9,9	0.24	0
8	EDO	B	815	-	3,3,3	0.42	0	2,2,2	0.40	0
8	EDO	D	809	-	3,3,3	0.42	0	2,2,2	0.42	0
7	PEG	D	815	-	6,6,6	0.72	0	5,5,5	0.56	0
8	EDO	D	812	-	3,3,3	0.43	0	2,2,2	0.36	0
8	EDO	B	808	-	3,3,3	0.42	0	2,2,2	0.40	0
7	PEG	D	813	-	6,6,6	0.72	0	5,5,5	0.54	0
7	PEG	A	409	-	6,6,6	0.71	0	5,5,5	0.58	0
8	EDO	D	810	-	3,3,3	0.43	0	2,2,2	0.36	0
8	EDO	B	807	-	3,3,3	0.41	0	2,2,2	0.41	0
7	PEG	B	811	-	6,6,6	0.70	0	5,5,5	0.59	0
8	EDO	D	816	-	3,3,3	0.43	0	2,2,2	0.33	0
8	EDO	D	818	-	3,3,3	0.42	0	2,2,2	0.41	0
8	EDO	D	814	-	3,3,3	0.43	0	2,2,2	0.38	0
8	EDO	A	410	-	3,3,3	0.42	0	2,2,2	0.41	0
8	EDO	B	814	-	3,3,3	0.43	0	2,2,2	0.28	0
9	PGE	B	806	-	9,9,9	0.33	0	8,8,8	0.25	0
7	PEG	B	809	-	6,6,6	0.72	0	5,5,5	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	HEZ	D	805	-	-	1/5/5/5	-
8	EDO	C	408	-	-	0/1/1/1	-
9	PGE	B	813	-	-	4/7/7/7	-
5	HEZ	A	407	-	-	4/5/5/5	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	HEZ	D	804	-	-	4/5/5/5	-
7	PEG	B	810	-	-	0/4/4/4	-
8	EDO	C	403	-	-	0/1/1/1	-
6	TRS	A	411	-	-	2/9/9/9	-
8	EDO	C	406	3	-	1/1/1/1	-
7	PEG	B	805	-	-	3/4/4/4	-
8	EDO	D	811	-	-	0/1/1/1	-
6	TRS	D	807	-	-	0/9/9/9	-
7	PEG	C	404	-	-	1/4/4/4	-
8	EDO	D	808	-	-	0/1/1/1	-
8	EDO	D	817	-	-	0/1/1/1	-
6	TRS	C	407	-	-	0/9/9/9	-
5	HEZ	D	806	-	-	3/5/5/5	-
7	PEG	B	812	-	-	1/4/4/4	-
7	PEG	C	405	-	-	2/4/4/4	-
5	HEZ	D	803	-	-	0/5/5/5	-
6	TRS	A	408	-	-	2/9/9/9	-
8	EDO	B	815	-	-	0/1/1/1	-
8	EDO	D	809	-	-	0/1/1/1	-
7	PEG	D	815	-	-	0/4/4/4	-
8	EDO	D	812	-	-	0/1/1/1	-
8	EDO	B	808	-	-	0/1/1/1	-
7	PEG	D	813	-	-	1/4/4/4	-
7	PEG	A	409	-	-	1/4/4/4	-
8	EDO	D	810	-	-	0/1/1/1	-
8	EDO	B	807	-	-	0/1/1/1	-
7	PEG	B	811	-	-	1/4/4/4	-
8	EDO	D	816	-	-	0/1/1/1	-
8	EDO	D	818	-	-	0/1/1/1	-
8	EDO	D	814	-	-	0/1/1/1	-
8	EDO	A	410	-	-	0/1/1/1	-
8	EDO	B	814	-	-	0/1/1/1	-
9	PGE	B	806	-	-	1/7/7/7	-
7	PEG	B	809	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 34 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	408	TRS	N-C-C2-O2
7	C	405	PEG	O2-C3-C4-O4
5	A	407	HEZ	C1-C2-C3-C4
5	D	804	HEZ	C2-C3-C4-C5
5	D	804	HEZ	C1-C2-C3-C4

There are no ring outliers.

14 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	B	810	PEG	2	0
6	A	411	TRS	1	0
7	C	404	PEG	2	0
6	C	407	TRS	1	0
5	D	806	HEZ	1	0
7	B	812	PEG	1	0
5	D	803	HEZ	1	0
6	A	408	TRS	2	0
8	B	815	EDO	1	0
7	D	815	PEG	1	0
8	D	812	EDO	1	0
8	B	807	EDO	2	0
8	B	814	EDO	1	0
9	B	806	PGE	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	241/332 (72%)	1.05	39 (16%) 4 5	37, 53, 91, 109	1 (0%)
1	C	323/332 (97%)	1.69	104 (32%) 1 1	39, 63, 146, 157	0
2	B	793/795 (99%)	1.10	124 (15%) 5 5	40, 60, 97, 118	0
2	D	794/795 (99%)	0.83	68 (8%) 16 17	33, 53, 75, 91	0
All	All	2151/2254 (95%)	1.08	335 (15%) 5 5	33, 57, 102, 157	1 (0%)

The worst 5 of 335 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	45	LEU	7.8
1	A	201	THR	6.7
2	B	791	ALA	6.7
1	C	27	VAL	6.5
1	C	60	ILE	6.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
8	EDO	D	816	4/4	0.36	0.26	66,68,70,87	0
6	TRS	D	807	8/8	0.41	0.23	64,79,83,84	0
8	EDO	D	818	4/4	0.64	0.20	71,71,71,83	0
6	TRS	C	407	8/8	0.65	0.19	64,74,86,87	0
6	TRS	A	411	8/8	0.67	0.19	75,78,83,83	0
8	EDO	B	815	4/4	0.67	0.26	65,70,73,79	0
7	PEG	D	815	7/7	0.69	0.20	66,76,80,81	0
8	EDO	C	406	4/4	0.70	0.21	67,68,76,77	0
7	PEG	C	404	7/7	0.71	0.23	51,64,78,81	0
7	PEG	B	811	7/7	0.71	0.20	75,79,81,89	0
9	PGE	B	813	10/10	0.72	0.22	62,72,83,92	0
8	EDO	B	807	4/4	0.73	0.18	61,66,71,75	0
3	MG	B	804	1/1	0.74	0.23	78,78,78,78	0
3	MG	D	802	1/1	0.74	0.36	91,91,91,91	0
7	PEG	B	809	7/7	0.75	0.19	54,63,76,78	0
8	EDO	D	814	4/4	0.76	0.21	66,69,71,77	0
7	PEG	B	812	7/7	0.76	0.20	66,68,77,86	0
6	TRS	A	408	8/8	0.77	0.17	50,62,67,68	0
7	PEG	C	405	7/7	0.77	0.20	64,71,81,85	0
5	HEZ	D	804	8/8	0.78	0.20	52,67,76,82	0
8	EDO	B	814	4/4	0.78	0.22	53,63,67,67	0
5	HEZ	D	805	8/8	0.78	0.22	50,61,72,73	0
7	PEG	A	409	7/7	0.78	0.17	67,76,80,88	0
8	EDO	C	408	4/4	0.79	0.19	60,61,64,69	0
8	EDO	D	812	4/4	0.79	0.15	72,73,75,77	0
9	PGE	B	806	10/10	0.80	0.19	57,65,73,84	0
5	HEZ	A	407	8/8	0.82	0.18	46,60,64,66	0
8	EDO	C	403	4/4	0.82	0.17	50,51,69,84	0
7	PEG	B	810	7/7	0.82	0.19	60,68,75,75	0
8	EDO	B	808	4/4	0.82	0.22	57,66,71,73	0
3	MG	A	404	1/1	0.82	0.21	68,68,68,68	0
3	MG	C	401	1/1	0.83	0.14	73,73,73,73	0
5	HEZ	D	803	8/8	0.84	0.21	57,62,68,76	0
8	EDO	D	817	4/4	0.85	0.14	60,65,65,75	0
8	EDO	D	808	4/4	0.85	0.20	54,54,62,71	0
4	CL	A	406	1/1	0.86	0.14	104,104,104,104	0
3	MG	D	801	1/1	0.86	0.15	74,74,74,74	0
3	MG	B	801	1/1	0.88	0.21	69,69,69,69	0
3	MG	C	402	1/1	0.88	0.23	47,47,47,47	0
3	MG	B	803	1/1	0.88	0.12	62,62,62,62	0
3	MG	A	401	1/1	0.88	0.20	48,48,48,48	0
5	HEZ	D	806	8/8	0.89	0.14	45,50,55,59	0
7	PEG	B	805	7/7	0.89	0.13	60,64,71,73	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
8	EDO	D	810	4/4	0.90	0.13	61,61,64,71	0
8	EDO	A	410	4/4	0.90	0.11	54,61,63,64	0
8	EDO	D	811	4/4	0.91	0.14	61,63,66,69	0
7	PEG	D	813	7/7	0.91	0.14	40,55,61,65	0
3	MG	A	402	1/1	0.92	0.20	63,63,63,63	0
3	MG	B	802	1/1	0.93	0.18	58,58,58,58	0
3	MG	A	403	1/1	0.93	0.10	56,56,56,56	0
8	EDO	D	809	4/4	0.93	0.13	48,49,49,52	0
3	MG	A	405	1/1	0.94	0.14	73,73,73,73	0

6.5 Other polymers [i](#)

There are no such residues in this entry.