



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:15 PM UTC

PDB ID : 3P37 / pdb_00003p37
Title : Polo-like kinase I Polo-box domain in complex with FDPPLHSpTA phosphopeptide from PBIP1
Authors : Sledz, P.; Hyvonen, M.; Abell, C.
Deposited on : 2010-10-04
Resolution : 2.38 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.38 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7164 (2.40-2.36)
Clashscore	190562	7722 (2.40-2.36)
Ramachandran outliers	187476	7626 (2.40-2.36)
Sidechain outliers	187428	7627 (2.40-2.36)
RSRZ outliers	180081	7170 (2.40-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	232	2% 80% 16% • •
1	B	232	12% 81% 12% • 6%
1	C	232	17% 81% 13% • 5%
2	D	11	73% 27%
2	E	11	73% 18% 9%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	11	 73% 18% 9%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5596 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein kinase PLK1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	224	Total	C	N	O	S	0	1	0
			1759	1121	292	334	12			
1	B	219	Total	C	N	O	S	0	1	0
			1741	1111	302	317	11			
1	C	220	Total	C	N	O	S	0	1	0
			1727	1103	289	326	9			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP P53350
A	2	PRO	-	expression tag	UNP P53350
A	3	LEU	-	expression tag	UNP P53350
A	4	GLY	-	expression tag	UNP P53350
A	5	SER	-	expression tag	UNP P53350
A	6	PRO	-	expression tag	UNP P53350
A	7	GLU	-	expression tag	UNP P53350
A	8	PHE	-	expression tag	UNP P53350
B	1	GLY	-	expression tag	UNP P53350
B	2	PRO	-	expression tag	UNP P53350
B	3	LEU	-	expression tag	UNP P53350
B	4	GLY	-	expression tag	UNP P53350
B	5	SER	-	expression tag	UNP P53350
B	6	PRO	-	expression tag	UNP P53350
B	7	GLU	-	expression tag	UNP P53350
B	8	PHE	-	expression tag	UNP P53350
C	1	GLY	-	expression tag	UNP P53350
C	2	PRO	-	expression tag	UNP P53350
C	3	LEU	-	expression tag	UNP P53350
C	4	GLY	-	expression tag	UNP P53350
C	5	SER	-	expression tag	UNP P53350
C	6	PRO	-	expression tag	UNP P53350
C	7	GLU	-	expression tag	UNP P53350

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	8	PHE	-	expression tag	UNP P53350

- Molecule 2 is a protein called phosphopeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	10	Total 76	C 47	N 11	O 17	P 1	0	0	0
2	D	11	Total 77	C 47	N 12	O 17	P 1	0	0	1
2	F	10	Total 76	C 47	N 11	O 17	P 1	0	0	0

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	48	Total 48	O 48	0	0
4	B	35	Total 35	O 35	0	0
4	C	23	Total 23	O 23	0	0
4	E	3	Total 3	O 3	0	0
4	D	5	Total 5	O 5	0	0
4	F	2	Total 2	O 2	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Serine/threonine-protein kinase PLK1



- Molecule 1: Serine/threonine-protein kinase PLK1



- Molecule 1: Serine/threonine-protein kinase PLK1



- Molecule 2: phosphopeptide





- Molecule 2: phosphopeptide



- Molecule 2: phosphopeptide



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.97Å 88.66Å 67.59Å 90.00° 113.48° 90.00°	Depositor
Resolution (Å)	62.02 – 2.38 61.99 – 2.38	Depositor EDS
% Data completeness (in resolution range)	99.6 (62.02-2.38) 99.6 (61.99-2.38)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.49 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.232 , 0.288 0.241 , 0.296	Depositor DCC
R_{free} test set	1296 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	30.7	Xtriage
Anisotropy	0.391	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 37.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.037 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	5596	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 46.29 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.1687e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, GOL, ACE, NH2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.47	0/1796	0.76	0/2437
1	B	0.43	0/1779	0.79	0/2406
1	C	0.45	0/1766	0.76	0/2397
2	D	0.50	0/65	0.59	0/88
2	E	0.47	0/65	0.72	0/88
2	F	0.56	0/65	0.96	0/88
All	All	0.45	0/5536	0.77	0/7504

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1759	0	1681	19	0
1	B	1741	0	1721	15	0
1	C	1727	0	1647	22	0
2	D	77	0	64	1	0
2	E	76	0	64	1	0
2	F	76	0	64	1	0
3	A	12	0	16	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	6	0	8	0	0
3	C	6	0	8	0	0
4	A	48	0	0	0	0
4	B	35	0	0	0	0
4	C	23	0	0	0	0
4	D	5	0	0	0	0
4	E	3	0	0	0	0
4	F	2	0	0	0	0
All	All	5596	0	5273	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:496:ASN:HD22	1:C:496:ASN:H	1.25	0.85
1:B:408:ILE:HD13	1:B:500:ARG:HB3	1.65	0.78
1:B:377[A]:MET:HE1	1:B:530:VAL:HG21	1.70	0.74
1:A:377[B]:MET:HE1	1:A:530:VAL:HG21	1.72	0.72
1:A:408:ILE:HD13	1:A:500:ARG:HB3	1.75	0.69
1:B:443:ILE:HD11	1:B:510:TYR:HB3	1.78	0.66
1:C:497:ILE:HD13	1:C:559[A]:PHE:CD1	2.31	0.65
1:C:551:THR:HG21	1:C:559[A]:PHE:CE1	2.32	0.65
1:B:377[A]:MET:HE1	1:B:530:VAL:CG2	2.31	0.61
1:A:384:VAL:HA	1:A:568:GLU:HG2	1.84	0.59
1:C:496:ASN:HD22	1:C:496:ASN:N	1.95	0.59
1:B:434:VAL:HG12	1:B:479:LEU:HD13	1.86	0.58
1:B:452:GLN:HE21	1:B:454:ILE:HD11	1.69	0.58
1:C:444:LEU:HB2	1:C:479:LEU:HD21	1.87	0.57
1:A:397:GLN:HB3	1:A:572:CYS:HA	1.88	0.56
1:C:575:GLU:O	1:C:579:ARG:HG2	2.07	0.54
1:A:538:HIS:NE2	1:C:460:GLU:HG2	2.22	0.54
1:C:412:SER:OG	1:C:428:CYS:HA	2.08	0.53
1:B:460:GLU:HG2	1:B:507:ARG:HH21	1.74	0.52
1:A:377[B]:MET:HE1	1:A:530:VAL:CG2	2.40	0.51
1:B:575:GLU:O	1:B:579:ARG:HG2	2.11	0.51
1:C:443:ILE:HD11	1:C:510:TYR:HB3	1.94	0.50
1:A:555:GLU:CD	1:A:555:GLU:H	2.21	0.49
1:C:534:PHE:CE1	1:C:579:ARG:HD2	2.48	0.49
1:C:541:LEU:HD11	1:C:579:ARG:HB3	1.94	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:441:ARG:HD3	1:B:510:TYR:CD1	2.48	0.48
1:A:538:HIS:NE2	1:C:460:GLU:CG	2.77	0.47
1:C:490:LEU:HA	2:F:79:ALA:HA	1.96	0.47
1:A:377[B]:MET:HE3	1:A:587:VAL:CG2	2.45	0.47
1:C:483:ARG:HA	1:C:486:MET:HE3	1.98	0.46
1:A:377[A]:MET:SD	1:A:543:LEU:HB3	2.56	0.45
1:A:453:TYR:O	1:A:460:GLU:HA	2.17	0.45
1:B:445:TYR:HB2	1:B:450:SER:HB2	2.00	0.44
1:C:423:LEU:HD13	1:C:514:TRP:CD2	2.52	0.44
1:A:552:TYR:O	1:A:559:PHE:HA	2.17	0.44
1:B:541:LEU:HD11	1:B:579:ARG:HB3	1.98	0.44
2:E:73:PRO:HA	2:E:74:PRO:HD3	1.86	0.43
1:B:444:LEU:HB2	1:B:479:LEU:HD21	2.00	0.43
1:A:553:ILE:HA	1:A:558:ASP:O	2.19	0.42
1:C:497:ILE:CD1	1:C:559[A]:PHE:CD1	3.01	0.42
1:A:429:ASP:O	1:A:430:ASN:HB2	2.19	0.42
1:B:412:SER:HA	1:B:493:ALA:HB3	2.01	0.41
1:B:497:ILE:HG12	1:B:559:PHE:CD1	2.55	0.41
1:A:583:ALA:O	1:A:587:VAL:HG23	2.20	0.41
1:C:449:ASP:HA	1:C:465:VAL:HG12	2.02	0.41
2:D:73:PRO:HA	2:D:74:PRO:HD3	1.92	0.41
1:A:403:PRO:O	1:A:406:ILE:HG12	2.20	0.41
1:C:522:ILE:N	1:C:522:ILE:HD12	2.36	0.41
1:C:406:ILE:HG12	1:C:559[A]:PHE:CE2	2.55	0.41
1:C:406:ILE:HG12	1:C:559[A]:PHE:HE2	1.86	0.41
1:C:564:LEU:HD23	1:C:567:LEU:HD12	2.02	0.41
1:A:537:ASP:OD2	1:A:579:ARG:NH2	2.54	0.40
1:A:588:ASP:O	1:A:592:SER:HB3	2.21	0.40
1:C:521:ILE:CG2	1:C:533:ASN:HB2	2.51	0.40
1:A:485:TYR:O	1:A:489:HIS:HB2	2.21	0.40
1:B:522:ILE:N	1:B:522:ILE:HD12	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	223/232 (96%)	215 (96%)	8 (4%)	0	100	100
1	B	216/232 (93%)	209 (97%)	7 (3%)	0	100	100
1	C	217/232 (94%)	208 (96%)	9 (4%)	0	100	100
2	D	8/11 (73%)	6 (75%)	2 (25%)	0	100	100
2	E	7/11 (64%)	6 (86%)	1 (14%)	0	100	100
2	F	7/11 (64%)	6 (86%)	1 (14%)	0	100	100
All	All	678/729 (93%)	650 (96%)	28 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	186/209 (89%)	173 (93%)	13 (7%)	14	21
1	B	187/209 (90%)	179 (96%)	8 (4%)	26	41
1	C	181/209 (87%)	174 (96%)	7 (4%)	28	45
2	D	7/7 (100%)	7 (100%)	0	100	100
2	E	7/7 (100%)	7 (100%)	0	100	100
2	F	7/7 (100%)	6 (86%)	1 (14%)	3	4
All	All	575/648 (89%)	546 (95%)	29 (5%)	21	35

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	372	CYS
1	A	378	LEU
1	A	379	GLN
1	A	406	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	408	ILE
1	A	412	SER
1	A	434	VAL
1	A	463	LEU
1	A	473	MET
1	A	508	LEU
1	A	558	ASP
1	A	560	ARG
1	A	575	GLU
1	B	375	SER
1	B	378	LEU
1	B	434	VAL
1	B	460	GLU
1	B	491	LEU
1	B	516	ARG
1	B	560	ARG
1	B	594	ARG
1	C	383	SER
1	C	391	GLU
1	C	434	VAL
1	C	492	LYS
1	C	496	ASN
1	C	497	ILE
1	C	547	MET
2	F	72	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	430	ASN
1	A	524	HIS
1	B	385	ASN
1	B	430	ASN
1	C	496	ASN
1	C	536	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	TPO	F	78	2	8,10,11	0.76	0	10,14,16	1.06	0
2	TPO	D	78	2	8,10,11	0.88	0	10,14,16	1.28	1 (10%)
2	TPO	E	78	2	8,10,11	0.74	0	10,14,16	1.05	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TPO	F	78	2	-	0/9/11/13	-
2	TPO	D	78	2	-	0/9/11/13	-
2	TPO	E	78	2	-	1/9/11/13	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	78	TPO	O3P-P-O2P	2.24	116.21	107.80

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	E	78	TPO	O-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	B	2	-	5,5,5	0.40	0	5,5,5	0.20	0
3	GOL	A	5	-	5,5,5	0.38	0	5,5,5	0.13	0
3	GOL	C	1	-	5,5,5	0.45	0	5,5,5	0.44	0
3	GOL	A	3	-	5,5,5	0.32	0	5,5,5	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	2	-	-	0/4/4/4	-
3	GOL	A	5	-	-	0/4/4/4	-
3	GOL	C	1	-	-	4/4/4/4	-
3	GOL	A	3	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	3	GOL	O1-C1-C2-C3
3	C	1	GOL	O1-C1-C2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	C	1	GOL	C1-C2-C3-O3
3	A	3	GOL	O1-C1-C2-O2
3	C	1	GOL	O1-C1-C2-O2
3	C	1	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	224/232 (96%)	0.12	5 (2%) 62 61	8, 31, 52, 56	1 (0%)
1	B	219/232 (94%)	0.78	29 (13%) 7 6	14, 43, 93, 112	2 (0%)
1	C	220/232 (94%)	0.97	40 (18%) 3 3	11, 46, 101, 123	1 (0%)
2	D	8/11 (72%)	-0.40	0 100 100	6, 12, 21, 22	0
2	E	8/11 (72%)	-0.06	0 100 100	16, 20, 27, 27	0
2	F	8/11 (72%)	0.46	0 100 100	19, 28, 35, 36	0
All	All	687/729 (94%)	0.60	74 (10%) 11 9	6, 38, 86, 123	4 (0%)

All (74) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	394	LEU	5.0
1	C	472	LEU	4.9
1	A	468	HIS	4.5
1	C	395	VAL	4.2
1	C	505	LEU	4.1
1	B	394	LEU	4.0
1	C	459	THR	4.0
1	C	463	LEU	3.8
1	B	470	ASN	3.7
1	B	505	LEU	3.7
1	C	470	ASN	3.4
1	C	393	GLY	3.3
1	C	468	HIS	3.2
1	C	559[A]	PHE	3.2
1	C	462	TYR	3.2
1	C	471	SER	3.1
1	B	465	VAL	3.1
1	C	461	SER	3.0
1	C	469	PRO	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	463	LEU	3.0
1	B	469	PRO	2.9
1	C	478	LEU	2.9
1	C	473	MET	2.9
1	B	393	GLY	2.8
1	C	479	LEU	2.8
1	B	391	GLU	2.8
1	C	466	SER	2.8
1	C	396	ARG	2.8
1	A	505	LEU	2.8
1	C	501	GLU	2.7
1	B	449	ASP	2.6
1	C	458	GLY	2.6
1	B	384	VAL	2.6
1	B	570	TYR	2.6
1	C	477	THR	2.6
1	B	473	MET	2.6
1	B	389	PRO	2.6
1	C	570	TYR	2.6
1	B	569	GLU	2.5
1	C	504	GLU	2.5
1	A	372	CYS	2.5
1	C	386	ALA	2.5
1	B	398	GLU	2.5
1	C	464	THR	2.5
1	C	467	SER	2.5
1	C	389	PRO	2.5
1	C	391	GLU	2.4
1	B	395	VAL	2.4
1	C	455	GLU	2.4
1	B	593	SER	2.4
1	C	453	TYR	2.3
1	B	400	ALA	2.3
1	C	465	VAL	2.3
1	C	460	GLU	2.3
1	C	390	SER	2.3
1	B	386	ALA	2.3
1	B	467	SER	2.2
1	B	594	ARG	2.2
1	B	464	THR	2.2
1	C	387	SER	2.2
1	A	371	ASP	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	462	TYR	2.1
1	B	457	ASP	2.1
1	B	458	GLY	2.1
1	B	567	LEU	2.1
1	B	396	ARG	2.1
1	B	399	GLU	2.1
1	B	472	LEU	2.1
1	C	398	GLU	2.1
1	C	476	ILE	2.1
1	C	567	LEU	2.1
1	C	421	TYR	2.1
1	C	384	VAL	2.1
1	A	470	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	TPO	F	78	11/12	0.95	0.07	16,18,18,19	0
2	TPO	D	78	11/12	0.97	0.06	2,4,6,6	0
2	TPO	E	78	11/12	0.97	0.06	15,16,17,18	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	A	3	6/6	0.70	0.13	36,37,38,38	0
3	GOL	C	1	6/6	0.74	0.16	29,32,33,34	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	B	2	6/6	0.80	0.11	28,30,30,31	0
3	GOL	A	5	6/6	0.87	0.10	40,40,40,41	0

6.5 Other polymers [i](#)

There are no such residues in this entry.