



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 12:11 AM UTC

PDB ID : 6P3D / pdb_00006p3d
Title : The co-crystal structure of BRAF(V600E) with ponatinib
Authors : Agianian, B.; Gavathiotis, E.
Deposited on : 2019-05-23
Resolution : 2.11 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.11 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	8290 (2.14-2.10)
Clashscore	190562	8817 (2.14-2.10)
Ramachandran outliers	187476	8738 (2.14-2.10)
Sidechain outliers	187428	8739 (2.14-2.10)
RSRZ outliers	180081	8294 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	297	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NH4	A	1010	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 2293 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein kinase B-raf.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	264	Total	C	N	O	S	0	0	0
			2108	1333	374	387	14			

There are 38 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	427	MET	-	initiating methionine	UNP P15056
A	428	GLY	-	expression tag	UNP P15056
A	429	SER	-	expression tag	UNP P15056
A	430	SER	-	expression tag	UNP P15056
A	431	HIS	-	expression tag	UNP P15056
A	432	HIS	-	expression tag	UNP P15056
A	433	HIS	-	expression tag	UNP P15056
A	434	HIS	-	expression tag	UNP P15056
A	435	HIS	-	expression tag	UNP P15056
A	436	HIS	-	expression tag	UNP P15056
A	437	SER	-	expression tag	UNP P15056
A	438	SER	-	expression tag	UNP P15056
A	439	GLY	-	expression tag	UNP P15056
A	440	LEU	-	expression tag	UNP P15056
A	441	VAL	-	expression tag	UNP P15056
A	442	PRO	-	expression tag	UNP P15056
A	443	ARG	-	expression tag	UNP P15056
A	444	GLY	-	expression tag	UNP P15056
A	445	SER	-	expression tag	UNP P15056
A	446	LYS	-	expression tag	UNP P15056
A	447	MET	-	expression tag	UNP P15056
A	543	ALA	ILE	conflict	UNP P15056
A	544	SER	ILE	conflict	UNP P15056
A	551	LYS	ILE	conflict	UNP P15056
A	562	ARG	GLN	conflict	UNP P15056
A	588	ASN	LEU	conflict	UNP P15056
A	600	GLU	VAL	conflict	UNP P15056

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	630	SER	LYS	conflict	UNP P15056
A	667	GLU	PHE	conflict	UNP P15056
A	673	SER	TYR	conflict	UNP P15056
A	688	ARG	ALA	conflict	UNP P15056
A	706	SER	LEU	conflict	UNP P15056
A	709	ARG	GLN	conflict	UNP P15056
A	713	GLU	SER	conflict	UNP P15056
A	716	GLU	LEU	conflict	UNP P15056
A	720	GLU	SER	conflict	UNP P15056
A	722	SER	-	expression tag	UNP P15056
A	723	GLY	-	expression tag	UNP P15056

- Molecule 2 is 3-(imidazo[1,2-b]pyridazin-3-ylethynyl)-4-methyl-N-{4-[(4-methylpiperazin-1-yl)methyl]-3-(trifluoromethyl)phenyl}benzamide (CCD ID: 0LI) (formula: C₂₉H₂₇F₃N₆O) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	0	0
			39	29	3	6	1		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is AMMONIUM ION (CCD ID: NH4) (formula: H₄N).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	N		0	0
			1	1			

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	97	Total	O	0	0
			97	97		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Serine/threonine-protein kinase B-raf

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 2 2	Depositor
Cell constants a, b, c, α , β , γ	118.84Å 118.84Å 51.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	84.03 – 2.11 84.03 – 2.11	Depositor EDS
% Data completeness (in resolution range)	99.4 (84.03-2.11) 99.4 (84.03-2.11)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.42 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.185 , 0.245 0.195 , 0.256	Depositor DCC
R_{free} test set	1117 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	27.0	Xtriage
Anisotropy	0.024	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 35.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2293	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NH4, OLI, SO4, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.20	7/2150 (0.3%)	1.14	6/2895 (0.2%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	617	ILE	CA-C	8.65	1.65	1.52
1	A	499	LYS	N-CA	-5.24	1.40	1.46
1	A	639	VAL	C-O	-5.20	1.18	1.24
1	A	448	ASP	N-CA	5.19	1.52	1.46
1	A	569	ALA	C-O	-5.17	1.17	1.24
1	A	552	LYS	N-CA	-5.14	1.40	1.46
1	A	515	LEU	C-O	-5.13	1.17	1.24

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	707	PHE	CA-C-N	-6.45	111.89	119.05
1	A	707	PHE	C-N-CA	-6.45	111.89	119.05
1	A	625	ILE	N-CA-C	5.75	115.94	110.42
1	A	626	ARG	N-CA-C	5.21	117.04	111.36
1	A	544	SER	CA-C-N	5.09	127.61	120.28
1	A	544	SER	C-N-CA	5.09	127.61	120.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2108	0	2123	24	0
2	A	39	0	27	0	0
3	A	40	0	0	2	0
4	A	1	0	0	2	0
5	A	8	0	12	1	0
6	A	97	0	0	4	0
All	All	2293	0	2162	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:515:LEU:H	1:A:530:GLN:HE21	1.35	0.73
1:A:495:LEU:HD23	1:A:525:LEU:HD21	1.77	0.67
1:A:512:ASN:HD21	1:A:559:GLN:HE21	1.45	0.64
3:A:1008:SO4:O1	6:A:1101:HOH:O	2.14	0.62
1:A:544:SER:HB3	1:A:546:THR:HG22	1.85	0.58
1:A:614:SER:HB3	1:A:617:ILE:HG23	1.87	0.57
1:A:559:GLN:HE22	1:A:589:THR:HA	1.68	0.56
1:A:522:LYS:HA	1:A:523:PRO:C	2.34	0.53
1:A:706:SER:HB3	1:A:708:PRO:HD2	1.92	0.52
1:A:540:HIS:HD1	1:A:546:THR:HG21	1.73	0.52
1:A:512:ASN:ND2	1:A:559:GLN:HE21	2.08	0.51
1:A:539:HIS:ND1	1:A:600:GLU:OE1	2.27	0.50
1:A:653:GLN:HA	5:A:1011:EDO:H11	1.96	0.48
1:A:614:SER:C	1:A:616:SER:N	2.72	0.48
1:A:587:ASP:OD1	4:A:1010:NH4:N	2.46	0.47
1:A:614:SER:O	1:A:616:SER:N	2.43	0.47
1:A:636:GLN:HG2	6:A:1162:HOH:O	2.14	0.47
1:A:584:LEU:O	4:A:1010:NH4:N	2.47	0.47
1:A:684:ASN:ND2	3:A:1007:SO4:O2	2.47	0.46
1:A:486:ASN:OD1	1:A:486:ASN:N	2.43	0.46
1:A:695:GLU:OE1	1:A:709:ARG:NH2	2.50	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:559:GLN:NE2	1:A:590:VAL:H	2.16	0.44
1:A:700:LYS:HG2	6:A:1103:HOH:O	2.19	0.42
1:A:614:SER:C	1:A:616:SER:H	2.28	0.40
1:A:688:ARG:NH1	6:A:1102:HOH:O	2.23	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	260/297 (88%)	252 (97%)	7 (3%)	1 (0%)	30	28

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	523	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	232/260 (89%)	223 (96%)	9 (4%)	28	29

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	446	LYS
1	A	448	ASP
1	A	486	ASN
1	A	525	LEU
1	A	584	LEU
1	A	600	GLU
1	A	671	ARG
1	A	691	ARG
1	A	719	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	530	GLN
1	A	559	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 1 is modelled with single atom - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	1002	-	4,4,4	0.29	0	6,6,6	0.62	0
3	SO4	A	1006	-	4,4,4	0.46	0	6,6,6	0.39	0
2	OLI	A	1001	-	43,43,43	2.28	10 (23%)	59,62,62	2.76	16 (27%)
3	SO4	A	1004	-	4,4,4	0.66	0	6,6,6	0.38	0
3	SO4	A	1008	-	4,4,4	0.70	0	6,6,6	0.61	0
5	EDO	A	1011	-	3,3,3	0.67	0	2,2,2	0.27	0
5	EDO	A	1012	-	3,3,3	0.38	0	2,2,2	0.59	0
3	SO4	A	1005	-	4,4,4	0.50	0	6,6,6	0.52	0
3	SO4	A	1009	-	4,4,4	0.37	0	6,6,6	0.37	0
3	SO4	A	1007	-	4,4,4	0.55	0	6,6,6	0.41	0
3	SO4	A	1003	-	4,4,4	0.57	0	6,6,6	0.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	A	1012	-	-	1/1/1/1	-
5	EDO	A	1011	-	-	0/1/1/1	-
2	OLI	A	1001	-	-	2/21/33/33	0/5/5/5

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1001	OLI	C5-C4	-7.27	1.30	1.43
2	A	1001	OLI	C2-N82	-6.31	1.30	1.37
2	A	1001	OLI	C16-C17	5.40	1.47	1.40
2	A	1001	OLI	C5-C6	4.19	1.47	1.40
2	A	1001	OLI	C84-N82	-3.88	1.33	1.39
2	A	1001	OLI	C13-N2	-3.43	1.34	1.41
2	A	1001	OLI	C3-C2	-3.32	1.37	1.42
2	A	1001	OLI	C83-N81	3.05	1.35	1.31
2	A	1001	OLI	C84-N1	2.93	1.37	1.33
2	A	1001	OLI	N82-N81	2.27	1.40	1.37

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1001	OLI	C1-C2-N82	14.14	108.61	103.61
2	A	1001	OLI	C1-C2-C3	-7.35	118.73	130.61
2	A	1001	OLI	C3-C2-N82	5.95	132.16	122.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1001	OLI	C9-C10-C5	4.59	124.68	120.60
2	A	1001	OLI	C2-C1-N1	-3.84	108.23	112.35
2	A	1001	OLI	C22-C21-N3	-3.52	103.56	110.65
2	A	1001	OLI	O1-C12-C9	-3.43	114.11	120.90
2	A	1001	OLI	C9-C12-N2	3.23	123.75	115.90
2	A	1001	OLI	C24-C23-N4	-3.07	105.92	110.86
2	A	1001	OLI	C10-C5-C6	-3.05	115.10	121.44
2	A	1001	OLI	C14-C13-C18	-3.04	115.98	119.66
2	A	1001	OLI	C13-C18-C17	2.63	124.09	120.82
2	A	1001	OLI	F2-C19-C17	-2.58	108.07	112.65
2	A	1001	OLI	C84-N82-C2	2.12	108.20	106.06
2	A	1001	OLI	C7-C6-C5	2.06	120.09	117.80
2	A	1001	OLI	C18-C17-C19	2.00	121.46	116.60

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1001	OLI	C3-C4-C5-C6
5	A	1012	EDO	O1-C1-C2-O2
2	A	1001	OLI	C3-C4-C5-C10

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1008	SO4	1	0
5	A	1011	EDO	1	0
3	A	1007	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	264/297 (88%)	0.17	20 (7%)	20 21	17, 29, 68, 83	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	487	VAL	5.6
1	A	488	THR	4.9
1	A	616	SER	4.3
1	A	546	THR	4.2
1	A	486	ASN	3.9
1	A	630	SER	3.8
1	A	489	ALA	3.8
1	A	523	PRO	3.6
1	A	662	ARG	3.4
1	A	721	LEU	3.2
1	A	615	GLY	2.9
1	A	446	LYS	2.8
1	A	628	GLN	2.8
1	A	544	SER	2.5
1	A	627	MET	2.5
1	A	716	GLU	2.4
1	A	614	SER	2.4
1	A	629	ASP	2.1
1	A	682	ARG	2.1
1	A	722	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates


There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	A	1007	5/5	0.75	0.12	75,82,83,89	0
3	SO4	A	1004	5/5	0.81	0.14	70,71,83,90	0
3	SO4	A	1006	5/5	0.85	0.10	70,72,84,85	0
3	SO4	A	1005	5/5	0.85	0.09	74,75,78,78	0
5	EDO	A	1011	4/4	0.90	0.13	42,44,45,51	0
4	NH4	A	1010	1/1	0.93	0.10	24,24,24,24	0
3	SO4	A	1008	5/5	0.93	0.11	28,31,33,33	5
5	EDO	A	1012	4/4	0.94	0.14	32,34,38,39	0
3	SO4	A	1003	5/5	0.96	0.11	47,50,53,56	0
3	SO4	A	1002	5/5	0.96	0.14	39,39,49,54	0
3	SO4	A	1009	5/5	0.96	0.08	36,38,42,44	5
2	OLI	A	1001	39/39	0.97	0.05	17,20,23,24	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.