



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 20, 2026 – 01:56 PM UTC

PDB ID : 8P3J / pdb_00008p3j
Title : Fusion hSlp2-a_Rab27A non-covalent complex with hit fragment IMP-1704
Authors : De Vita, E.; Brustur, D.; Tersa, M.; Petracca, R.; Morgan, R.M.L.; Lanyon-Hogg, T.; Norman, J.C.; Cota, E.; Tate, E.W.
Deposited on : 2023-05-17
Resolution : 2.16 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.16 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	230	53% 75% 17% 8%
1	B	230	60% 69% 24% 7%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	WT6	A	201	-	X	-	X
2	WT6	A	202	-	-	-	X
2	WT6	B	201	-	X	-	-
2	WT6	B	203	-	X	-	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3686 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Synaptotagmin-like protein 2,Ras-related protein Rab-27A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	212	Total	C	N	O	S	0	2	0
			1710	1083	295	324	8			
1	B	214	Total	C	N	O	S	0	2	0
			1719	1085	298	328	8			

There are 26 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-38	GLY	-	expression tag	UNP Q9HCH5
A	-37	HIS	-	expression tag	UNP Q9HCH5
A	-36	MET	-	expression tag	UNP Q9HCH5
A	-6	GLY	-	linker	UNP Q9HCH5
A	-5	SER	-	linker	UNP Q9HCH5
A	-4	GLY	-	linker	UNP Q9HCH5
A	-3	SER	-	linker	UNP Q9HCH5
A	-2	GLY	-	linker	UNP Q9HCH5
A	-1	SER	-	linker	UNP Q9HCH5
A	0	GLY	-	linker	UNP Q9HCH5
A	78	LEU	GLN	engineered mutation	UNP P51159
A	123	ALA	CYS	engineered mutation	UNP P51159
A	188	SER	CYS	engineered mutation	UNP P51159
B	-38	GLY	-	expression tag	UNP Q9HCH5
B	-37	HIS	-	expression tag	UNP Q9HCH5
B	-36	MET	-	expression tag	UNP Q9HCH5
B	-6	GLY	-	linker	UNP Q9HCH5
B	-5	SER	-	linker	UNP Q9HCH5
B	-4	GLY	-	linker	UNP Q9HCH5
B	-3	SER	-	linker	UNP Q9HCH5
B	-2	GLY	-	linker	UNP Q9HCH5
B	-1	SER	-	linker	UNP Q9HCH5
B	0	GLY	-	linker	UNP Q9HCH5
B	78	LEU	GLN	engineered mutation	UNP P51159
B	123	ALA	CYS	engineered mutation	UNP P51159

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	188	SER	CYS	engineered mutation	UNP P51159

- Molecule 2 is {N}-(4-pyrrolidin-1-ylsulfonylphenyl)prop-2-enamide (CCD ID: WT6) (formula: C₁₃H₁₆N₂O₃S) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			19	13	2	3	1		
2	A	1	Total	C	N	O	S	0	0
			19	13	2	3	1		
2	B	1	Total	C	N	O	S	0	0
			19	13	2	3	1		
2	B	1	Total	C	N	O	S	0	0
			19	13	2	3	1		

- Molecule 3 is PHOSPHOAMINOPHOSPHONIC ACID-GUANYLATE ESTER (CCD ID: GNP) (formula: C₁₀H₁₇N₆O₁₃P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			32	10	6	13	3		
3	B	1	Total	C	N	O	P	0	0
			32	10	6	13	3		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mg	0	0
			1	1		
5	B	1	Total	Mg	0	0
			1	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	68	Total	O	0	0
			68	68		
6	B	29	Total	O	0	0
			29	29		

- Molecule 1: Synaptotagmin-like protein 2, Ras-related protein Rab-27A



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	62.44Å 75.83Å 118.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	63.88 – 2.16 63.88 – 2.16	Depositor EDS
% Data completeness (in resolution range)	96.4 (63.88-2.16) 96.5 (63.88-2.16)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.96 (at 2.16Å)	Xtriage
Refinement program	PHENIX (1.17_3644: ???)	Depositor
R, R_{free}	0.284 , 0.351 0.285 , 0.349	Depositor DCC
R_{free} test set	1579 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	4.5	Xtriage
Anisotropy	0.244	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 33.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.74	EDS
Total number of atoms	3686	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, GNP, WT6, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	0/1741	0.55	0/2340
1	B	0.36	0/1745	0.58	0/2345
All	All	0.36	0/3486	0.57	0/4685

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	7	ASP	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1710	0	1688	41	3
1	B	1719	0	1695	53	3
2	A	38	0	0	2	0
2	B	38	0	0	6	0
3	A	32	0	13	4	0
3	B	32	0	13	1	0
4	A	6	0	8	0	0
4	B	12	0	15	3	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	68	0	0	3	0
6	B	29	0	0	1	0
All	All	3686	0	3432	78	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 78 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:125:ASN:ND2	2:B:203:WT6:C04	1.68	1.55
1:A:124:GLU:CG	1:B:-13:ARG:HD2	1.61	1.28
1:B:125:ASN:ND2	2:B:203:WT6:C05	1.98	1.27
1:A:124:GLU:HG3	1:B:-13:ARG:CD	1.73	1.17
1:A:124:GLU:HG3	1:B:-13:ARG:HD2	1.14	1.11

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:138:GLU:OE2	1:B:138:GLU:OE1[2_554]	1.46	0.74
1:A:138:GLU:CD	1:B:138:GLU:OE1[2_554]	1.87	0.33
1:A:67:ARG:NH2	1:B:115:SER:OG[4_455]	2.15	0.05

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	210/230 (91%)	205 (98%)	4 (2%)	1 (0%)	24	20
1	B	212/230 (92%)	198 (93%)	11 (5%)	3 (1%)	9	4
All	All	422/460 (92%)	403 (96%)	15 (4%)	4 (1%)	12	9

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	-9	GLU
1	B	60	GLY
1	A	42	VAL
1	B	62	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	180/192 (94%)	180 (100%)	0	100	100
1	B	180/192 (94%)	180 (100%)	0	100	100
All	All	360/384 (94%)	360 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	35	ASN
1	A	133	ASN
1	B	-20	GLN
1	B	105	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 11 ligands modelled in this entry, 2 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	B	204	-	5,5,5	0.94	0	5,5,5	0.81	0
3	GNP	A	203	5	34,34,34	1.31	5 (14%)	47,54,54	0.86	3 (6%)
4	GOL	B	205	-	5,5,5	1.35	1 (20%)	5,5,5	1.39	1 (20%)
2	WT6	B	201	-	20,20,20	4.05	13 (65%)	28,28,28	3.60	12 (42%)
2	WT6	B	203	-	20,20,20	4.12	13 (65%)	28,28,28	6.17	18 (64%)
2	WT6	A	201	-	20,20,20	4.17	14 (70%)	28,28,28	4.78	10 (35%)
3	GNP	B	202	5	34,34,34	1.40	5 (14%)	47,54,54	0.84	1 (2%)
4	GOL	A	204	-	5,5,5	0.65	0	5,5,5	1.62	1 (20%)
2	WT6	A	202	-	20,20,20	3.89	11 (55%)	28,28,28	3.53	9 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	B	204	-	-	1/4/4/4	-
3	GNP	A	203	5	-	8/18/38/38	0/3/3/3
4	GOL	B	205	-	-	3/4/4/4	-
2	WT6	B	201	-	-	5/18/25/25	0/2/2/2
2	WT6	B	203	-	-	13/18/25/25	0/2/2/2
2	WT6	A	201	-	-	8/18/25/25	0/2/2/2
3	GNP	B	202	5	-	4/18/38/38	0/3/3/3
4	GOL	A	204	-	-	2/4/4/4	-
2	WT6	A	202	-	-	5/18/25/25	0/2/2/2

The worst 5 of 62 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	202	WT6	S02-N14	10.34	1.78	1.63
2	B	201	WT6	S02-N14	10.11	1.77	1.63
2	A	201	WT6	S02-N14	10.02	1.77	1.63
2	B	203	WT6	S02-N14	9.71	1.77	1.63
2	B	203	WT6	C08-N07	8.44	1.51	1.35

The worst 5 of 55 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	201	WT6	C09-C08-N07	15.94	124.67	113.84
2	B	203	WT6	C13-C03-S02	-15.31	104.53	119.73
2	B	203	WT6	C03-S02-N14	15.10	125.51	107.28
2	A	201	WT6	O19-S02-O01	-12.33	100.34	119.59
2	B	201	WT6	O19-S02-O01	-12.03	100.81	119.59

There are no chirality outliers.

5 of 49 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	202	WT6	N07-C08-C09-C10
2	A	202	WT6	O11-C08-C09-C10
2	B	201	WT6	N07-C08-C09-C10
2	B	201	WT6	O11-C08-C09-C10
2	B	201	WT6	C15-N14-S02-C03

There are no ring outliers.


7 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	204	GOL	1	0
3	A	203	GNP	4	0
4	B	205	GOL	2	0
2	B	201	WT6	1	0
2	B	203	WT6	5	0
3	B	202	GNP	1	0
2	A	202	WT6	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand WT6 B 201




Bond lengths



Bond angles




Torsions




Rings


Ligand WT6 B 203




Bond lengths





Bond angles




Torsions



Rings





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	212/230 (92%)	2.42	122 (57%) 0 0	13, 23, 44, 91	2 (0%)
1	B	214/230 (93%)	2.84	137 (64%) 0 0	10, 25, 63, 93	2 (0%)
All	All	426/460 (92%)	2.63	259 (60%) 0 0	10, 24, 60, 93	4 (0%)

The worst 5 of 259 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	58	PRO	10.2
1	B	60	GLY	9.5
1	B	61	ALA	9.3
1	B	57	GLY	8.8
1	A	58	PRO	8.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	WT6	A	201	19/19	0.36	0.41	30,41,57,71	19
2	WT6	A	202	19/19	0.39	0.42	19,52,84,96	19
2	WT6	B	203	19/19	0.39	0.31	37,53,71,82	19
2	WT6	B	201	19/19	0.49	0.33	29,35,50,59	9
4	GOL	B	204	6/6	0.52	0.25	20,27,39,48	0
4	GOL	B	205	6/6	0.68	0.20	26,29,36,42	0
4	GOL	A	204	6/6	0.70	0.23	27,28,29,29	0
5	MG	A	205	1/1	0.70	0.09	21,21,21,21	0
3	GNP	B	202	32/32	0.90	0.11	10,17,22,30	0
3	GNP	A	203	32/32	0.92	0.10	8,17,23,28	0
5	MG	B	206	1/1	0.95	0.09	8,8,8,8	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around WT6 A 202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around WT6 B 203:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around WT6 B 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around GNP B 202:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.