



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 10:03 AM UTC

PDB ID : 4P4B / pdb_00004p4b
Title : X-ray structure of human glutamate carboxypeptidase II (GCPII) in complex with a phosphoramidate inhibitor CTT54
Authors : Barinka, C.
Deposited on : 2014-03-12
Resolution : 1.93 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION




The reported resolution of this entry is 1.93 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1452 (1.94-1.94)
Clashscore	190562	1494 (1.94-1.94)
Ramachandran outliers	187476	1479 (1.94-1.94)
Sidechain outliers	187428	1479 (1.94-1.94)
RSRZ outliers	180081	1453 (1.94-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	707	 86% 11% .
2	B	2	 50% 50%
2	C	2	 100%
2	D	2	 100%
3	E	3	 100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	F	4	 25% 75%

2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 6529 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate carboxypeptidase 2.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	694	Total	C	N	O	S	0	50	0
			5782	3719	964	1077	22			

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	D	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	F	4	Total	C	N	O	0	0	0
			50	28	2	20			

- Molecule 5 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Zn	0	0
			2	2		

- Molecule 6 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Ca	0	0
			1	1		

- Molecule 7 is CHLORIDE ION (CCD ID: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Cl	0	0
			1	1		

- Molecule 8 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	A	1	Total	C	N	O	0	0
			14	8	1	5		
8	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 9 is L-gamma-glutamyl-O-[(S)-{[(1S)-1,3-dicarboxypropyl]amino}(hydroxy)phosphoryl]-L-serine (CCD ID: 2G2) (formula: C₁₃H₂₂N₃O₁₂P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
9	A	1	Total	C	N	O	P	0	0
			29	13	3	12	1		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	513	Total	O	0	0
			513	513		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glutamate carboxypeptidase 2




- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  100%

MAG1
MAG2
BMA3

- Molecule 4: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  25% 75%

MAG1
MAG2
BMA3
MAN4

4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	101.63Å 130.31Å 158.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.82 – 1.93 19.82 – 1.93	Depositor EDS
% Data completeness (in resolution range)	99.1 (19.82-1.93) 99.5 (19.82-1.93)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.43 (at 1.92Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.155 , 0.178 0.155 , 0.179	Depositor DCC
R_{free} test set	2380 reflections (2.98%)	wwPDB-VP
Wilson B-factor (Å ²)	23.7	Xtriage
Anisotropy	0.047	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 50.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6529	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, BMA, ZN, CL, 2G2, CA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.05	5/6080 (0.1%)	0.95	7/8235 (0.1%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	705	PHE	N-CA	5.37	1.50	1.46
1	A	554[A]	SER	CA-C	5.02	1.59	1.52
1	A	554[B]	SER	CA-C	5.02	1.59	1.52
1	A	239[A]	VAL	CA-CB	5.00	1.60	1.54
1	A	239[B]	VAL	CA-CB	5.00	1.60	1.54

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	549[A]	TYR	CA-C-N	5.76	125.43	119.56
1	A	549[A]	TYR	C-N-CA	5.76	125.43	119.56
1	A	549[B]	TYR	CA-C-N	5.76	125.43	119.56
1	A	549[B]	TYR	C-N-CA	5.76	125.43	119.56
1	A	146	PRO	CA-C-N	-5.46	115.73	119.66
1	A	146	PRO	C-N-CA	-5.46	115.73	119.66
1	A	652	ASP	N-CA-C	5.18	119.75	113.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5782	0	5644	53	0
2	B	28	0	25	1	0
2	C	28	0	25	0	0
2	D	28	0	25	0	0
3	E	39	0	34	0	0
4	F	50	0	43	0	0
5	A	2	0	0	0	0
6	A	1	0	0	0	0
7	A	1	0	0	0	0
8	A	28	0	26	0	0
9	A	29	0	17	1	0
10	A	513	0	0	12	0
All	All	6529	0	5839	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:660[A]:VAL:O	1:A:664[A]:MET:HG2	1.46	1.16
1:A:412:PRO:HA	1:A:589[B]:ASN:HD21	1.25	0.98
1:A:656[B]:SER:O	10:A:902:HOH:O	1.96	0.84
1:A:90[B]:PHE:CE2	1:A:94:LYS:HE2	2.15	0.82
1:A:617[B]:LYS:NZ	10:A:1313:HOH:O	2.12	0.79
1:A:486[A]:GLU:OE1	10:A:1198:HOH:O	2.01	0.78
1:A:506:PHE:HB2	1:A:509:MET:HG3	1.72	0.72
1:A:400:ARG:O	1:A:404[B]:THR:HG23	1.90	0.72
1:A:656[B]:SER:OG	10:A:902:HOH:O	2.12	0.67
1:A:623:LYS:NZ	10:A:906:HOH:O	2.26	0.66
1:A:688[B]:ARG:NH2	10:A:1050:HOH:O	2.28	0.65
1:A:688[B]:ARG:NH1	10:A:907:HOH:O	2.32	0.63
1:A:612:TYR:CZ	1:A:616:MET:HG3	2.35	0.61
1:A:198:GLY:O	1:A:223:LYS:HE2	2.03	0.58
1:A:308:LYS:HB2	10:A:1255:HOH:O	2.04	0.58
1:A:641:GLU:HG3	10:A:1408:HOH:O	2.04	0.56
1:A:688[B]:ARG:CZ	10:A:907:HOH:O	2.53	0.56
1:A:239[A]:VAL:HG22	1:A:247:ASN:ND2	2.24	0.52
1:A:412:PRO:CA	1:A:589[B]:ASN:HD21	2.10	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:551:LEU:HD22	1:A:556:TYR:HB2	1.93	0.50
1:A:169:MET:HE3	1:A:347:HIS:NE2	2.25	0.50
1:A:201[A]:VAL:HG21	1:A:217:ALA:HB1	1.93	0.50
1:A:258:ILE:HD13	1:A:294:VAL:HB	1.94	0.49
1:A:610:LYS:CE	10:A:1145:HOH:O	2.60	0.49
1:A:174[A]:LEU:HG	1:A:202:ILE:HG22	1.95	0.48
1:A:413:ARG:H	1:A:589[B]:ASN:ND2	2.11	0.48
1:A:90[B]:PHE:CZ	1:A:94:LYS:HE2	2.48	0.48
1:A:169:MET:HA	1:A:344:MET:O	2.14	0.48
1:A:188:LEU:CD2	1:A:329:VAL:HG11	2.47	0.45
1:A:239[A]:VAL:HG22	1:A:247:ASN:CG	2.41	0.45
1:A:205:TYR:CE1	1:A:254:GLN:HB3	2.51	0.45
1:A:188:LEU:HD21	1:A:329:VAL:HG11	1.99	0.44
1:A:133:GLU:HG2	1:A:339:THR:HB	2.00	0.44
1:A:177:VAL:HG12	1:A:188:LEU:HD11	2.00	0.44
1:A:687:TYR:CZ	1:A:694:PRO:HG2	2.53	0.44
1:A:645[B]:LYS:HD3	1:A:645[B]:LYS:HA	1.85	0.43
1:A:309[B]:MET:HE3	1:A:309[B]:MET:HB3	1.92	0.43
1:A:659[B]:ILE:HD13	1:A:659[B]:ILE:HA	1.76	0.43
1:A:552:TYR:CD2	9:A:820:2G2:H2	2.54	0.43
1:A:377:HIS:CE1	1:A:388:PRO:HB3	2.54	0.42
1:A:169:MET:HE2	1:A:345:HIS:HD2	1.83	0.42
1:A:79[B]:GLN:OE1	2:B:1:NAG:H81	2.19	0.42
1:A:96[B]:ILE:HD13	10:A:1064:HOH:O	2.20	0.41
1:A:263:GLY:HA2	1:A:522:GLU:OE1	2.20	0.41
1:A:324:LYS:HA	1:A:324:LYS:HD3	1.89	0.41
1:A:551:LEU:HD23	1:A:551:LEU:HA	1.84	0.41
1:A:469:LEU:O	1:A:595:PHE:HA	2.21	0.41
1:A:295:HIS:HB2	1:A:346:ILE:HD11	2.02	0.41
1:A:58[A]:MET:HE2	1:A:58[A]:MET:HB3	1.77	0.41
1:A:177:VAL:CG1	1:A:201[B]:VAL:HG22	2.51	0.41
1:A:171[A]:GLU:HG2	1:A:343:LYS:HG3	2.03	0.40
1:A:174[B]:LEU:HD11	1:A:342:VAL:HG21	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	738/707 (104%)	720 (98%)	16 (2%)	2 (0%)	36	30

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	335	GLY
1	A	382	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	640/603 (106%)	626 (98%)	14 (2%)	45	35

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	55	LYS
1	A	239[A]	VAL
1	A	239[B]	VAL
1	A	267	PRO
1	A	303	GLN
1	A	336	ASN
1	A	489	GLU
1	A	503	SER
1	A	505	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	537	TYR
1	A	542	GLU
1	A	549[A]	TYR
1	A	549[B]	TYR
1	A	600	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	95	GLN
1	A	97	GLN
1	A	443	GLN
1	A	618	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

13 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
8	NAG	A	811	1	14,14,15	0.54	0	17,19,21	2.59	2 (11%)
4	NAG	F	1	1,4	14,14,15	0.80	0	17,19,21	1.33	3 (17%)
4	NAG	F	2	4	14,14,15	0.72	0	17,19,21	1.68	2 (11%)
8	NAG	A	812	1	14,14,15	0.73	0	17,19,21	1.54	1 (5%)
4	MAN	F	4	4	11,11,12	0.65	0	15,15,17	1.18	1 (6%)
2	NAG	D	1	2,1	14,14,15	0.60	0	17,19,21	1.40	1 (5%)
2	NAG	D	2	2	14,14,15	0.58	0	17,19,21	1.25	2 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	C	1	2,1	14,14,15	0.91	1 (7%)	17,19,21	1.77	6 (35%)
2	NAG	B	1	2,1	14,14,15	0.57	0	17,19,21	1.28	3 (17%)
2	NAG	B	2	2	14,14,15	0.57	0	17,19,21	1.20	1 (5%)
2	NAG	C	2	2	14,14,15	0.65	0	17,19,21	1.41	2 (11%)
3	NAG	E	1	1,3	14,14,15	1.11	1 (7%)	17,19,21	0.97	0
3	NAG	E	2	3	14,14,15	0.61	0	17,19,21	1.15	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	NAG	A	811	1	-	2/6/23/26	0/1/1/1
4	NAG	F	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	F	2	4	-	2/6/23/26	0/1/1/1
8	NAG	A	812	1	-	0/6/23/26	0/1/1/1
4	MAN	F	4	4	-	0/2/19/22	0/1/1/1
2	NAG	D	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	D	2	2	-	0/6/23/26	0/1/1/1
2	NAG	C	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	B	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	1/6/23/26	0/1/1/1
3	NAG	E	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	E	2	3	-	2/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1	NAG	O5-C1	-2.38	1.39	1.43
3	E	1	NAG	O7-C7	2.14	1.28	1.23

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	A	811	NAG	C1-O5-C5	9.60	125.05	112.19
8	A	812	NAG	C1-O5-C5	4.48	118.20	112.19
4	F	2	NAG	C3-C4-C5	-4.05	102.88	110.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	2	NAG	C1-O5-C5	4.02	117.58	112.19
2	D	1	NAG	C1-O5-C5	3.71	117.16	112.19
3	E	2	NAG	C8-C7-N2	3.22	121.46	116.12
2	C	1	NAG	C2-N2-C7	-2.97	118.92	122.90
2	C	1	NAG	C4-C3-C2	2.83	115.16	111.02
2	D	2	NAG	O4-C4-C5	2.82	116.26	109.32
4	F	4	MAN	O5-C5-C6	2.76	113.03	107.66
4	F	2	NAG	C8-C7-N2	2.75	120.68	116.12
4	F	1	NAG	C1-O5-C5	2.74	115.85	112.19
2	C	1	NAG	O5-C5-C6	2.66	112.83	107.66
8	A	811	NAG	C4-C3-C2	-2.52	107.33	111.02
2	C	1	NAG	C1-C2-N2	-2.45	106.57	110.43
2	D	2	NAG	C3-C4-C5	-2.45	105.79	110.23
4	F	1	NAG	C2-N2-C7	-2.43	119.64	122.90
2	B	1	NAG	C1-O5-C5	2.29	115.26	112.19
2	B	1	NAG	O4-C4-C3	-2.21	105.16	110.38
2	B	2	NAG	O4-C4-C5	2.20	114.74	109.32
2	B	1	NAG	O5-C1-C2	-2.11	108.02	111.29
2	C	2	NAG	O4-C4-C5	2.06	114.39	109.32
2	C	1	NAG	O3-C3-C2	-2.05	105.15	109.40
4	F	1	NAG	O4-C4-C5	-2.04	104.31	109.32
2	C	1	NAG	C6-C5-C4	2.02	117.97	113.02

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	A	811	NAG	C4-C5-C6-O6
3	E	1	NAG	C8-C7-N2-C2
3	E	1	NAG	O7-C7-N2-C2
3	E	2	NAG	C8-C7-N2-C2
3	E	2	NAG	O7-C7-N2-C2
4	F	2	NAG	C8-C7-N2-C2
4	F	2	NAG	O7-C7-N2-C2
2	C	2	NAG	O5-C5-C6-O6
8	A	811	NAG	O5-C5-C6-O6
4	F	1	NAG	C4-C5-C6-O6
4	F	1	NAG	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1	NAG	1	0

5.5 Carbohydrates [i](#)

13 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	B	1	2,1	14,14,15	0.57	0	17,19,21	1.28	3 (17%)
2	NAG	B	2	2	14,14,15	0.57	0	17,19,21	1.20	1 (5%)
2	NAG	C	1	2,1	14,14,15	0.91	1 (7%)	17,19,21	1.77	6 (35%)
2	NAG	C	2	2	14,14,15	0.65	0	17,19,21	1.41	2 (11%)
2	NAG	D	1	2,1	14,14,15	0.60	0	17,19,21	1.40	1 (5%)
2	NAG	D	2	2	14,14,15	0.58	0	17,19,21	1.25	2 (11%)
3	NAG	E	1	1,3	14,14,15	1.11	1 (7%)	17,19,21	0.97	0
3	NAG	E	2	3	14,14,15	0.61	0	17,19,21	1.15	1 (5%)
3	BMA	E	3	3	11,11,12	0.80	0	15,15,17	1.45	2 (13%)
4	NAG	F	1	1,4	14,14,15	0.80	0	17,19,21	1.33	3 (17%)
4	NAG	F	2	4	14,14,15	0.72	0	17,19,21	1.68	2 (11%)
4	BMA	F	3	4	11,11,12	0.56	0	15,15,17	0.98	0
4	MAN	F	4	4	11,11,12	0.65	0	15,15,17	1.18	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
2	NAG	C	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	1/6/23/26	0/1/1/1
2	NAG	D	1	2,1	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	D	2	2	-	0/6/23/26	0/1/1/1
3	NAG	E	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	E	2	3	-	2/6/23/26	0/1/1/1
3	BMA	E	3	3	-	0/2/19/22	0/1/1/1
4	NAG	F	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	F	2	4	-	2/6/23/26	0/1/1/1
4	BMA	F	3	4	-	0/2/19/22	0/1/1/1
4	MAN	F	4	4	-	0/2/19/22	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1	NAG	O5-C1	-2.38	1.39	1.43
3	E	1	NAG	O7-C7	2.14	1.28	1.23

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	F	2	NAG	C3-C4-C5	-4.05	102.88	110.23
2	C	2	NAG	C1-O5-C5	4.02	117.58	112.19
2	D	1	NAG	C1-O5-C5	3.71	117.16	112.19
3	E	2	NAG	C8-C7-N2	3.22	121.46	116.12
3	E	3	BMA	C3-C4-C5	3.12	115.89	110.23
2	C	1	NAG	C2-N2-C7	-2.97	118.92	122.90
2	C	1	NAG	C4-C3-C2	2.83	115.16	111.02
2	D	2	NAG	O4-C4-C5	2.82	116.26	109.32
3	E	3	BMA	C1-O5-C5	2.79	115.93	112.19
4	F	4	MAN	O5-C5-C6	2.76	113.03	107.66
4	F	2	NAG	C8-C7-N2	2.75	120.68	116.12
4	F	1	NAG	C1-O5-C5	2.74	115.85	112.19
2	C	1	NAG	O5-C5-C6	2.66	112.83	107.66
2	C	1	NAG	C1-C2-N2	-2.45	106.57	110.43
2	D	2	NAG	C3-C4-C5	-2.45	105.79	110.23
4	F	1	NAG	C2-N2-C7	-2.43	119.64	122.90
2	B	1	NAG	C1-O5-C5	2.29	115.26	112.19
2	B	1	NAG	O4-C4-C3	-2.21	105.16	110.38
2	B	2	NAG	O4-C4-C5	2.20	114.74	109.32
2	B	1	NAG	O5-C1-C2	-2.11	108.02	111.29
2	C	2	NAG	O4-C4-C5	2.06	114.39	109.32
2	C	1	NAG	O3-C3-C2	-2.05	105.15	109.40
4	F	1	NAG	O4-C4-C5	-2.04	104.31	109.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	NAG	C6-C5-C4	2.02	117.97	113.02

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	E	1	NAG	C8-C7-N2-C2
3	E	1	NAG	O7-C7-N2-C2
3	E	2	NAG	C8-C7-N2-C2
3	E	2	NAG	O7-C7-N2-C2
4	F	2	NAG	C8-C7-N2-C2
4	F	2	NAG	O7-C7-N2-C2
2	C	2	NAG	O5-C5-C6-O6
4	F	1	NAG	C4-C5-C6-O6
4	F	1	NAG	O5-C5-C6-O6


There are no ring outliers.


1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 4 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
8	NAG	A	811	1	14,14,15	0.54	0	17,19,21	2.59	2 (11%)
9	2G2	A	820	5	27,28,28	1.73	7 (25%)	33,38,38	1.37	4 (12%)
8	NAG	A	812	1	14,14,15	0.73	0	17,19,21	1.54	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	NAG	A	811	1	-	2/6/23/26	0/1/1/1
9	2G2	A	820	5	-	1/36/37/37	-
8	NAG	A	812	1	-	0/6/23/26	0/1/1/1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	A	820	2G2	CAP-CBB	3.40	1.61	1.53
9	A	820	2G2	CAN-CAV	3.26	1.57	1.51
9	A	820	2G2	CBA-NAR	3.16	1.52	1.45
9	A	820	2G2	PBC-OAG	2.84	1.50	1.46
9	A	820	2G2	CB-CA	2.38	1.58	1.53
9	A	820	2G2	CAM-CAU	2.19	1.55	1.50
9	A	820	2G2	CBB-NAS	2.14	1.49	1.47

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	A	811	NAG	C1-O5-C5	9.60	125.05	112.19
8	A	812	NAG	C1-O5-C5	4.48	118.20	112.19
9	A	820	2G2	OAG-PBC-NAS	-3.22	107.66	112.75
9	A	820	2G2	PBC-NAS-CBB	-3.08	117.16	124.03
8	A	811	NAG	C4-C3-C2	-2.52	107.33	111.02

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	A	820	2G2	OAL-PBC-OAG	2.43	115.08	109.87
9	A	820	2G2	CB-CAN-CAV	-2.43	107.65	113.06

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
8	A	811	NAG	C4-C5-C6-O6
8	A	811	NAG	O5-C5-C6-O6
9	A	820	2G2	OAK-CAY-CBB-NAS

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	A	820	2G2	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	694/707 (98%)	-0.60	6 (0%) 81 85	8, 24, 46, 67	49 (7%)

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	546	PHE	4.4
1	A	656[A]	SER	4.1
1	A	506	PHE	2.4
1	A	545	LYS	2.4
1	A	653	PHE	2.2
1	A	549[A]	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	C	2	14/15	0.63	0.15	73,78,79,80	0
8	NAG	A	811	14/15	0.72	0.14	65,72,74,75	0
2	NAG	C	1	14/15	0.78	0.14	49,53,65,68	0
2	NAG	B	2	14/15	0.81	0.11	43,50,56,56	0
2	NAG	D	2	14/15	0.83	0.10	52,62,68,68	0
3	NAG	E	2	14/15	0.83	0.12	44,51,57,62	0
4	MAN	F	4	11/12	0.86	0.11	49,53,55,56	0
4	NAG	F	2	14/15	0.87	0.12	36,43,50,51	0
8	NAG	A	812	14/15	0.89	0.09	32,48,51,53	0
2	NAG	D	1	14/15	0.92	0.08	41,46,49,54	0
3	NAG	E	1	14/15	0.94	0.07	29,33,38,39	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NAG	B	1	14/15	0.94	0.07	35,42,45,48	0
4	NAG	F	1	14/15	0.95	0.06	20,29,42,45	0

6.3 Carbohydrates




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NAG	C	2	14/15	0.63	0.15	73,78,79,80	0
3	BMA	E	3	11/12	0.70	0.15	66,72,78,80	0
2	NAG	C	1	14/15	0.78	0.14	49,53,65,68	0
2	NAG	B	2	14/15	0.81	0.11	43,50,56,56	0
3	NAG	E	2	14/15	0.83	0.12	44,51,57,62	0
2	NAG	D	2	14/15	0.83	0.10	52,62,68,68	0
4	MAN	F	4	11/12	0.86	0.11	49,53,55,56	0
4	NAG	F	2	14/15	0.87	0.12	36,43,50,51	0
4	BMA	F	3	11/12	0.90	0.09	41,44,47,48	0
2	NAG	D	1	14/15	0.92	0.08	41,46,49,54	0
3	NAG	E	1	14/15	0.94	0.07	29,33,38,39	0
2	NAG	B	1	14/15	0.94	0.07	35,42,45,48	0
4	NAG	F	1	14/15	0.95	0.06	20,29,42,45	0




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain B:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain C:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain D:


$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)



In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
8	NAG	A	811	14/15	0.72	0.14	65,72,74,75	0
8	NAG	A	812	14/15	0.89	0.09	32,48,51,53	0
9	2G2	A	820	29/29	0.98	0.06	14,19,39,41	0
7	CL	A	804	1/1	1.00	0.01	22,22,22,22	0
5	ZN	A	801	1/1	1.00	0.01	19,19,19,19	0
5	ZN	A	802	1/1	1.00	0.01	20,20,20,20	0
6	CA	A	803	1/1	1.00	0.01	17,17,17,17	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 2G2 A 820:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.