



wwPDB EM Validation Summary Report ⓘ

Mar 5, 2026 – 06:36 PM UTC

PDB ID : 6P4K / pdb_00006p4k
EMDB ID : EMD-20251
Title : mouse norovirus complexed with TCA
Authors : Smith, T.J.
Deposited on : 2019-05-27
Resolution : 3.10 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY


The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	229148	23984
Ramachandran outliers	224038	23583
Sidechain outliers	223484	23102

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	516	 81% 73% 26%
1	B	516	 80% 75% 24% .
1	C	516	 78% 79% 18% .

2 Entry composition [i](#)


There are 2 unique types of molecules in this entry. The entry contains 11963 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Capsid protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	514	Total	C	N	O	S	0	0
			3948	2544	650	738	16		
1	B	513	Total	C	N	O	S	0	0
			3941	2539	649	737	16		
1	C	502	Total	C	N	O	S	0	0
			3864	2493	636	719	16		

- Molecule 2 is TAUROCHOLIC ACID (CCD ID: TCH) (formula: C₂₆H₄₅NO₇S).




Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf
2	C	1	Total	C	N	O	S	0
			35	26	1	7	1	
2	C	1	Total	C	N	O	S	0
			35	26	1	7	1	



3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Capsid protein



• Molecule 1: Capsid protein



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	25000	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	28	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	OTHER	Depositor
Maximum map value	19.155	Depositor
Minimum map value	-13.334	Depositor
Average map value	0.005	Depositor
Map value standard deviation	0.647	Depositor
Recommended contour level	1	Depositor
Map size ($\text{\AA}$)	624.456, 624.456, 624.456	wwPDB
Map dimensions	540, 540, 540	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	1.1564, 1.1564, 1.1564	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TCH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.49	0/4056	0.52	0/5561
1	B	0.50	0/4050	0.54	4/5554 (0.1%)
1	C	0.49	0/3972	0.51	0/5447
All	All	0.49	0/12078	0.52	4/16562 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5
1	C	0	2
All	All	0	7

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	469	GLN	CA-C-N	-6.79	113.35	121.90
1	B	469	GLN	C-N-CA	-6.79	113.35	121.90
1	B	215	PHE	CA-C-N	-5.40	117.68	122.97
1	B	215	PHE	C-N-CA	-5.40	117.68	122.97

There are no chirality outliers.

5 of 7 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	173	GLN	Peptide

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	A	221	PRO	Peptide
1	A	224	ARG	Peptide
1	A	225	THR	Peptide
1	A	458	VAL	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3948	0	3893	90	0
1	B	3941	0	3883	82	0
1	C	3864	0	3812	63	0
2	A	70	0	76	21	0
2	B	70	0	76	24	0
2	C	70	0	78	20	0
All	All	11963	0	11818	287	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 287 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:602:TCH:C17	2:A:602:TCH:C16	1.76	1.63
2:B:601:TCH:C7	2:B:601:TCH:C6	1.75	1.63
2:A:602:TCH:C7	2:A:602:TCH:C6	1.76	1.62
2:B:602:TCH:C16	2:B:602:TCH:C17	1.76	1.61
2:B:601:TCH:C17	2:B:601:TCH:C16	1.76	1.60

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	510/516 (99%)	453 (89%)	55 (11%)	2 (0%)	30	61
1	B	511/516 (99%)	445 (87%)	65 (13%)	1 (0%)	43	73
1	C	500/516 (97%)	440 (88%)	57 (11%)	3 (1%)	21	52
All	All	1521/1548 (98%)	1338 (88%)	177 (12%)	6 (0%)	31	61

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	225	THR
1	A	224	ARG
1	C	194	PRO
1	B	196	ASP
1	C	132	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	426/427 (100%)	425 (100%)	1 (0%)	87	89
1	B	425/427 (100%)	422 (99%)	3 (1%)	76	82
1	C	417/427 (98%)	416 (100%)	1 (0%)	87	89
All	All	1268/1281 (99%)	1263 (100%)	5 (0%)	81	86

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	138	THR
1	B	79	LEU
1	B	100	THR
1	B	112	LEU
1	C	414	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 26 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	236	GLN
1	B	340	GLN
1	C	409	ASN
1	B	270	HIS
1	B	455	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	TCH	C	602	-	38,38,38	8.78	30 (78%)	59,60,60	2.31	22 (37%)
2	TCH	A	601	-	38,38,38	8.75	30 (78%)	59,60,60	2.65	29 (49%)
2	TCH	B	602	-	38,38,38	8.78	30 (78%)	59,60,60	2.36	24 (40%)
2	TCH	B	601	-	38,38,38	8.76	30 (78%)	59,60,60	2.58	21 (35%)
2	TCH	A	602	-	38,38,38	8.78	30 (78%)	59,60,60	2.47	21 (35%)
2	TCH	C	601	-	38,38,38	8.79	30 (78%)	59,60,60	2.45	22 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TCH	C	602	-	-	11/16/81/81	0/4/4/4
2	TCH	A	601	-	-	10/16/81/81	0/4/4/4
2	TCH	B	602	-	-	9/16/81/81	0/4/4/4
2	TCH	B	601	-	-	9/16/81/81	0/4/4/4
2	TCH	A	602	-	-	6/16/81/81	0/4/4/4
2	TCH	C	601	-	-	11/16/81/81	0/4/4/4

The worst 5 of 180 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	TCH	C8-C14	-25.75	1.04	1.53
2	A	602	TCH	C8-C14	-25.69	1.04	1.53
2	C	601	TCH	C8-C14	-25.66	1.04	1.53
2	B	602	TCH	C8-C14	-25.57	1.04	1.53
2	C	602	TCH	C8-C14	-25.40	1.05	1.53

The worst 5 of 139 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	TCH	C14-C13-C12	8.32	115.02	107.42
2	A	602	TCH	C14-C13-C12	7.26	114.05	107.42
2	A	601	TCH	C13-C17-C20	-6.93	111.09	119.48
2	C	601	TCH	C14-C13-C12	6.93	113.75	107.42
2	B	602	TCH	C13-C17-C20	-6.89	111.13	119.48

There are no chirality outliers.

5 of 56 torsion outliers are listed below:




Mol	Chain	Res	Type	Atoms
2	A	601	TCH	C13-C17-C20-C21
2	A	601	TCH	N24-C25-C26-S26
2	A	602	TCH	N24-C25-C26-S26
2	B	601	TCH	N24-C25-C26-S26
2	B	602	TCH	C13-C17-C20-C21


There are no ring outliers.

6 monomers are involved in 65 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	602	TCH	9	0
2	A	601	TCH	12	0
2	B	602	TCH	12	0
2	B	601	TCH	12	0
2	A	602	TCH	9	0
2	C	601	TCH	11	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	296:GLU	C	297:PHE	N	5.74


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-20251. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 270



Y Index: 270




Z Index: 270


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 345



Y Index: 345




Z Index: 195


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z


The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)


6.5.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 1.0. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 4629 nm^3 ; this corresponds to an approximate mass of 4181 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.323 Å⁻¹

8 Fourier-Shell correlation ⓘ

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-20251 and PDB model 6P4K. Per-residue inclusion information can be found in section 3 on page 5.

9.1 Map-model overlay [i](#)


This section was not generated.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (1).

9.4 Atom inclusion [i](#)



At the recommended contour level, 18% of all backbone atoms, 18% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (1) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.1830	-0.0050
A	0.1790	-0.0030
B	0.1880	-0.0110
C	0.1810	-0.0000

