



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:40 PM UTC

PDB ID : 7P4W / pdb_00007p4w
Title : Crystal structure of alpha-amylase from *Aspergillus oryzae* in space group I222
Authors : Bellini, D.; Gorrec, F.
Deposited on : 2021-07-13
Resolution : 2.28 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.28 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	9078 (2.30-2.26)
Clashscore	190562	9802 (2.30-2.26)
Ramachandran outliers	187476	9690 (2.30-2.26)
Sidechain outliers	187428	9691 (2.30-2.26)
RSRZ outliers	180081	9085 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	476	8% 84% 14%
2	E	2	50% 50%

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 3785 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-amylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	476	Total	C	N	O	S	0	0	0
			3686	2336	595	737	18			

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	E	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca	0	0
			1	1		

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	58	Total	O	0	0
			58	58		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Alpha-amylase



• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	65.72Å 140.40Å 154.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	51.91 – 2.28 51.91 – 2.28	Depositor EDS
% Data completeness (in resolution range)	99.1 (51.91-2.28) 99.1 (51.91-2.28)	Depositor EDS
R_{merge}	9.90	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.96 (at 2.27Å)	Xtriage
Refinement program	REFMAC 1.17.1_3660, PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.228 , 0.285 0.238 , 0.289	Depositor DCC
R_{free} test set	1606 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å ²)	49.2	Xtriage
Anisotropy	0.301	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 24.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3785	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, CA, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/3781	0.66	2/5164 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	445	VAL	CA-C-N	-5.13	114.62	122.06
1	A	445	VAL	C-N-CA	-5.13	114.62	122.06

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	446	GLY	Peptide
1	A	51	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3686	0	3478	53	0
2	E	28	0	25	1	0
3	A	1	0	0	0	0
4	A	12	0	12	0	0
5	A	58	0	0	2	0
All	All	3785	0	3515	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:85:GLN:HA	1:A:123:MET:HE3	1.58	0.85
1:A:157:ASP:OD1	1:A:159:THR:HG22	1.90	0.72
1:A:216:TRP:HB2	1:A:245:VAL:HG12	1.80	0.63
1:A:49:LEU:HB2	1:A:110:ARG:NH1	2.14	0.62
1:A:310:LEU:O	1:A:314:VAL:HG23	2.05	0.57
1:A:65:VAL:HG12	1:A:90:LEU:HD22	1.86	0.57
1:A:50:ASP:OD1	1:A:110:ARG:NH2	2.38	0.56
1:A:162:GLU:OE2	1:A:212:GLN:HA	2.05	0.56
1:A:87:ILE:HG12	1:A:139:PRO:HD3	1.86	0.56
1:A:236:PRO:HG3	1:A:253:PRO:HB3	1.88	0.55
1:A:51:TYR:C	1:A:51:TYR:CD1	2.85	0.55
1:A:96:THR:HG22	1:A:98:ASP:H	1.72	0.54
1:A:205:ILE:HD12	1:A:246:MET:HE1	1.91	0.51
1:A:204:ARG:NH2	1:A:230:GLU:HG3	2.26	0.50
1:A:382:TYR:CE2	1:A:398:LYS:HA	2.47	0.50
1:A:129:GLY:HA3	1:A:149:PHE:CE1	2.45	0.50
1:A:14:LEU:HD11	1:A:62:ILE:HG22	1.93	0.50
1:A:334:HIS:HE1	5:A:618:HOH:O	1.95	0.50
1:A:193:SER:OG	2:E:1:NAG:N2	2.44	0.49
1:A:6:TRP:CD1	1:A:226:TYR:HB3	2.48	0.48
1:A:51:TYR:CD2	1:A:352:TYR:HB3	2.48	0.48
1:A:79:TYR:HD2	1:A:79:TYR:H	1.62	0.48
1:A:61:TRP:C	1:A:61:TRP:CD1	2.92	0.47
1:A:140:PHE:HA	1:A:145:TYR:CD2	2.48	0.47
1:A:18:ARG:NH2	1:A:79:TYR:CD1	2.83	0.46
1:A:1:ALA:HA	1:A:201:ASP:O	2.16	0.46
1:A:115:MET:HA	1:A:202:GLY:O	2.15	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:147:HIS:NE2	1:A:177:ASP:HA	2.30	0.45
1:A:371:TYR:CE1	1:A:472:SER:HB3	2.51	0.44
1:A:322:ASP:CG	1:A:384:ASN:H	2.26	0.44
1:A:79:TYR:CD2	1:A:79:TYR:N	2.86	0.44
1:A:216:TRP:HB2	1:A:245:VAL:CG1	2.47	0.44
1:A:51:TYR:C	1:A:51:TYR:HD1	2.24	0.44
1:A:176:LEU:HD13	1:A:187:TRP:HE1	1.83	0.43
1:A:134:TYR:CD1	1:A:134:TYR:N	2.86	0.43
1:A:382:TYR:CD2	1:A:398:LYS:HA	2.53	0.43
1:A:129:GLY:O	1:A:132:VAL:HG23	2.19	0.43
1:A:32:THR:HB	1:A:341:PRO:O	2.19	0.43
1:A:134:TYR:CD2	1:A:143:GLN:HA	2.54	0.43
1:A:162:GLU:CD	1:A:212:GLN:HA	2.43	0.43
1:A:272:LEU:O	1:A:276:ILE:HG13	2.19	0.43
1:A:243:GLN:HB3	1:A:284:PRO:HG2	2.01	0.43
1:A:262:PHE:CB	1:A:314:VAL:HG22	2.49	0.42
1:A:74:ALA:HB3	1:A:170:THR:CG2	2.49	0.42
1:A:55:MET:HG2	1:A:359:TYR:CD1	2.54	0.41
1:A:205:ILE:HD12	1:A:246:MET:CE	2.50	0.41
1:A:147:HIS:CD2	1:A:177:ASP:HA	2.56	0.41
1:A:65:VAL:CG1	1:A:90:LEU:HD22	2.48	0.41
1:A:140:PHE:HA	1:A:145:TYR:HD2	1.84	0.41
1:A:84:GLN:HG2	5:A:649:HOH:O	2.20	0.41
1:A:445:VAL:O	1:A:450:ASN:O	2.39	0.40
1:A:15:LEU:HD22	1:A:63:THR:HG21	2.03	0.40
1:A:134:TYR:N	1:A:134:TYR:HD1	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	474/476 (100%)	456 (96%)	18 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	397/397 (100%)	391 (98%)	6 (2%)	57	72

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	51	TYR
1	A	61	TRP
1	A	79	TYR
1	A	134	TYR
1	A	314	VAL
1	A	359	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	43	GLN
1	A	85	GLN
1	A	93	ASN
1	A	298	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	E	1	2,1	14,14,15	0.69	1 (7%)	17,19,21	0.83	0
2	NAG	E	2	2	14,14,15	0.68	0	17,19,21	1.12	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	E	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	E	2	2	-	2/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	1	NAG	O5-C1	2.29	1.47	1.43

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	2	NAG	C1-O5-C5	3.99	117.53	112.19

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	E	2	NAG	O5-C5-C6-O6
2	E	2	NAG	C4-C5-C6-O6
2	E	1	NAG	O5-C5-C6-O6
2	E	1	NAG	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	MES	A	502	-	12,12,12	2.12	1 (8%)	15,16,16	2.23	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MES	A	502	-	-	5/6/14/14	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	502	MES	C8-S	-6.86	1.68	1.77

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	502	MES	C5-N4-C3	5.54	120.78	108.84
4	A	502	MES	O2S-S-C8	3.96	112.72	106.73
4	A	502	MES	O1S-S-C8	3.35	111.80	106.73

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	502	MES	N4-C7-C8-S
4	A	502	MES	C7-C8-S-O2S
4	A	502	MES	C7-C8-S-O3S
4	A	502	MES	C7-C8-S-O1S

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	A	502	MES	C8-C7-N4-C3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	476/476 (100%)	0.59	39 (8%) 17 18	37, 51, 75, 90	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	155	TYR	4.1
1	A	219	TYR	3.5
1	A	151	PHE	3.3
1	A	182	VAL	3.3
1	A	149	PHE	3.3
1	A	218	GLY	3.2
1	A	193	SER	3.0
1	A	446	GLY	3.0
1	A	139	PRO	3.0
1	A	178	THR	2.9
1	A	138	LYS	2.9
1	A	140	PHE	2.9
1	A	212	GLN	2.8
1	A	215	PHE	2.8
1	A	1	ALA	2.8
1	A	185	ASN	2.7
1	A	179	THR	2.6
1	A	150	CYS	2.6
1	A	163	ASP	2.6
1	A	190	TRP	2.5
1	A	159	THR	2.4
1	A	180	LYS	2.3
1	A	88	TYR	2.3
1	A	145	TYR	2.3
1	A	187	TRP	2.3
1	A	476	SER	2.3
1	A	471	GLY	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	474	ILE	2.2
1	A	213	LYS	2.2
1	A	225	VAL	2.2
1	A	216	TRP	2.2
1	A	152	ILE	2.1
1	A	147	HIS	2.1
1	A	222	ALA	2.1
1	A	238	TYR	2.1
1	A	374	SER	2.1
1	A	244	ASN	2.1
1	A	211	VAL	2.1
1	A	31	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	E	1	14/15	-	-	64,81,90,93	0
2	NAG	E	2	14/15	-	-	85,99,106,106	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MES	A	502	12/12	0.68	0.23	58,62,95,98	0
3	CA	A	501	1/1	0.98	0.04	65,65,65,65	0

6.5 Other polymers [i](#)

There are no such residues in this entry.