



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 11:51 AM UTC

PDB ID : 1P53 / pdb\_00001p53  
Title : The Crystal Structure of ICAM-1 D3-D5 fragment  
Authors : Yang, Y.; Jun, C.D.; Liu, J.H.; Zhang, R.; Jochimiak, A.; Springer, T.A.; Wang, J.H.  
Deposited on : 2003-04-24  
Resolution : 3.06 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	2469 (3.10-3.02)
Clashscore	190562	2569 (3.10-3.02)
Ramachandran outliers	187476	2424 (3.10-3.02)
Sidechain outliers	187428	2423 (3.10-3.02)
RSRZ outliers	180081	2469 (3.10-3.02)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	266	
1	B	266	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	1269	X	-	-	-
2	NAG	B	2240	X	-	-	-

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 3955 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Intercellular adhesion molecule-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	250	Total 1903	C 1185	N 328	O 381	S 9	0	0	0
1	B	250	Total 1903	C 1185	N 328	O 381	S 9	0	0	0

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	Total 14	C 8	N 1	O 5	0	0
2	A	1	Total 14	C 8	N 1	O 5	0	0
2	A	1	Total 14	C 8	N 1	O 5	0	0
2	B	1	Total 14	C 8	N 1	O 5	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	42	Total	O	0	0
			42	42		
3	B	23	Total	O	0	0
			23	23		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Intercellular adhesion molecule-1



- Molecule 1: Intercellular adhesion molecule-1



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	193.70Å 193.70Å 175.30Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 3.06 20.00 – 3.06	Depositor EDS
% Data completeness (in resolution range)	89.9 (20.00-3.06) 94.8 (20.00-3.06)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.32 (at 3.04Å)	Xtrriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.220 , 0.252 0.222 , 0.251	Depositor DCC
$R_{free}$ test set	2263 reflections (9.94%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	44.6	Xtrriage
Anisotropy	0.004	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 29.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	3955	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 34.41 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.8551e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.49	0/1939	0.99	11/2650 (0.4%)
1	B	0.45	0/1939	0.97	6/2650 (0.2%)
All	All	0.47	0/3878	0.98	17/5300 (0.3%)

There are no bond length outliers.

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	376	CYS	CA-C-N	7.66	127.71	119.90
1	A	376	CYS	C-N-CA	7.66	127.71	119.90
1	B	289	ILE	N-CA-C	6.54	117.29	107.80
1	B	358	ASN	N-CA-C	6.48	119.01	108.26
1	A	324	GLY	CA-C-N	6.45	126.36	119.85
1	A	324	GLY	C-N-CA	6.45	126.36	119.85
1	B	236	VAL	N-CA-C	6.22	117.11	108.89
1	A	396	GLY	N-CA-C	6.07	119.98	111.00
1	A	266	ILE	N-CA-C	5.69	116.08	108.11
1	A	373	GLU	N-CA-C	-5.69	105.44	112.38
1	B	373	GLU	N-CA-C	-5.60	105.71	112.54
1	B	187	LEU	CA-C-N	5.25	125.26	119.90
1	B	187	LEU	C-N-CA	5.25	125.26	119.90
1	A	341	SER	N-CA-C	5.24	117.44	108.90
1	A	382	TRP	CA-C-N	5.14	124.90	119.76
1	A	382	TRP	C-N-CA	5.14	124.90	119.76
1	A	412	LEU	N-CA-C	-5.02	103.45	109.72

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1903	0	1876	42	0
1	B	1903	0	1876	57	0
2	A	42	0	39	1	0
2	B	42	0	39	1	0
3	A	42	0	0	3	0
3	B	23	0	0	0	0
All	All	3955	0	3830	100	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (100) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:412:LEU:HD12	1:B:413:PRO:HD2	1.50	0.92
1:A:350:VAL:HG13	1:A:355:ILE:HD13	1.66	0.78
1:A:231:ARG:HD2	1:A:231:ARG:H	1.50	0.77
1:B:288:VAL:HG12	1:B:359:GLN:HG3	1.74	0.69
1:A:337:ASP:HA	1:A:340:ARG:HD2	1.75	0.68
1:B:288:VAL:CG1	1:B:361:ARG:HB2	2.24	0.67
1:A:251:VAL:HG13	1:A:255:ASP:HB2	1.77	0.67
1:B:215:LEU:HD23	1:B:218:VAL:HG23	1.79	0.65
1:B:252:THR:HG22	1:B:253:ALA:H	1.65	0.62
1:B:288:VAL:HG13	1:B:361:ARG:HB2	1.82	0.61
1:B:252:THR:HB	1:B:254:GLU:HG2	1.82	0.61
1:B:373:GLU:HG3	1:B:441:ARG:NH2	2.16	0.61
1:A:291:THR:HG23	1:A:302:THR:HB	1.83	0.60
1:A:326:ARG:O	1:B:346:ALA:HA	2.01	0.60
1:A:397:ASN:HD22	1:A:398:PRO:HA	1.68	0.59
1:A:251:VAL:CG1	1:A:255:ASP:HB2	2.34	0.58
1:A:231:ARG:HD2	1:A:231:ARG:N	2.18	0.58
1:A:251:VAL:HG11	1:A:280:ILE:HG12	1.85	0.58
1:A:384:GLU:HB3	1:A:450:PRO:HD3	1.86	0.58
1:B:232:LEU:O	1:B:234:PRO:HD3	2.04	0.57
1:B:369:PRO:HG3	1:B:434:SER:HB3	1.87	0.57

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:280:ILE:HD12	1:A:280:ILE:N	2.20	0.56
1:B:382:TRP:CG	1:B:388:GLN:HG3	2.40	0.56
1:B:406:LYS:HG2	1:B:410:PHE:HB3	1.87	0.56
1:A:215:LEU:HD23	1:A:215:LEU:H	1.72	0.55
1:A:198:ARG:HD3	3:A:4:HOH:O	2.07	0.54
1:A:397:ASN:ND2	1:A:398:PRO:HA	2.22	0.54
1:A:390:PRO:HB2	1:A:414:ILE:HA	1.90	0.54
1:B:288:VAL:HG11	1:B:361:ARG:HB2	1.90	0.54
1:A:291:THR:CG2	1:A:302:THR:HB	2.38	0.54
1:B:286:PRO:O	1:B:359:GLN:HB3	2.08	0.53
1:B:200:LEU:O	1:B:280:ILE:HA	2.07	0.53
1:A:187:LEU:HD22	1:A:267:LEU:HB2	1.90	0.53
1:A:258:THR:HG23	1:A:277:THR:HG23	1.91	0.52
1:B:227:LEU:HD22	1:B:249:VAL:HG11	1.92	0.51
1:A:348:LEU:HD12	1:A:349:GLU:H	1.76	0.51
1:B:234:PRO:HA	1:B:247:ALA:HB2	1.93	0.51
1:B:397:ASN:HA	1:B:398:PRO:C	2.34	0.51
1:B:376:CYS:HB2	1:B:392:CYS:HA	1.92	0.51
1:A:200:LEU:O	1:A:280:ILE:HA	2.11	0.50
1:A:399:LEU:HD12	1:A:400:PRO:HD2	1.93	0.50
1:A:304:LYS:HD3	1:A:328:GLN:NE2	2.27	0.50
1:A:422:ARG:NH2	1:A:447:VAL:O	2.44	0.49
1:A:185:PHE:N	3:A:33:HOH:O	2.46	0.49
1:B:373:GLU:HG3	1:B:441:ARG:HH22	1.78	0.48
1:A:397:ASN:HA	1:A:398:PRO:C	2.37	0.48
1:B:281:TYR:CZ	1:B:355:ILE:HD13	2.49	0.48
1:B:384:GLU:HB2	1:B:448:LEU:O	2.14	0.48
1:B:212:LEU:HB3	1:B:243:PHE:CE1	2.48	0.48
2:A:1358:NAG:H4	3:A:14:HOH:O	2.14	0.47
1:B:234:PRO:HB3	1:B:245:ALA:HB1	1.94	0.47
1:B:292:LYS:HD2	1:B:293:PRO:HD2	1.96	0.47
1:B:403:LYS:O	1:B:430:CYS:HA	2.14	0.47
1:A:283:PHE:CE2	1:A:357:LYS:HB2	2.49	0.47
1:A:192:PRO:HA	1:A:212:LEU:HD13	1.96	0.47
1:B:251:VAL:HG11	1:B:280:ILE:HD13	1.97	0.47
1:B:293:PRO:HB2	1:B:364:ARG:HH22	1.79	0.47
1:A:295:VAL:CG1	1:A:299:THR:HB	2.44	0.47
1:B:293:PRO:HB2	1:B:364:ARG:NH2	2.29	0.47
1:B:431:ARG:HG3	1:B:440:THR:HG22	1.96	0.47
2:B:2240:NAG:H83	2:B:2240:NAG:O3	2.14	0.47
1:A:251:VAL:HG13	1:A:255:ASP:CB	2.45	0.46

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:232:LEU:HD21	1:B:249:VAL:HG13	1.97	0.46
1:B:194:LEU:HB3	1:B:276:GLN:NE2	2.31	0.46
1:B:252:THR:HG22	1:B:253:ALA:N	2.32	0.45
1:B:404:CYS:O	1:B:411:PRO:HA	2.16	0.45
1:B:290:LEU:H	1:B:361:ARG:NH1	2.15	0.45
1:B:449:SER:OG	1:B:450:PRO:HD2	2.17	0.45
1:A:348:LEU:HD12	1:A:349:GLU:N	2.32	0.45
1:B:199:VAL:HG12	1:B:279:THR:HB	1.99	0.45
1:B:349:GLU:HA	1:B:353:GLN:O	2.16	0.45
1:B:260:ARG:HG3	1:B:275:LEU:HD13	1.98	0.44
1:B:275:LEU:HG	1:B:276:GLN:N	2.31	0.44
1:A:295:VAL:HG13	1:A:299:THR:HB	1.98	0.44
1:B:292:LYS:HG3	1:B:294:GLU:O	2.18	0.44
1:A:251:VAL:HG12	1:A:252:THR:N	2.33	0.43
1:B:212:LEU:HB3	1:B:243:PHE:HE1	1.82	0.43
1:A:304:LYS:HD3	1:A:328:GLN:HE21	1.83	0.43
1:B:338:ASN:HB2	1:B:365:VAL:HB	1.99	0.43
1:B:275:LEU:HD12	1:B:276:GLN:H	1.83	0.43
1:B:225:LEU:HB3	1:B:232:LEU:HB2	2.01	0.43
1:A:367:TYR:CD1	1:A:367:TYR:N	2.87	0.43
1:B:394:ALA:HB1	1:B:400:PRO:HB2	2.00	0.42
1:B:422:ARG:C	1:B:424:LEU:H	2.26	0.42
1:A:194:LEU:HB3	1:A:276:GLN:NE2	2.35	0.41
1:A:379:ASN:C	1:A:379:ASN:HD22	2.29	0.41
1:B:383:PRO:HA	1:B:448:LEU:HB2	2.03	0.41
1:B:402:LEU:HD12	1:B:403:LYS:H	1.86	0.41
1:B:236:VAL:HA	1:B:244:SER:O	2.21	0.41
1:B:288:VAL:HG12	1:B:359:GLN:CG	2.48	0.41
1:B:375:ASP:HB3	1:B:393:GLN:O	2.21	0.41
1:B:382:TRP:HA	1:B:383:PRO:HD3	1.91	0.41
1:A:355:ILE:N	1:A:355:ILE:HD12	2.36	0.41
1:A:358:ASN:OD1	1:A:358:ASN:C	2.65	0.40
1:A:259:GLN:HG2	1:A:280:ILE:HD13	2.02	0.40
1:A:266:ILE:HG12	1:A:271:SER:HB2	2.02	0.40
1:B:414:ILE:H	1:B:414:ILE:HG13	1.65	0.40
1:A:243:PHE:CD1	1:A:243:PHE:C	2.98	0.40
1:B:382:TRP:CD2	1:B:388:GLN:HG3	2.57	0.40
1:B:390:PRO:HG2	1:B:413:PRO:HG2	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	246/266 (92%)	236 (96%)	7 (3%)	3 (1%)	10	32
1	B	246/266 (92%)	233 (95%)	11 (4%)	2 (1%)	16	42
All	All	492/532 (92%)	469 (95%)	18 (4%)	5 (1%)	12	37

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	240	ASN
1	B	228	GLY
1	B	368	GLY
1	A	368	GLY
1	A	408	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	218/231 (94%)	207 (95%)	11 (5%)	22	49
1	B	218/231 (94%)	204 (94%)	14 (6%)	16	40
All	All	436/462 (94%)	411 (94%)	25 (6%)	18	45

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	190	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	193	GLN
1	A	258	THR
1	A	291	THR
1	A	295	VAL
1	A	350	VAL
1	A	367	TYR
1	A	373	GLU
1	A	379	ASN
1	A	385	ASN
1	A	438	GLU
1	B	225	LEU
1	B	236	VAL
1	B	249	VAL
1	B	260	ARG
1	B	279	THR
1	B	288	VAL
1	B	295	VAL
1	B	296	SER
1	B	300	GLU
1	B	328	GLN
1	B	358	ASN
1	B	385	ASN
1	B	434	SER
1	B	439	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	276	GLN
1	A	379	ASN
1	A	388	GLN
1	A	397	ASN
1	A	436	GLN
1	B	224	HIS
1	B	356	HIS
1	B	359	GLN
1	B	388	GLN
1	B	397	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	B	2358	1	14,14,15	0.76	0	17,19,21	0.80	0
2	NAG	A	1269	1	14,14,15	0.72	1 (7%)	17,19,21	0.74	0
2	NAG	A	1358	1	14,14,15	0.65	0	17,19,21	0.70	0
2	NAG	B	2240	1	14,14,15	1.00	1 (7%)	17,19,21	0.79	0
2	NAG	B	2269	1	14,14,15	0.67	0	17,19,21	0.57	0
2	NAG	A	1240	1	14,14,15	0.78	0	17,19,21	1.29	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	2358	1	-	4/6/23/26	0/1/1/1
2	NAG	A	1269	1	1/1/5/7	3/6/23/26	0/1/1/1
2	NAG	A	1358	1	-	4/6/23/26	0/1/1/1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	2240	1	1/1/5/7	4/6/23/26	0/1/1/1
2	NAG	B	2269	1	-	3/6/23/26	0/1/1/1
2	NAG	A	1240	1	-	2/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2240	NAG	C1-C2	3.01	1.56	1.52
2	A	1269	NAG	C1-C2	2.11	1.55	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1240	NAG	C3-C4-C5	2.37	114.53	110.23
2	A	1240	NAG	O5-C1-C2	-2.36	107.64	111.29
2	A	1240	NAG	C2-N2-C7	-2.19	119.97	122.90
2	A	1240	NAG	C4-C3-C2	-2.04	108.03	111.02

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	1269	NAG	C1
2	B	2240	NAG	C1

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1240	NAG	C8-C7-N2-C2
2	A	1240	NAG	O7-C7-N2-C2
2	A	1269	NAG	C8-C7-N2-C2
2	A	1269	NAG	O7-C7-N2-C2
2	A	1358	NAG	C8-C7-N2-C2
2	A	1358	NAG	O7-C7-N2-C2
2	B	2358	NAG	C8-C7-N2-C2
2	B	2358	NAG	O7-C7-N2-C2
2	B	2240	NAG	C8-C7-N2-C2
2	B	2240	NAG	O7-C7-N2-C2
2	B	2269	NAG	C3-C2-N2-C7
2	B	2269	NAG	C8-C7-N2-C2
2	B	2269	NAG	O7-C7-N2-C2
2	B	2358	NAG	O5-C5-C6-O6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
2	B	2358	NAG	C4-C5-C6-O6
2	A	1358	NAG	C4-C5-C6-O6
2	A	1358	NAG	O5-C5-C6-O6
2	B	2240	NAG	C3-C2-N2-C7
2	B	2240	NAG	C4-C5-C6-O6
2	A	1269	NAG	C4-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1358	NAG	1	0
2	B	2240	NAG	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	250/266 (93%)	-0.21	8 (3%) 50 28	9, 26, 52, 59	0
1	B	250/266 (93%)	-0.03	3 (1%) 76 55	9, 34, 57, 59	0
All	All	500/532 (93%)	-0.12	11 (2%) 62 39	9, 30, 55, 59	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	324	GLY	3.6
1	B	450	PRO	3.3
1	A	368	GLY	3.2
1	A	326	ARG	3.2
1	B	287	ASN	2.6
1	B	229	ASP	2.6
1	A	450	PRO	2.4
1	A	307	ALA	2.3
1	A	240	ASN	2.3
1	A	306	GLU	2.1
1	A	325	PRO	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	A	1240	14/15	0.47	0.26	58,58,58,58	0
2	NAG	B	2240	14/15	0.64	0.26	58,58,58,58	0
2	NAG	B	2358	14/15	0.70	0.25	58,58,58,58	0
2	NAG	A	1358	14/15	0.71	0.18	58,58,58,58	0
2	NAG	B	2269	14/15	0.78	0.23	58,58,58,58	0
2	NAG	A	1269	14/15	0.82	0.17	58,58,58,58	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.