



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 12:43 PM UTC

PDB ID : 6P50 / pdb_00006p50
Title : Crystal Structure of a Complex of human IL-7Ralpha with an anti-IL-7Ralpha Fab 4A10
Authors : Walsh, S.T.R.; Kashi, L.; Kohnhorst, C.L.
Deposited on : 2019-05-29
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	223	
2	H	225	
3	L	213	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3990 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interleukin-7 receptor subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	90	Total	C	N	O	S	0	0	0
			679	440	110	123	6			

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-3	GLY	-	expression tag	UNP P16871
C	-2	SER	-	expression tag	UNP P16871
C	-1	HIS	-	expression tag	UNP P16871
C	0	MET	-	expression tag	UNP P16871
C	46	THR	ILE	conflict	UNP P16871

- Molecule 2 is a protein called anti-IL-7R 4A10 Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	221	Total	C	N	O	S	0	0	0
			1668	1059	269	331	9			

- Molecule 3 is a protein called anti-IL-7R 4A10 Fab light chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	211	Total	C	N	O	S	0	0	0
			1629	1019	278	326	6			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	C	1	14	8	1	5	0	0

- Molecule 1: Interleukin-7 receptor subunit alpha



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	145.11Å 40.17Å 105.10Å 90.00° 105.48° 90.00°	Depositor
Resolution (Å)	38.61 – 2.90 38.61 – 2.90	Depositor EDS
% Data completeness (in resolution range)	96.0 (38.61-2.90) 96.3 (38.61-2.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.33 (at 2.90Å)	Xtriage
Refinement program	PHENIX 1.16rc1_3535	Depositor
R, R_{free}	0.285 , 0.330 0.291 , 0.342	Depositor DCC
R_{free} test set	644 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	62.5	Xtriage
Anisotropy	0.506	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 65.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	3990	wwPDB-VP
Average B, all atoms (Å ²)	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	C	0.09	0/689	0.30	0/931
2	H	0.09	0/1714	0.26	0/2345
3	L	0.09	0/1667	0.32	0/2265
All	All	0.09	0/4070	0.29	0/5541

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	679	0	664	11	0
2	H	1668	0	1603	13	0
3	L	1629	0	1564	13	0
4	C	14	0	13	0	0
All	All	3990	0	3844	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:11:LEU:HD22	2:H:156:PRO:HG3	1.70	0.71
2:H:99:ARG:NH2	2:H:102:SER:O	2.32	0.62
3:L:18:LYS:HA	3:L:75:ILE:O	2.01	0.60
3:L:165:GLN:NE2	3:L:170:SER:OG	2.36	0.58
1:C:73:PHE:HE1	1:C:75:GLU:HB2	1.69	0.57
3:L:49:HIS:O	3:L:53:THR:OG1	2.19	0.57
2:H:38:LYS:HB2	2:H:48:ILE:HD11	1.86	0.57
1:C:76:THR:HG22	1:C:78:LYS:H	1.71	0.56
1:C:51:PHE:O	1:C:64:ASN:ND2	2.41	0.53
2:H:152:LYS:HG2	2:H:185:THR:HG23	1.91	0.53
3:L:67:SER:O	3:L:69:ARG:N	2.42	0.52
1:C:99:LYS:HD3	1:C:100:LYS:H	1.74	0.51
2:H:144:MET:HE3	2:H:191:THR:HG22	1.94	0.50
3:L:149:ILE:HD11	3:L:178:LEU:HD21	1.94	0.49
3:L:19:VAL:HG13	3:L:78:LEU:HD11	1.94	0.48
3:L:183:ASP:O	3:L:187:ARG:HG3	2.14	0.48
1:C:35:LEU:HB3	1:C:74:ILE:HG23	1.96	0.48
3:L:83:ILE:HG21	3:L:165:GLN:HB3	1.96	0.47
2:H:128:PRO:HA	2:H:154:TYR:HB3	1.97	0.47
3:L:50:TYR:O	3:L:52:SER:N	2.44	0.47
2:H:208:HIS:CE1	2:H:210:ALA:HB3	2.50	0.47
1:C:44:VAL:HG12	1:C:46:THR:H	1.80	0.46
3:L:168:LYS:C	3:L:170:SER:H	2.24	0.45
2:H:174:THR:HG23	2:H:186:LEU:HD21	1.97	0.45
1:C:25:GLN:HB2	1:C:34:SER:OG	2.16	0.45
3:L:192:THR:HG23	3:L:207:SER:HB2	1.99	0.44
1:C:53:ILE:HD11	1:C:85:SER:HB3	2.00	0.43
1:C:90:LYS:HE3	2:H:103:ASN:ND2	2.34	0.43
2:H:156:PRO:HB2	2:H:157:GLU:H	1.72	0.41
1:C:84:LYS:O	2:H:105:TYR:OH	2.33	0.41
1:C:47:THR:C	1:C:49:LEU:H	2.28	0.41
2:H:160:THR:OG1	2:H:207:ALA:HB3	2.20	0.41
3:L:129:ALA:O	3:L:180:LEU:N	2.54	0.41
2:H:55:ASP:OD1	2:H:55:ASP:N	2.52	0.40
3:L:3:GLN:HB3	3:L:26:SER:HB3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	86/223 (39%)	75 (87%)	11 (13%)	0	100	100
2	H	217/225 (96%)	202 (93%)	13 (6%)	2 (1%)	14	41
3	L	209/213 (98%)	191 (91%)	16 (8%)	2 (1%)	12	39
All	All	512/661 (78%)	468 (91%)	40 (8%)	4 (1%)	16	44

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	L	68	GLY
2	H	221	PRO
3	L	137	ASN
2	H	156	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	74/204 (36%)	70 (95%)	4 (5%)	20	51
2	H	188/195 (96%)	187 (100%)	1 (0%)	81	93
3	L	183/189 (97%)	179 (98%)	4 (2%)	45	77
All	All	445/588 (76%)	436 (98%)	9 (2%)	48	78

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	61	LYS
1	C	63	LEU
1	C	64	ASN
1	C	99	LYS
2	H	6	GLN
3	L	23	CYS
3	L	69	ARG
3	L	141	ARG
3	L	148	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	C	301	1	14,14,15	0.28	0	17,19,21	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	C	301	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	C	301	NAG	O5-C5-C6-O6
4	C	301	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	C	90/223 (40%)	0.99	6 (6%) 24 19	66, 82, 104, 127	0
2	H	221/225 (98%)	1.03	24 (10%) 10 9	38, 66, 94, 121	0
3	L	211/213 (99%)	1.01	24 (11%) 10 8	35, 68, 92, 107	0
All	All	522/661 (78%)	1.01	54 (10%) 12 10	35, 70, 98, 127	0

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	L	129	ALA	4.9
3	L	116	ILE	4.7
3	L	204	ILE	3.9
3	L	115	SER	3.3
2	H	112	TRP	3.2
3	L	35	TRP	3.2
3	L	88	CYS	3.2
2	H	198	PRO	3.1
3	L	74	SER	2.9
2	H	208	HIS	2.9
3	L	38	HIS	2.9
2	H	18	VAL	2.8
3	L	26	SER	2.7
2	H	204	CYS	2.7
2	H	146	THR	2.7
2	H	136	GLY	2.7
2	H	35	HIS	2.6
1	C	54	CYS	2.6
3	L	85	THR	2.6
2	H	37	VAL	2.6
2	H	125	THR	2.6
3	L	94	LEU	2.6
1	C	41	ASP	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	L	21	ILE	2.5
3	L	143	ILE	2.5
3	L	73	PHE	2.5
2	H	9	ALA	2.5
3	L	148	LYS	2.5
2	H	79	ALA	2.4
3	L	103	LEU	2.4
2	H	121	SER	2.3
1	C	63	LEU	2.3
2	H	178	VAL	2.3
3	L	1	ASP	2.3
3	L	25	ALA	2.3
2	H	159	VAL	2.3
3	L	118	PRO	2.3
2	H	111	TYR	2.2
3	L	132	VAL	2.2
2	H	107	TYR	2.2
1	C	33	HIS	2.2
3	L	136	ASN	2.2
2	H	149	CYS	2.2
3	L	23	CYS	2.2
1	C	81	LEU	2.2
2	H	30	THR	2.1
1	C	60	VAL	2.1
3	L	201	THR	2.1
2	H	88	SER	2.1
2	H	2	VAL	2.1
2	H	151	VAL	2.1
3	L	95	LEU	2.1
2	H	34	MET	2.0
2	H	216	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	C	301	14/15	0.60	0.14	76,81,86,88	0

6.5 Other polymers

There are no such residues in this entry.