



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 11:55 PM UTC

PDB ID : 8P5I / pdb_00008p5i
Title : Kinase domain of mutant human ULK1 in complex with compound XMD-17-51
Authors : Battista, T.; Semrau, M.S.; Heroux, A.; Lolli, G.; Storici, P.
Deposited on : 2023-05-24
Resolution : 1.83 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.83 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1112 (1.82-1.82)
Clashscore	190562	1148 (1.82-1.82)
Ramachandran outliers	187476	1140 (1.82-1.82)
Sidechain outliers	187428	1140 (1.82-1.82)
RSRZ outliers	180081	1112 (1.82-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	284	
1	B	284	
1	C	284	
1	D	284	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 19508 atoms, of which 9429 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Serine/threonine-protein kinase ULK1.

Mol	Chain	Residues	Atoms								ZeroOcc	AltConf	Trace
1	A	269	Total	C	H	N	O	P	S		63	17	0
			4548	1442	2316	380	392	1	17				
1	B	273	Total	C	H	N	O	P	S		62	16	0
			4602	1457	2341	390	397	1	16				
1	C	273	Total	C	H	N	O	P	S		61	18	0
			4639	1468	2365	393	396	1	16				
1	D	267	Total	C	H	N	O	P	S		61	15	0
			4505	1428	2295	376	388	1	17				

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP O75385
A	245	ALA	ARG	engineered mutation	UNP O75385
A	246	ALA	GLU	engineered mutation	UNP O75385
B	0	SER	-	expression tag	UNP O75385
B	245	ALA	ARG	engineered mutation	UNP O75385
B	246	ALA	GLU	engineered mutation	UNP O75385
C	0	SER	-	expression tag	UNP O75385
C	245	ALA	ARG	engineered mutation	UNP O75385
C	246	ALA	GLU	engineered mutation	UNP O75385
D	0	SER	-	expression tag	UNP O75385
D	245	ALA	ARG	engineered mutation	UNP O75385
D	246	ALA	GLU	engineered mutation	UNP O75385

- Molecule 2 is 5,11-dimethyl-2-[(1-piperidin-4-ylpyrazol-4-yl)amino]pyrimido[4,5-b][1,4]benzodiazepin-6-one (CCD ID: WYX) (formula: C₂₁H₂₄N₈O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	0	0
			54	21	24	8	1		
2	B	1	Total	C	H	N	O	0	0
			54	21	24	8	1		
2	C	1	Total	C	H	N	O	0	0
			54	21	24	8	1		
2	D	1	Total	C	H	N	O	0	0
			54	21	24	8	1		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Mg	0	0
			2	2		
3	B	2	Total	Mg	0	0
			2	2		
3	C	2	Total	Mg	0	0
			2	2		
3	D	2	Total	Mg	0	0
			2	2		

- Molecule 4 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Na	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Na	0	0
			1	1		
4	D	1	Total	Na	0	0
			1	1		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	H	O	2	0
			14	3	8	3		
5	C	1	Total	C	H	O	2	0
			14	3	8	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	243	Total	O	0	0
			243	243		
6	B	265	Total	O	0	0
			265	265		
6	C	233	Total	O	0	0
			233	233		
6	D	218	Total	O	0	0
			218	218		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Serine/threonine-protein kinase ULK1




- Molecule 1: Serine/threonine-protein kinase ULK1




- Molecule 1: Serine/threonine-protein kinase ULK1



- Molecule 1: Serine/threonine-protein kinase ULK1





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	85.47Å 109.54Å 96.52Å 90.00° 92.84° 90.00°	Depositor
Resolution (Å)	96.40 – 1.83 96.40 – 1.83	Depositor EDS
% Data completeness (in resolution range)	99.9 (96.40-1.83) 99.9 (96.40-1.83)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.31 (at 1.83Å)	Xtriage
Refinement program	REFMAC 5.8.0405	Depositor
R, R_{free}	0.166 , 0.189 0.166 , 0.190	Depositor DCC
R_{free} test set	7671 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	27.6	Xtriage
Anisotropy	0.028	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 40.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.056 for h,-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	19508	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, MG, NA, WYX, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.70	0/2314	0.95	2/3119 (0.1%)
1	B	0.70	0/2341	0.95	1/3154 (0.0%)
1	C	0.67	0/2360	0.94	3/3181 (0.1%)
1	D	0.68	0/2288	0.95	0/3082
All	All	0.69	0/9303	0.95	6/12536 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	2
1	C	0	1
All	All	0	4

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	156	PRO	N-CA-C	5.74	122.13	113.81
1	C	203	ASP	CA-CB-CG	5.37	117.97	112.60
1	C	155	ASN	CB-CA-C	5.33	116.54	108.86
1	B	203	ASP	CA-CB-CG	5.28	117.88	112.60
1	A	8	THR	CA-CB-OG1	-5.25	101.72	109.60
1	C	108	ARG	CB-CG-CD	5.10	123.02	111.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	279	ASP	Peptide
1	B	153	ARG	Sidechain
1	B	36	ARG	Sidechain
1	C	153	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2232	2316	2318	24	0
1	B	2261	2341	2343	20	0
1	C	2274	2365	2369	18	0
1	D	2210	2295	2297	22	0
2	A	30	24	0	0	0
2	B	30	24	0	0	0
2	C	30	24	0	0	0
2	D	30	24	0	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	D	1	0	0	0	0
5	B	6	8	8	0	0
5	C	6	8	8	0	0
6	A	243	0	0	6	0
6	B	265	0	0	8	0
6	C	233	0	0	3	0
6	D	218	0	0	5	0
All	All	10079	9429	9343	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:184[A]:SER:OG	6:B:401:HOH:O	1.62	1.00
1:C:175:ASN:O	1:C:176[B]:MET:HE2	1.69	0.92
1:A:184[A]:SER:OG	6:A:401:HOH:O	1.95	0.84
1:A:176[B]:MET:HE1	6:A:591:HOH:O	1.82	0.78
1:B:121[B]:GLN:HE22	1:B:160:ARG:HA	1.52	0.75
1:D:176[A]:MET:HE1	6:D:563:HOH:O	1.86	0.74
1:A:176[B]:MET:SD	6:A:598:HOH:O	2.46	0.72
1:B:168[B]:PHE:CD2	6:B:470:HOH:O	2.43	0.72
1:A:13[A]:LYS:HD2	1:A:35[A]:HIS:NE2	2.05	0.72
1:D:173:GLN:CB	1:D:176[A]:MET:HE3	2.20	0.72
1:B:168[B]:PHE:HD2	6:B:470:HOH:O	1.72	0.71
1:B:74:ASN:HD21	1:B:121[B]:GLN:HE21	1.38	0.71
1:D:176[A]:MET:SD	6:D:577:HOH:O	2.49	0.70
1:A:173:GLN:CB	1:A:176[B]:MET:HE3	2.21	0.70
1:A:268:ASP:HB2	6:A:559:HOH:O	1.92	0.69
1:D:221:PHE:CE1	1:D:239[B]:LEU:HD11	2.29	0.67
1:D:268:ASP:HB2	6:D:549:HOH:O	1.94	0.67
1:D:175:ASN:O	1:D:176[B]:MET:HE2	1.95	0.66
1:D:173:GLN:HB3	1:D:176[A]:MET:HE3	1.79	0.63
1:A:173:GLN:HB2	1:A:176[B]:MET:HE3	1.81	0.63
1:A:267[B]:MET:HE1	1:A:275:HIS:CD2	2.34	0.62
1:D:173:GLN:HB2	1:D:176[A]:MET:HE3	1.83	0.59
1:A:173:GLN:HB3	1:A:176[B]:MET:HE3	1.83	0.59
1:A:205:TRP:O	1:A:209[A]:THR:HG23	2.04	0.58
1:C:205:TRP:O	1:C:209[B]:THR:HG23	2.03	0.58
1:C:227:GLN:O	1:C:231:LEU:HD23	2.05	0.57
1:D:205:TRP:O	1:D:209[A]:THR:HG23	2.04	0.57
1:B:205:TRP:O	1:B:209[A]:THR:HG23	2.04	0.57
1:C:149:PRO:O	1:C:153:ARG:HD3	2.05	0.56
1:A:48:ILE:HD11	1:A:88[B]:VAL:HG12	1.87	0.55
1:A:227:GLN:HG3	1:A:230:ARG:NH2	2.22	0.55
1:C:21:LEU:CD1	1:C:29[A]:VAL:HG13	2.38	0.54
1:D:48:ILE:HD11	1:D:88[B]:VAL:HG12	1.90	0.54
1:C:209[B]:THR:HG21	6:C:455:HOH:O	2.08	0.54
1:B:21:LEU:CD1	1:B:29[A]:VAL:HG13	2.37	0.53
1:B:74:ASN:ND2	1:B:121[B]:GLN:HE21	2.04	0.53
1:D:184[A]:SER:OG	6:D:401:HOH:O	2.12	0.53
1:B:62:LYS:NZ	6:B:402:HOH:O	2.41	0.53
1:A:76[B]:VAL:HG11	1:A:164:ALA:HB2	1.92	0.52
1:B:145:LEU:O	1:B:146[A]:LEU:HD13	2.10	0.51
1:C:62:LYS:NZ	6:C:401:HOH:O	2.45	0.50
1:C:257:ALA:HB3	1:C:267:MET:HE3	1.92	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:48:ILE:HD11	1:C:88[B]:VAL:HG12	1.93	0.49
1:B:268:ASP:HB2	6:B:573:HOH:O	2.12	0.49
1:B:21:LEU:HD11	1:B:29[A]:VAL:HG13	1.94	0.49
1:A:173:GLN:CD	1:A:176[B]:MET:HE2	2.38	0.48
1:D:267[B]:MET:HE1	1:D:275:HIS:CD2	2.49	0.48
1:C:243:ILE:O	1:C:252[B]:ARG:NH1	2.47	0.48
1:A:35[A]:HIS:HD1	1:A:37:GLU:H	1.62	0.47
1:A:173:GLN:NE2	1:A:176[B]:MET:HE2	2.29	0.47
1:D:257:ALA:HB3	1:D:267[A]:MET:HE3	1.97	0.47
1:B:48:ILE:HD11	1:B:88[B]:VAL:HG12	1.96	0.46
1:B:149:PRO:O	1:B:153:ARG:HD3	2.15	0.46
1:C:76[B]:VAL:HG11	1:C:164:ALA:HB2	1.97	0.46
1:D:267[B]:MET:HE1	1:D:275:HIS:HD2	1.81	0.46
1:C:21:LEU:HD11	1:C:29[A]:VAL:HG13	1.96	0.46
1:D:121:GLN:HB2	1:D:161[A]:VAL:CG2	2.45	0.46
1:D:173:GLN:CD	1:D:176[A]:MET:HE2	2.41	0.46
1:C:13:LYS:HA	1:C:13:LYS:HE2	1.98	0.45
1:D:173:GLN:NE2	1:D:176[A]:MET:HE2	2.31	0.45
1:B:184[B]:SER:HB2	6:B:401:HOH:O	2.15	0.45
1:A:127:ARG:HA	1:A:273:PHE:CZ	2.52	0.44
1:D:21:LEU:CD1	1:D:29:VAL:HG13	2.48	0.44
1:D:127:ARG:HA	1:D:273:PHE:CZ	2.52	0.43
1:A:257:ALA:HB3	1:A:267[A]:MET:CE	2.48	0.43
1:D:76[B]:VAL:HG11	1:D:164:ALA:HB2	2.00	0.43
1:A:186:MET:HB2	6:A:614:HOH:O	2.19	0.43
1:D:173:GLN:HB3	1:D:176[A]:MET:CE	2.48	0.43
1:B:142:GLN:HG2	6:B:547:HOH:O	2.18	0.43
1:B:121[A]:GLN:OE1	1:B:161[A]:VAL:HG13	2.19	0.42
1:C:144:ILE:HG23	1:C:161[B]:VAL:HG13	2.01	0.42
1:A:21:LEU:CD1	1:A:29[A]:VAL:HG13	2.49	0.42
1:B:257:ALA:HB3	1:B:267:MET:HE3	2.02	0.42
1:B:71:LYS:NZ	6:B:406:HOH:O	2.51	0.41
1:A:173:GLN:HB3	1:A:176[B]:MET:CE	2.50	0.41
1:B:153:ARG:HG2	1:B:158:SER:CB	2.50	0.41
1:C:127[A]:ARG:HA	1:C:273:PHE:CZ	2.56	0.41
1:C:127[B]:ARG:HA	1:C:273:PHE:CZ	2.56	0.41
1:A:29[B]:VAL:HG23	1:A:47:CYS:HB2	2.01	0.40
1:A:249:ALA:HB3	1:A:250:PRO:HD3	2.02	0.40
1:D:209[A]:THR:HG21	6:D:501:HOH:O	2.22	0.40
1:C:268:ASP:HB2	6:C:555:HOH:O	2.20	0.40
1:A:176[A]:MET:HE3	6:A:529:HOH:O	2.21	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:160:ARG:HD3	1:C:160:ARG:HA	1.93	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	281/284 (99%)	273 (97%)	8 (3%)	0	100	100
1	B	286/284 (101%)	275 (96%)	11 (4%)	0	100	100
1	C	288/284 (101%)	277 (96%)	11 (4%)	0	100	100
1	D	277/284 (98%)	268 (97%)	9 (3%)	0	100	100
All	All	1132/1136 (100%)	1093 (97%)	39 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	247/240 (103%)	240 (97%)	7 (3%)	38	21
1	B	248/240 (103%)	243 (98%)	5 (2%)	48	35
1	C	250/240 (104%)	244 (98%)	6 (2%)	43	27
1	D	245/240 (102%)	239 (98%)	6 (2%)	43	27

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	990/960 (103%)	966 (98%)	24 (2%)	50 27

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	HIS
1	A	147	SER
1	A	155	ASN
1	A	161[A]	VAL
1	A	161[B]	VAL
1	A	162	LYS
1	A	224	SER
1	B	146[A]	LEU
1	B	146[B]	LEU
1	B	161[A]	VAL
1	B	161[B]	VAL
1	B	218	LYS
1	C	127[A]	ARG
1	C	127[B]	ARG
1	C	155	ASN
1	C	161[A]	VAL
1	C	161[B]	VAL
1	C	218	LYS
1	D	50	LYS
1	D	147	SER
1	D	155	ASN
1	D	162	LYS
1	D	239[A]	LEU
1	D	239[B]	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	86	ASN
1	A	157	ASN
1	A	173	GLN
1	B	86	ASN
1	B	173	GLN
1	B	263	HIS
1	C	86	ASN
1	C	173	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	263	HIS
1	D	86	ASN
1	D	173	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	TPO	C	180	1	8,10,11	1.63	1 (12%)	10,14,16	1.34	3 (30%)
1	TPO	A	180	1	8,10,11	1.22	1 (12%)	10,14,16	1.39	2 (20%)
1	TPO	D	180	1	8,10,11	1.38	2 (25%)	10,14,16	1.40	2 (20%)
1	TPO	B	180	1	8,10,11	1.50	1 (12%)	10,14,16	1.15	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	C	180	1	-	0/9/11/13	-
1	TPO	A	180	1	-	0/9/11/13	-
1	TPO	D	180	1	-	0/9/11/13	-
1	TPO	B	180	1	-	2/9/11/13	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	180	TPO	P-OG1	3.76	1.66	1.59
1	C	180	TPO	P-OG1	3.52	1.65	1.59
1	D	180	TPO	P-O3P	-2.41	1.45	1.54
1	D	180	TPO	P-OG1	2.22	1.63	1.59
1	A	180	TPO	P-O3P	-2.01	1.47	1.54

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	180	TPO	OG1-P-O1P	-2.68	99.78	109.33
1	D	180	TPO	O3P-P-O2P	2.50	117.16	107.80
1	D	180	TPO	P-OG1-CB	-2.35	116.94	123.33
1	C	180	TPO	P-OG1-CB	-2.27	117.15	123.33
1	C	180	TPO	OG1-P-O1P	-2.22	101.43	109.33
1	B	180	TPO	P-OG1-CB	-2.17	117.43	123.33
1	A	180	TPO	P-OG1-CB	-2.12	117.56	123.33
1	C	180	TPO	O3P-P-O2P	2.11	115.73	107.80

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	180	TPO	CB-OG1-P-O2P
1	B	180	TPO	O-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 11 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	WYX	B	301	-	30,34,34	0.57	0	37,49,49	1.83	6 (16%)
5	GOL	C	302	-	5,5,5	0.42	0	5,5,5	0.62	0
2	WYX	D	301	-	30,34,34	0.65	1 (3%)	37,49,49	2.03	7 (18%)
2	WYX	A	301	-	30,34,34	0.73	1 (3%)	37,49,49	1.88	5 (13%)
5	GOL	B	302	-	5,5,5	0.50	0	5,5,5	0.71	0
2	WYX	C	301	-	30,34,34	0.54	0	37,49,49	1.57	4 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	WYX	B	301	-	-	0/8/16/16	0/5/5/5
5	GOL	C	302	-	-	0/4/4/4	-
2	WYX	D	301	-	-	3/8/16/16	0/5/5/5
2	WYX	A	301	-	-	1/8/16/16	0/5/5/5
5	GOL	B	302	-	-	0/4/4/4	-
2	WYX	C	301	-	-	0/8/16/16	0/5/5/5

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	WYX	C2-N1	2.66	1.46	1.43
2	D	301	WYX	C2-N1	2.20	1.45	1.43

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	301	WYX	C9-C12-N1	9.44	126.21	120.14
2	A	301	WYX	C9-C12-N1	8.62	125.68	120.14
2	B	301	WYX	C9-C12-N1	7.52	124.98	120.14
2	C	301	WYX	C9-C12-N1	7.17	124.75	120.14
2	B	301	WYX	C14-N6-N7	-3.83	109.88	112.45
2	D	301	WYX	C15-C13-N5	3.80	136.44	125.55
2	B	301	WYX	C15-N7-N6	3.61	105.99	104.14
2	C	301	WYX	C15-N7-N6	3.35	105.86	104.14
2	A	301	WYX	C9-C12-N4	-2.94	118.66	123.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	WYX	C15-N7-N6	2.75	105.55	104.14
2	D	301	WYX	C21-N2-C9	-2.73	114.10	118.38
2	D	301	WYX	C9-C12-N4	-2.58	119.23	123.32
2	D	301	WYX	C17-C16-N6	2.55	114.44	110.95
2	D	301	WYX	C1-N1-C2	2.54	117.81	115.95
2	A	301	WYX	C20-C19-N8	2.35	115.25	110.81
2	B	301	WYX	C17-C18-N8	2.34	115.22	110.81
2	B	301	WYX	C19-C20-C16	2.34	115.03	110.77
2	A	301	WYX	C14-N6-N7	-2.27	110.92	112.45
2	D	301	WYX	C15-N7-N6	2.12	105.23	104.14
2	C	301	WYX	C14-N6-N7	-2.12	111.03	112.45
2	C	301	WYX	C1-N1-C2	2.08	117.47	115.95
2	B	301	WYX	C9-C12-N4	-2.04	120.08	123.32

There are no chirality outliers.


All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	301	WYX	C14-C13-N5-C11
2	D	301	WYX	C14-C13-N5-C11
2	D	301	WYX	C17-C16-N6-C14
2	D	301	WYX	C17-C16-N6-N7


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	268/284 (94%)	-0.16	9 (3%)	48	55	12, 28, 56, 83	17 (6%)
1	B	272/284 (95%)	-0.31	6 (2%)	62	69	10, 27, 58, 111	16 (5%)
1	C	272/284 (95%)	-0.16	11 (4%)	42	48	11, 28, 63, 113	18 (6%)
1	D	266/284 (93%)	-0.05	13 (4%)	35	41	12, 29, 67, 81	15 (5%)
All	All	1078/1136 (94%)	-0.17	39 (3%)	46	53	10, 28, 63, 113	66 (6%)

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	280	ALA	7.4
1	A	149	PRO	5.7
1	D	149	PRO	5.7
1	A	154	ALA	5.2
1	B	280	ALA	4.8
1	C	280	ALA	4.1
1	C	155	ASN	4.0
1	D	41	LEU	3.8
1	B	151	GLY	3.8
1	C	231	LEU	3.8
1	D	24	HIS	3.5
1	D	156	PRO	3.4
1	D	8	THR	3.4
1	C	8	THR	3.3
1	B	8	THR	3.2
1	A	24	HIS	3.0
1	A	8	THR	2.8
1	C	38	LYS	2.8
1	C	154	ALA	2.7
1	A	50	LYS	2.7
1	C	151	GLY	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	51	LYS	2.6
1	D	35	HIS	2.6
1	C	41	LEU	2.5
1	B	152	ARG	2.4
1	D	36	ARG	2.4
1	A	222	GLN	2.4
1	D	222	GLN	2.3
1	D	155	ASN	2.3
1	D	39	HIS	2.3
1	B	231	LEU	2.2
1	B	154	ALA	2.2
1	C	152	ARG	2.2
1	A	19	LYS	2.2
1	C	235	LYS	2.2
1	D	37	GLU	2.2
1	D	40	ASP	2.2
1	D	22	ILE	2.1
1	A	186	MET	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	TPO	D	180	11/12	0.98	0.05	19,20,23,24	0
1	TPO	B	180	11/12	0.99	0.03	18,20,21,22	0
1	TPO	C	180	11/12	0.99	0.04	19,21,22,23	0
1	TPO	A	180	11/12	0.99	0.04	18,20,22,22	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NA	B	305	1/1	0.80	0.29	59,59,59,59	0
4	NA	A	304	1/1	0.87	0.23	53,53,53,53	0
5	GOL	B	302	6/6	0.88	0.13	37,41,46,46	2
5	GOL	C	302	6/6	0.91	0.12	38,46,47,51	2
4	NA	D	304	1/1	0.93	0.14	41,41,41,41	0
2	WYX	D	301	30/30	0.95	0.10	25,34,80,81	0
2	WYX	A	301	30/30	0.97	0.08	22,31,67,69	0
2	WYX	B	301	30/30	0.98	0.04	18,21,24,24	0
2	WYX	C	301	30/30	0.98	0.04	20,23,26,27	0
3	MG	C	304	1/1	0.99	0.12	30,30,30,30	0
3	MG	D	302	1/1	0.99	0.09	30,30,30,30	0
3	MG	D	303	1/1	0.99	0.09	33,33,33,33	0
3	MG	A	302	1/1	0.99	0.07	29,29,29,29	0
3	MG	A	303	1/1	0.99	0.11	32,32,32,32	0
3	MG	B	303	1/1	0.99	0.05	35,35,35,35	0
3	MG	B	304	1/1	0.99	0.09	30,30,30,30	0
3	MG	C	303	1/1	0.99	0.06	34,34,34,34	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around WYX D 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around WYX A 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around WYX B 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around WYX C 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.