



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 12:28 PM UTC

PDB ID : 8P6G / pdb_00008p6g
Title : Crystal structure of the improved version of the Genetically Encoded Green Calcium Indicator YTnC2-5
Authors : Boyko, K.M.; Nikolaeva, A.Y.; Subach, O.M.; Vlaskina, A.V.; Agapova, Y.K.; Popov, V.O.; Subach, F.V.
Deposited on : 2023-05-26
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	314	
1	D	314	

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 4137 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Genetically Encoded Green Calcium Indicator YTnC2-5 based on Troponin C from toadfish.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	245	Total	C	N	O	S	0	2	0
			1959	1245	336	369	9			
1	D	245	Total	C	N	O	S	0	3	0
			1950	1242	334	365	9			

- Molecule 2 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			7	4	3		
2	D	1	Total	C	O	0	0
			7	4	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	106	Total 106	O 106	0	0
3	D	108	Total 108	O 108	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Genetically Encoded Green Calcium Indicator YTnC2-5 based on Troponin C from toadfish



- Molecule 1: Genetically Encoded Green Calcium Indicator YTnC2-5 based on Troponin C from toadfish



4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, α , β , γ	65.74Å 65.74Å 315.04Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.14 – 1.95 46.14 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.8 (46.14-1.95) 99.8 (46.14-1.95)	Depositor EDS
R_{merge}	0.31	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.29 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.8.0411	Depositor
R, R_{free}	0.183 , 0.229 0.188 , 0.229	Depositor DCC
R_{free} test set	2837 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	30.2	Xtriage
Anisotropy	0.284	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 15.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.479 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4137	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, CRO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	B	0.78	0/1992	1.32	11/2689 (0.4%)
1	D	0.78	0/1986	1.29	11/2683 (0.4%)
All	All	0.78	0/3978	1.30	22/5372 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	3

There are no bond length outliers.

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	219	VAL	N-CA-CB	-8.47	100.66	112.35
1	B	219	VAL	CB-CA-C	8.38	120.02	110.88
1	D	219	VAL	N-CA-CB	-7.93	102.04	112.26
1	B	293	ASP	CA-CB-CG	6.94	119.54	112.60
1	B	212	PHE	N-CA-C	6.20	117.70	111.07
1	B	274	ASP	CA-CB-CG	5.98	118.58	112.60
1	D	211	GLU	CA-C-O	-5.92	114.28	120.55
1	D	212	PHE	N-CA-CB	5.87	118.69	109.94
1	D	257	ASP	CA-CB-CG	5.82	118.42	112.60
1	D	219	VAL	CB-CA-C	5.78	118.36	111.20
1	D	232	ASP	CA-CB-CG	5.71	118.31	112.60
1	B	280	HIS	CA-CB-CG	5.65	119.45	113.80
1	D	43	TYR	CA-CB-CG	5.51	123.82	113.90
1	D	137	ASP	CA-CB-CG	5.36	117.96	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	103	PHE	CA-CB-CG	5.34	119.14	113.80
1	D	136	GLU	CB-CG-CD	5.33	121.67	112.60
1	B	212	PHE	N-CA-CB	-5.25	102.39	110.01
1	B	267	ASP	CA-CB-CG	5.17	117.77	112.60
1	B	264	PRO	N-CA-CB	5.11	107.86	103.31
1	D	225	HIS	CB-CA-C	5.11	118.31	110.14
1	B	37	GLY	CA-C-O	-5.09	117.40	120.91
1	B	241	HIS	CB-CG-CD2	-5.00	124.69	131.20

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	126[A]	ARG	Sidechain
1	D	126[B]	ARG	Sidechain
1	D	279	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1959	0	1864	19	0
1	D	1950	0	1855	19	0
2	B	7	0	10	3	0
2	D	7	0	10	2	0
3	B	106	0	0	2	0
3	D	108	0	0	1	0
All	All	4137	0	3739	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:209:PHE:O	1:B:213:LEU:HD23	1.97	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:296:VAL:HG13	2:D:401:PEG:H31	1.79	0.63
1:D:275:ASN:ND2	3:D:503:HOH:O	2.34	0.61
1:B:49:LYS:HB2	2:B:401:PEG:H32	1.89	0.55
1:D:50:LEU:HG	1:D:68:LEU:HD13	1.89	0.55
1:B:116:VAL:HA	1:B:124:VAL:O	2.08	0.54
1:D:97:VAL:O	1:D:262:ASN:HA	2.10	0.51
1:B:49:LYS:HB2	2:B:401:PEG:H22	1.94	0.48
1:D:219:VAL:HG22	1:D:281:GLN:HB3	1.94	0.48
1:D:70:CRO:N2	1:D:299:GLU:OE2	2.46	0.48
1:B:215:MET:HE1	3:B:521:HOH:O	2.14	0.48
1:B:241:HIS:HE1	1:B:257:ASP:OD1	1.97	0.47
1:B:210:GLY:HA2	1:D:77[A]:ARG:NH2	2.31	0.46
1:D:51:ILE:HD13	1:D:292:ARG:CZ	2.46	0.46
1:D:243:GLU:CG	1:D:255:LEU:HD21	2.46	0.46
1:D:124:VAL:HG12	1:D:126[B]:ARG:HG3	1.99	0.45
1:D:241:HIS:HE1	1:D:257:ASP:OD1	1.99	0.45
1:B:49:LYS:CB	2:B:401:PEG:H22	2.47	0.45
1:B:292[A]:ARG:HE	1:B:292[A]:ARG:HB2	1.55	0.44
1:B:101:THR:HG21	1:B:103:PHE:CZ	2.53	0.43
1:D:135:ARG:NH1	1:D:137:ASP:OD1	2.51	0.43
1:B:24:GLY:HA2	1:B:129:LEU:O	2.18	0.43
1:B:241:HIS:CE1	1:B:257:ASP:OD1	2.72	0.43
1:D:46:LEU:C	1:D:46:LEU:HD22	2.44	0.43
1:B:10:GLU:HA	1:B:13:THR:HG23	2.00	0.42
1:B:219:VAL:HA	1:B:222:ILE:HD12	2.01	0.42
1:B:220:ARG:HA	3:B:547:HOH:O	2.19	0.42
1:B:208:ASP:O	1:B:212:PHE:HB3	2.20	0.42
1:D:209:PHE:O	1:D:210:GLY:C	2.62	0.42
1:D:48:LEU:HD21	1:D:72:MET:HE1	2.01	0.41
1:B:61:TRP:CE3	1:B:295:MET:HE2	2.56	0.41
1:D:46:LEU:HD22	1:D:46:LEU:O	2.21	0.41
1:D:49:LYS:HB2	2:D:401:PEG:H32	2.02	0.41
1:D:18:ILE:HG21	1:D:48:LEU:HD21	2.03	0.41
1:D:18:ILE:HG21	1:D:48:LEU:CD2	2.51	0.40
1:B:39:GLY:HA3	1:B:75:PHE:CD1	2.56	0.40
1:B:138:GLY:O	1:B:142:GLY:HA3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	240/314 (76%)	235 (98%)	5 (2%)	0	100	100
1	D	241/314 (77%)	233 (97%)	7 (3%)	1 (0%)	30	21
All	All	481/628 (77%)	468 (97%)	12 (2%)	1 (0%)	43	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	151	LEU

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	207/271 (76%)	200 (97%)	7 (3%)	32	23
1	D	205/271 (76%)	199 (97%)	6 (3%)	37	29
All	All	412/542 (76%)	399 (97%)	13 (3%)	34	25

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	117	LYS
1	B	216	VAL
1	B	219	VAL
1	B	227	VAL
1	B	263	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	270	VAL
1	B	272	LEU
1	D	46	LEU
1	D	56	GLU
1	D	149	PHE
1	D	219	VAL
1	D	272	LEU
1	D	284	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	81	HIS
1	B	236	ASN
1	B	241	HIS
1	B	280	HIS
1	D	29	HIS
1	D	125	ASN
1	D	241	HIS
1	D	275	ASN
1	D	280	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CRO	D	70	1	20,20,24	1.01	1 (5%)	25,27,34	1.83	7 (28%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CRO	B	70	1	20,20,24	0.98	1 (5%)	25,27,34	1.57	4 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CRO	D	70	1	-	1/6/25/32	0/2/2/2
1	CRO	B	70	1	-	2/6/25/32	0/2/2/2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	70	CRO	CA2-C2	-2.63	1.45	1.48
1	D	70	CRO	CA2-C2	-2.52	1.45	1.48

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	70	CRO	CG2-CB2-CA2	5.04	135.86	129.87
1	B	70	CRO	C1-CA1-N1	-4.33	103.28	112.85
1	B	70	CRO	O2-C2-CA2	-4.18	128.35	131.02
1	D	70	CRO	CB2-CA2-N2	3.40	133.37	128.76
1	D	70	CRO	CB2-CA2-C2	-2.87	118.88	122.36
1	D	70	CRO	C1-CA1-N1	-2.86	106.51	112.85
1	D	70	CRO	CA2-C2-N3	2.50	105.60	103.50
1	D	70	CRO	CA1-C1-N3	2.31	125.61	122.52
1	B	70	CRO	O3-C3-CA3	-2.27	115.25	125.47
1	D	70	CRO	O3-C3-CA3	-2.09	116.06	125.47
1	B	70	CRO	CB2-CA2-N2	2.06	131.56	128.76

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	70	CRO	C2-CA2-CB2-CG2
1	B	70	CRO	N2-CA2-CB2-CG2
1	D	70	CRO	C3-CA3-N3-C2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	D	70	CRO	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PEG	D	401	-	6,6,6	0.32	0	5,5,5	0.30	0
2	PEG	B	401	-	6,6,6	0.64	0	5,5,5	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PEG	D	401	-	-	3/4/4/4	-
2	PEG	B	401	-	-	1/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	401	PEG	O1-C1-C2-O2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	D	401	PEG	O2-C3-C4-O4
2	D	401	PEG	C4-C3-O2-C2
2	B	401	PEG	O2-C3-C4-O4

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	401	PEG	2	0
2	B	401	PEG	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	B	244/314 (77%)	-1.22	0 100 100	21, 32, 51, 100	2 (0%)
1	D	244/314 (77%)	-1.22	0 100 100	16, 32, 50, 74	3 (1%)
All	All	488/628 (77%)	-1.22	0 100 100	16, 32, 51, 100	5 (1%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CRO	B	70	19/23	0.99	0.05	23,29,32,34	0
1	CRO	D	70	19/23	0.99	0.05	20,24,29,30	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PEG	B	401	7/7	0.98	0.06	33,41,49,50	0
2	PEG	D	401	7/7	0.98	0.05	36,39,41,43	0

6.5 Other polymers [i](#)

There are no such residues in this entry.