



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 12:56 PM UTC

PDB ID : 4P7O / pdb_00004p7o
Title : Structure of Escherichia coli PgaB C-terminal domain, P1 crystal form
Authors : Little, D.J.; Li, G.; Ing, C.; DiFrancesco, B.; Bamford, N.C.; Robinson, H.; Nitz, M.; Pomes, R.; Howell, P.L.
Deposited on : 2014-03-27
Resolution : 1.48 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6779 (1.50-1.46)
Clashscore	190562	7025 (1.50-1.46)
Ramachandran outliers	187476	6917 (1.50-1.46)
Sidechain outliers	187428	6914 (1.50-1.46)
RSRZ outliers	180081	6781 (1.50-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	367	
1	B	367	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6424 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Poly-beta-1,6-N-acetyl-D-glucosamine N-deacetylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	345	Total	C	N	O	S	0	0	0
			2801	1799	477	516	9			
1	B	346	Total	C	N	O	S	0	0	0
			2809	1805	477	518	9			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	306	GLY	-	expression tag	UNP P75906
A	307	SER	-	expression tag	UNP P75906
A	308	HIS	-	expression tag	UNP P75906
A	309	MET	-	expression tag	UNP P75906
B	306	GLY	-	expression tag	UNP P75906
B	307	SER	-	expression tag	UNP P75906
B	308	HIS	-	expression tag	UNP P75906
B	309	MET	-	expression tag	UNP P75906


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	402	Total	O	0	0
			402	402		
2	B	412	Total	O	0	0
			412	412		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Poly-beta-1,6-N-acetyl-D-glucosamine N-deacetylase



- Molecule 1: Poly-beta-1,6-N-acetyl-D-glucosamine N-deacetylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	41.62Å 54.02Å 86.46Å 101.70° 98.30° 90.19°	Depositor
Resolution (Å)	41.86 – 1.48 41.86 – 1.48	Depositor EDS
% Data completeness (in resolution range)	92.1 (41.86-1.48) 85.6 (41.86-1.48)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.85 (at 1.48Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1615)	Depositor
R, R_{free}	0.221 , 0.253 0.222 , 0.253	Depositor DCC
R_{free} test set	2000 reflections (1.73%)	wwPDB-VP
Wilson B-factor (Å ²)	14.2	Xtriage
Anisotropy	0.653	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 52.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6424	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 19.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/2878	0.74	0/3920
1	B	0.41	0/2885	0.71	2/3927 (0.1%)
All	All	0.41	0/5763	0.72	2/7847 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	312	SER	CA-C-N	-5.25	114.83	120.03
1	B	312	SER	C-N-CA	-5.25	114.83	120.03

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2801	0	2691	19	0
1	B	2809	0	2708	22	0
2	A	402	0	0	9	0
2	B	412	0	0	3	0
All	All	6424	0	5399	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:631:MET:HE1	1:B:659:ILE:HG23	1.66	0.77
1:A:466:ASP:OD2	2:A:949:HOH:O	2.03	0.75
1:B:628:ALA:HA	1:B:631:MET:HE3	1.71	0.71
1:B:627:LEU:HG	1:B:631:MET:HE2	1.78	0.66
1:A:611:GLN:OE1	2:A:897:HOH:O	2.13	0.65
1:A:605:ILE:HD13	1:A:641:ASN:HB3	1.78	0.65
1:A:486:GLN:NE2	2:A:986:HOH:O	2.29	0.64
1:A:390:ARG:NH1	2:A:1010:HOH:O	2.36	0.58
1:A:556:ASN:ND2	2:A:1009:HOH:O	2.38	0.56
1:B:502:GLN:NE2	2:B:940:HOH:O	2.39	0.52
1:B:631:MET:HE1	1:B:659:ILE:HD12	1.92	0.52
1:B:390:ARG:HH11	1:B:460:ASP:CG	2.19	0.50
1:B:640:LYS:NZ	2:B:705:HOH:O	2.43	0.50
1:B:605:ILE:HD13	1:B:641:ASN:HB3	1.92	0.50
1:A:358:ASP:OD2	2:A:952:HOH:O	2.19	0.50
1:B:336:ILE:O	1:B:340:ILE:HG12	2.12	0.50
1:B:561:LEU:HD22	1:B:603:LYS:HG3	1.94	0.49
1:A:356:PHE:CD1	1:A:402:PRO:HD3	2.47	0.49
1:A:319:ILE:HD13	1:A:339:LEU:HD11	1.97	0.47
1:B:356:PHE:CD1	1:B:402:PRO:HD3	2.50	0.47
1:A:365:VAL:HG21	1:A:402:PRO:HG2	1.98	0.46
1:B:561:LEU:HB3	1:B:603:LYS:HE2	1.97	0.46
1:A:321:LEU:O	1:A:324:VAL:HG22	2.16	0.46
1:A:536:ALA:HA	1:A:566:TRP:O	2.17	0.45
1:B:648:ASN:HB3	1:B:653:GLN:HB2	1.99	0.45
1:B:611:GLN:NE2	1:B:619:HIS:O	2.49	0.44
1:B:390:ARG:HH12	1:B:459:PHE:HA	1.82	0.44
1:B:319:ILE:HD13	1:B:339:LEU:HD11	1.99	0.43
1:A:387:TRP:CZ2	1:A:391:THR:HG21	2.54	0.43
1:B:422:GLU:H	1:B:422:GLU:CD	2.27	0.43
1:A:577:GLY:O	1:B:423:LYS:HD2	2.18	0.42
1:B:512:ARG:NH1	2:B:993:HOH:O	2.51	0.42
1:A:565:ASP:OD1	2:A:701:HOH:O	2.22	0.42
1:A:334:ARG:NH2	2:A:1055:HOH:O	2.53	0.42
1:A:336:ILE:O	1:A:340:ILE:HG12	2.20	0.41
1:B:330:GLN:HG2	1:B:334:ARG:NH2	2.34	0.41
1:B:539:ILE:HB	1:B:560:PHE:CE2	2.55	0.41
1:A:516:ASP:HB2	2:A:1035:HOH:O	2.21	0.41
1:A:351:VAL:HB	1:A:397:ILE:HD13	2.03	0.40
1:B:574:TYR:CE1	1:B:580:GLU:HA	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	339/367 (92%)	336 (99%)	3 (1%)	0	100	100
1	B	340/367 (93%)	336 (99%)	4 (1%)	0	100	100
All	All	679/734 (92%)	672 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	294/318 (92%)	293 (100%)	1 (0%)	86	75
1	B	295/318 (93%)	294 (100%)	1 (0%)	86	75
All	All	589/636 (93%)	587 (100%)	2 (0%)	86	75

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	546	GLN
1	B	620	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	347	GLN
1	A	611	GLN
1	B	538	ASN
1	B	620	GLN
1	B	626	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	345/367 (94%)	0.80	26 (7%) 20 22	12, 19, 32, 52	0
1	B	346/367 (94%)	0.91	37 (10%) 11 12	13, 19, 36, 57	0
All	All	691/734 (94%)	0.86	63 (9%) 15 16	12, 19, 35, 57	0

All (63) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	668	TYR	5.5
1	B	598	PRO	4.9
1	A	613	TRP	4.6
1	B	621	ALA	4.3
1	A	619	HIS	4.2
1	A	669	PRO	4.0
1	B	597	ILE	3.8
1	B	669	PRO	3.8
1	A	668	TYR	3.7
1	A	621	ALA	3.6
1	B	620	GLN	3.6
1	B	613	TRP	3.6
1	B	599	GLN	3.5
1	B	636	LEU	3.4
1	B	596	ASN	3.4
1	A	636	LEU	3.3
1	A	566	TRP	3.3
1	B	622	ILE	3.2
1	A	433	HIS	3.2
1	A	387	TRP	3.2
1	A	364	LEU	3.1
1	A	324	VAL	3.1
1	A	597	ILE	3.0
1	B	324	VAL	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	665	THR	2.9
1	B	619	HIS	2.9
1	B	394	GLY	2.9
1	B	312	SER	2.9
1	B	653	GLN	2.9
1	A	323	TYR	2.8
1	A	491	GLY	2.7
1	B	390	ARG	2.7
1	A	432	TYR	2.7
1	B	362	ASP	2.6
1	B	614	GLN	2.6
1	B	493	LEU	2.6
1	A	362	ASP	2.6
1	A	361	GLY	2.5
1	B	649	PHE	2.5
1	B	639	VAL	2.5
1	B	665	THR	2.5
1	A	312	SER	2.4
1	B	487	ALA	2.4
1	A	365	VAL	2.3
1	B	334	ARG	2.3
1	B	323	TYR	2.3
1	B	651	HIS	2.2
1	B	321	LEU	2.2
1	B	486	GLN	2.2
1	A	487	ALA	2.1
1	A	423	LYS	2.1
1	B	387	TRP	2.1
1	A	313	PRO	2.1
1	A	606	LEU	2.1
1	A	532	HIS	2.1
1	B	433	HIS	2.1
1	B	667	TRP	2.1
1	B	360	ASP	2.1
1	B	488	GLY	2.0
1	B	359	PRO	2.0
1	A	368	VAL	2.0
1	B	489	PHE	2.0
1	B	432	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.