



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 09:27 PM EDT

PDB ID : 3P8A / pdb_00003p8a
Title : Crystal Structure of a hypothetical protein from Staphylococcus aureus
Authors : Lam, R.; Qiu, W.; Battaile, K.; Lam, K.; Romanov, V.; Chan, T.; Pai, E.; Chirgadze, N.Y.
Deposited on : 2010-10-13
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	274	11% 79% 10% 8%
1	B	274	9% 79% 12% 8%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4605 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called uncharacterized protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	251	Total	C	N	O	S	Se	0	2	0
			2115	1367	349	391	1	7			
1	B	252	Total	C	N	O	S	Se	0	4	0
			2130	1377	354	393	1	5			

There are 40 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	-	initiating methionine	UNP Q8KYA6
A	2	GLY	-	expression tag	UNP Q8KYA6
A	3	SER	-	expression tag	UNP Q8KYA6
A	4	SER	-	expression tag	UNP Q8KYA6
A	5	HIS	-	expression tag	UNP Q8KYA6
A	6	HIS	-	expression tag	UNP Q8KYA6
A	7	HIS	-	expression tag	UNP Q8KYA6
A	8	HIS	-	expression tag	UNP Q8KYA6
A	9	HIS	-	expression tag	UNP Q8KYA6
A	10	HIS	-	expression tag	UNP Q8KYA6
A	11	SER	-	expression tag	UNP Q8KYA6
A	12	SER	-	expression tag	UNP Q8KYA6
A	13	GLY	-	expression tag	UNP Q8KYA6
A	14	LEU	-	expression tag	UNP Q8KYA6
A	15	VAL	-	expression tag	UNP Q8KYA6
A	16	PRO	-	expression tag	UNP Q8KYA6
A	17	ARG	-	expression tag	UNP Q8KYA6
A	18	GLY	-	expression tag	UNP Q8KYA6
A	19	SER	-	expression tag	UNP Q8KYA6
A	20	HIS	-	expression tag	UNP Q8KYA6
B	1	MSE	-	initiating methionine	UNP Q8KYA6
B	2	GLY	-	expression tag	UNP Q8KYA6
B	3	SER	-	expression tag	UNP Q8KYA6
B	4	SER	-	expression tag	UNP Q8KYA6
B	5	HIS	-	expression tag	UNP Q8KYA6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	6	HIS	-	expression tag	UNP Q8KYA6
B	7	HIS	-	expression tag	UNP Q8KYA6
B	8	HIS	-	expression tag	UNP Q8KYA6
B	9	HIS	-	expression tag	UNP Q8KYA6
B	10	HIS	-	expression tag	UNP Q8KYA6
B	11	SER	-	expression tag	UNP Q8KYA6
B	12	SER	-	expression tag	UNP Q8KYA6
B	13	GLY	-	expression tag	UNP Q8KYA6
B	14	LEU	-	expression tag	UNP Q8KYA6
B	15	VAL	-	expression tag	UNP Q8KYA6
B	16	PRO	-	expression tag	UNP Q8KYA6
B	17	ARG	-	expression tag	UNP Q8KYA6
B	18	GLY	-	expression tag	UNP Q8KYA6
B	19	SER	-	expression tag	UNP Q8KYA6
B	20	HIS	-	expression tag	UNP Q8KYA6

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is 2-[BIS-(2-HYDROXY-ETHYL)-AMINO]-2-HYDROXYMETHYL-PROPAN E-1,3-DIOL (CCD ID: BTB) (formula: $C_8H_{19}NO_5$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			13	8	5		

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Cl	0	0
			1	1		
5	B	2	Total	Cl	0	0
			2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	122	Total	O	0	0
			122	122		
6	B	158	Total	O	0	0
			158	158		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: uncharacterized protein



- Molecule 1: uncharacterized protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	148.18Å 49.73Å 67.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.15 – 1.95 47.15 – 1.95	Depositor EDS
% Data completeness (in resolution range)	(Not available) (47.15-1.95) 98.0 (47.15-1.95)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.81 (at 1.95Å)	Xtriage
Refinement program	BUSTER 2.8.0	Depositor
R, R_{free}	0.238 , 0.270 0.234 , 0.270	Depositor DCC
R_{free} test set	1837 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	16.4	Xtriage
Anisotropy	0.036	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 55.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4605	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 50.05 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.8346e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CL, PG4, BTB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.80	1/2165 (0.0%)	1.20	5/2905 (0.2%)
1	B	0.80	0/2187	1.18	9/2937 (0.3%)
All	All	0.80	1/4352 (0.0%)	1.19	14/5842 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	88	ILE	CG1-CD1	6.73	1.77	1.51

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	129	VAL	N-CA-CB	7.58	119.64	111.00
1	A	232	ASP	CA-CB-CG	6.59	119.19	112.60
1	B	232	ASP	CA-CB-CG	6.33	118.92	112.60
1	B	245	TYR	N-CA-C	-5.98	100.54	109.15
1	A	158	ASP	CA-CB-CG	5.51	118.11	112.60
1	A	245	TYR	N-CA-C	-5.42	101.34	109.15
1	A	74	ASP	CA-CB-CG	5.26	117.86	112.60
1	B	74	ASP	CA-CB-CG	5.22	117.82	112.60
1	B	272	PRO	CA-C-N	5.15	130.97	121.70
1	B	272	PRO	C-N-CA	5.15	130.97	121.70
1	B	244	LYS	N-CA-C	5.08	116.25	108.42
1	B	256	THR	N-CA-C	-5.07	107.64	113.88
1	B	241	LYS	N-CA-C	-5.06	107.06	113.18
1	B	144	ASP	CA-CB-CG	5.02	117.62	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2115	0	2093	20	0
1	B	2130	0	2113	11	0
2	A	12	0	15	1	0
2	B	24	0	32	3	0
3	A	14	0	19	4	0
3	B	14	0	19	0	0
4	A	13	0	18	4	0
5	A	1	0	0	0	0
5	B	2	0	0	0	0
6	A	122	0	0	0	0
6	B	158	0	0	2	0
All	All	4605	0	4309	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:88:ILE:CD1	1:A:88:ILE:CG1	1.78	1.60
4:A:277:PG4:H51	4:A:277:PG4:H81	1.78	0.64
1:A:105:GLN:H	3:A:276:BTB:H12	1.63	0.63
1:A:23:LEU:HD23	1:A:114:THR:HG23	1.84	0.60
1:A:104:GLU:HA	3:A:276:BTB:H41	1.84	0.58
1:A:202:SER:OG	4:A:277:PG4:H62	2.04	0.56
1:A:60:ASN:ND2	2:A:275:GOL:H11	2.22	0.55
1:B:121:LYS:HD2	1:B:134:PRO:HD3	1.89	0.55
1:A:121:LYS:HD2	1:A:134:PRO:HD3	1.88	0.54
1:A:88:ILE:HD12	1:A:88:ILE:H	1.76	0.51
1:B:141:THR:HG23	1:B:143:LYS:H	1.74	0.51
4:A:277:PG4:H81	4:A:277:PG4:H32	1.95	0.49
1:A:173:SER:H	1:A:176:MSE:HE2	1.76	0.49
1:B:139:ARG:NH2	2:B:278:GOL:O1	2.39	0.48
1:A:121:LYS:HG3	1:A:131:VAL:HG21	1.96	0.47
1:A:172:GLU:HG3	1:A:176:MSE:HE3	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:51:GLY:H	1:B:60:ASN:ND2	2.13	0.47
1:A:201:ARG:CZ	4:A:277:PG4:H71	2.46	0.46
1:A:114:THR:HG21	1:A:120:VAL:CG2	2.46	0.45
3:A:276:BTB:H62	3:A:276:BTB:H82	1.98	0.45
1:A:114:THR:HG21	1:A:120:VAL:HG21	1.98	0.45
1:A:175:SER:HB3	1:A:178:THR:HB	1.99	0.45
1:A:174:ASP:HA	1:A:175:SER:HA	1.68	0.44
1:B:84:MSE:O	1:B:90:GLY:HA3	2.18	0.44
1:A:105:GLN:H	3:A:276:BTB:C1	2.29	0.44
2:B:281:GOL:H2	6:B:404:HOH:O	2.18	0.44
1:B:244:LYS:NZ	6:B:339:HOH:O	2.46	0.43
1:A:88:ILE:CD1	1:A:88:ILE:H	2.32	0.42
1:B:201:ARG:CZ	2:B:281:GOL:H11	2.49	0.42
1:B:177:ARG:CZ	1:B:181:LEU:HD11	2.50	0.42
1:A:117:ILE:HB	1:A:169:GLN:HB2	2.02	0.42
1:B:50:GLY:HA3	1:B:60:ASN:ND2	2.35	0.41
1:B:117:ILE:HB	1:B:169:GLN:HB2	2.03	0.41
1:B:93:ALA:HA	1:B:153:TYR:CZ	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	249/274 (91%)	241 (97%)	8 (3%)	0	100	100
1	B	254/274 (93%)	247 (97%)	7 (3%)	0	100	100
All	All	503/548 (92%)	488 (97%)	15 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	235/245 (96%)	221 (94%)	14 (6%)	17	7
1	B	236/245 (96%)	222 (94%)	14 (6%)	18	7
All	All	471/490 (96%)	443 (94%)	28 (6%)	17	7

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	ILE
1	A	44	VAL
1	A	68	ASN
1	A	75	VAL
1	A	79	GLU
1	A	88	ILE
1	A	114	THR
1	A	129	VAL
1	A	157	GLN
1	A	158	ASP
1	A	171	GLU
1	A	175	SER
1	A	213	ASP
1	A	232	ASP
1	B	22	ILE
1	B	44	VAL
1	B	60	ASN
1	B	68	ASN
1	B	75	VAL
1	B	130	GLU
1	B	142	HIS
1	B	143	LYS
1	B	155	MSE
1	B	157	GLN
1	B	179	LYS
1	B	183	LYS
1	B	213	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	232	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	78	ASN
1	A	97	GLN
1	A	125	GLN
1	B	60	ASN
1	B	68	ASN
1	B	78	ASN
1	B	97	GLN
1	B	182	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 3 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	A	275	-	5,5,5	0.86	0	5,5,5	1.29	0
2	GOL	B	276	-	5,5,5	0.69	0	5,5,5	0.49	0
3	BTB	B	275	-	13,13,13	2.02	4 (30%)	7,16,16	1.23	0
2	GOL	A	279	-	5,5,5	1.26	1 (20%)	5,5,5	1.10	0
3	BTB	A	276	-	13,13,13	1.81	3 (23%)	7,16,16	1.32	1 (14%)
4	PG4	A	277	-	12,12,12	1.14	0	11,11,11	1.19	2 (18%)
2	GOL	B	278	-	5,5,5	0.73	0	5,5,5	1.50	1 (20%)
2	GOL	B	280	-	5,5,5	0.94	0	5,5,5	0.93	0
2	GOL	B	281	-	5,5,5	0.63	0	5,5,5	0.94	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	275	-	-	4/4/4/4	-
2	GOL	B	276	-	-	2/4/4/4	-
3	BTB	B	275	-	-	14/21/21/21	-
2	GOL	A	279	-	-	0/4/4/4	-
3	BTB	A	276	-	-	8/21/21/21	-
4	PG4	A	277	-	-	4/10/10/10	-
2	GOL	B	278	-	-	4/4/4/4	-
2	GOL	B	280	-	-	2/4/4/4	-
2	GOL	B	281	-	-	2/4/4/4	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	275	BTB	C2-N	4.49	1.57	1.48
3	A	276	BTB	C4-C2	3.67	1.57	1.53
3	B	275	BTB	C4-C2	3.58	1.57	1.53
3	B	275	BTB	C3-C2	3.14	1.57	1.53
3	A	276	BTB	C7-N	3.07	1.52	1.48
3	A	276	BTB	C2-N	2.90	1.54	1.48
3	B	275	BTB	C1-C2	2.17	1.55	1.53
2	A	279	GOL	O2-C2	2.02	1.49	1.43

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	276	BTB	O1-C1-C2	-2.37	105.83	111.40
4	A	277	PG4	O4-C6-C5	2.27	120.69	110.35
4	A	277	PG4	C7-O4-C6	-2.11	104.01	113.26
2	B	278	GOL	O2-C2-C3	2.02	117.56	109.18

There are no chirality outliers.

All (40) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	275	GOL	O1-C1-C2-C3
2	A	275	GOL	C1-C2-C3-O3
2	B	278	GOL	O1-C1-C2-C3
2	B	278	GOL	C1-C2-C3-O3
2	B	280	GOL	O1-C1-C2-C3
3	A	276	BTB	O1-C1-C2-C3
3	A	276	BTB	C1-C2-C3-O3
3	A	276	BTB	C4-C2-C3-O3
3	A	276	BTB	N-C2-C3-O3
3	A	276	BTB	C8-C7-N-C5
3	B	275	BTB	O1-C1-C2-C4
3	B	275	BTB	O1-C1-C2-N
3	B	275	BTB	C4-C2-C3-O3
3	B	275	BTB	N-C2-C3-O3
3	B	275	BTB	C1-C2-C4-O4
3	B	275	BTB	C3-C2-C4-O4
3	B	275	BTB	N-C2-C4-O4
3	B	275	BTB	C1-C2-N-C5
3	B	275	BTB	C3-C2-N-C5
3	B	275	BTB	C3-C2-N-C7
3	B	275	BTB	C4-C2-N-C7
4	A	277	PG4	C4-C3-O2-C2
2	A	275	GOL	O2-C2-C3-O3
4	A	277	PG4	O3-C5-C6-O4
2	B	276	GOL	O1-C1-C2-C3
2	B	281	GOL	C1-C2-C3-O3
2	A	275	GOL	O1-C1-C2-O2
2	B	276	GOL	O1-C1-C2-O2
2	B	278	GOL	O1-C1-C2-O2
2	B	278	GOL	O2-C2-C3-O3
3	B	275	BTB	C8-C7-N-C2
2	B	280	GOL	O1-C1-C2-O2
3	A	276	BTB	N-C7-C8-O8
3	A	276	BTB	C3-C2-N-C7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	B	275	BTB	C1-C2-N-C7
3	B	275	BTB	C4-C2-N-C5
2	B	281	GOL	O2-C2-C3-O3
4	A	277	PG4	C6-C5-O3-C4
3	A	276	BTB	O1-C1-C2-N
4	A	277	PG4	O2-C3-C4-O3

There are no ring outliers.

5 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	275	GOL	1	0
3	A	276	BTB	4	0
4	A	277	PG4	4	0
2	B	278	GOL	1	0
2	B	281	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	245/274 (89%)	0.74	29 (11%) 9 10	8, 20, 47, 67	1 (0%)
1	B	247/274 (90%)	0.55	25 (10%) 12 14	4, 18, 44, 65	4 (1%)
All	All	492/548 (89%)	0.64	54 (10%) 10 12	4, 19, 47, 67	5 (1%)

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	273	LEU	7.9
1	A	273	LEU	6.2
1	B	142	HIS	5.1
1	A	174	ASP	4.7
1	A	179	LYS	4.4
1	B	178	THR	4.1
1	B	157	GLN	4.0
1	A	173	SER	3.8
1	A	180	LYS	3.4
1	B	179	LYS	3.3
1	A	181	LEU	3.3
1	B	181	LEU	3.3
1	B	144	ASP	3.2
1	A	22	ILE	3.2
1	B	174	ASP	3.1
1	B	79	GLU	3.1
1	B	180	LYS	3.1
1	B	141	THR	3.1
1	B	175	SER	3.0
1	B	256	THR	3.0
1	A	175	SER	3.0
1	A	157	GLN	3.0
1	B	143	LYS	2.9
1	A	88	ILE	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	255	ALA	2.8
1	A	142	HIS	2.8
1	A	146	LYS	2.8
1	A	143	LYS	2.7
1	A	256	THR	2.6
1	A	54	HIS	2.5
1	B	138	GLU	2.5
1	A	147	VAL	2.5
1	B	136	GLN	2.5
1	A	141	THR	2.5
1	B	185	PHE	2.4
1	B	173	SER	2.4
1	B	232	ASP	2.4
1	A	65	ILE	2.4
1	A	114	THR	2.3
1	A	86	LYS	2.3
1	A	254	GLN	2.3
1	B	37	ARG	2.3
1	B	145	GLY	2.3
1	A	182	GLN	2.3
1	A	178	THR	2.2
1	B	254	GLN	2.2
1	A	232	ASP	2.2
1	A	83	LYS	2.2
1	B	159	ASP	2.2
1	A	140	ASP	2.1
1	A	177	ARG	2.1
1	A	255	ALA	2.1
1	A	158	ASP	2.1
1	B	115	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	BTB	A	276	14/14	0.63	0.21	32,34,37,39	0
3	BTB	B	275	14/14	0.64	0.22	39,41,44,44	0
4	PG4	A	277	13/13	0.73	0.16	30,31,33,37	0
2	GOL	B	280	6/6	0.76	0.15	34,35,36,38	0
2	GOL	B	281	6/6	0.80	0.17	26,27,27,28	0
2	GOL	A	279	6/6	0.84	0.11	28,29,30,30	0
2	GOL	B	276	6/6	0.85	0.11	27,28,28,29	0
2	GOL	A	275	6/6	0.89	0.12	23,23,24,26	0
2	GOL	B	278	6/6	0.90	0.09	24,24,24,25	0
5	CL	B	279	1/1	0.90	0.11	45,45,45,45	0
5	CL	B	277	1/1	0.97	0.06	23,23,23,23	0
5	CL	A	278	1/1	0.98	0.05	23,23,23,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.