



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 04:07 AM UTC

PDB ID : 8P8X / pdb_00008p8x
Title : Crystal structure of a pathogenic mutant variant of human mitochondrial PheRS
Authors : Chen, W.; Kuhle, B.
Deposited on : 2023-06-03
Resolution : 1.46 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.46 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1756 (1.46-1.46)
Clashscore	190562	1795 (1.46-1.46)
Ramachandran outliers	187476	1776 (1.46-1.46)
Sidechain outliers	187428	1776 (1.46-1.46)
RSRZ outliers	180081	1756 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	415	4% 71% 26% ..

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3851 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Phenylalanine-tRNA ligase, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	404	3362	2155	594	601	12	0	2	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP O95363
A	2	PRO	-	expression tag	UNP O95363
A	162	LEU	ARG	conflict	UNP O95363

- Molecule 2 is PHENYLALANINE (CCD ID: PHE) (formula: $C_9H_{11}NO_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	12	9	1	2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	477	Total 477	O 477	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phenylalanine-tRNA ligase, mitochondrial



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	54.53Å 88.46Å 97.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.32 – 1.46 36.32 – 1.46	Depositor EDS
% Data completeness (in resolution range)	99.6 (36.32-1.46) 99.6 (36.32-1.46)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.40 (at 1.46Å)	Xtriage
Refinement program	PHENIX (1.18.2_3874: ???)	Depositor
R, R_{free}	0.145 , 0.167 0.158 , 0.171	Depositor DCC
R_{free} test set	2000 reflections (2.42%)	wwPDB-VP
Wilson B-factor (Å ²)	19.1	Xtriage
Anisotropy	0.346	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 40.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3851	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.78	90/3454 (2.6%)	1.26	13/4677 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

All (90) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	82	PRO	C-O	-8.31	1.13	1.24
1	A	350	ARG	C-O	-7.79	1.15	1.24
1	A	233	PRO	C-O	-7.53	1.14	1.24
1	A	96	PRO	C-O	-7.48	1.14	1.23
1	A	153	PRO	C-O	-7.39	1.15	1.24
1	A	159	GLU	C-O	-7.27	1.15	1.23
1	A	188	SER	C-O	-7.25	1.14	1.24
1	A	49	PRO	C-O	-7.23	1.15	1.24
1	A	216	ALA	C-O	-6.96	1.16	1.24
1	A	100	PRO	C-O	-6.82	1.15	1.24
1	A	167	HIS	C-O	-6.67	1.16	1.24
1	A	22	TYR	C-O	-6.65	1.19	1.25
1	A	296	PHE	C-O	-6.52	1.16	1.24
1	A	312	ILE	C-O	-6.50	1.15	1.24
1	A	352	ILE	C-O	-6.43	1.17	1.24
1	A	38	VAL	C-O	-6.40	1.17	1.24
1	A	291	PRO	C-O	-6.36	1.16	1.24
1	A	281	GLU	C-O	-6.19	1.16	1.24
1	A	230	CYS	C-O	-6.16	1.17	1.23
1	A	23	PRO	C-O	-6.10	1.16	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	369	PRO	C-O	-6.06	1.16	1.24
1	A	232	PHE	C-O	-6.00	1.16	1.24
1	A	320	PRO	C-O	-6.00	1.16	1.23
1	A	164	PHE	C-O	-5.99	1.16	1.23
1	A	129	HIS	CE1-NE2	-5.87	1.26	1.32
1	A	84	VAL	C-O	-5.86	1.17	1.23
1	A	245	PHE	C-O	-5.80	1.17	1.23
1	A	183	GLU	C-O	-5.79	1.17	1.23
1	A	228	VAL	C-O	-5.76	1.18	1.24
1	A	201	LYS	C-O	-5.72	1.17	1.24
1	A	121	SER	C-O	-5.69	1.16	1.24
1	A	152	TYR	C-O	-5.67	1.18	1.24
1	A	235	THR	C-O	-5.64	1.17	1.23
1	A	397	ILE	C-O	-5.63	1.17	1.24
1	A	91	ASP	CG-OD2	-5.58	1.14	1.25
1	A	86	THR	C-O	-5.54	1.17	1.24
1	A	198	GLU	C-O	-5.51	1.17	1.24
1	A	229	ASP	C-O	-5.50	1.17	1.23
1	A	405	ALA	C-O	-5.49	1.17	1.24
1	A	328	ILE	C-O	-5.47	1.18	1.24
1	A	166	LYS	C-O	-5.46	1.17	1.24
1	A	200	VAL	C-O	-5.45	1.17	1.24
1	A	88	GLN	C-O	-5.41	1.17	1.24
1	A	106	ASP	C-O	-5.40	1.16	1.24
1	A	301	GLU	C-O	-5.38	1.17	1.24
1	A	204	GLU	C-O	-5.38	1.17	1.24
1	A	248	GLU	C-O	-5.38	1.17	1.23
1	A	236	HIS	CD2-NE2	-5.36	1.31	1.37
1	A	345	PHE	C-O	-5.32	1.17	1.24
1	A	393	GLU	C-O	-5.31	1.17	1.24
1	A	76	VAL	C-O	-5.30	1.18	1.24
1	A	351	THR	C-O	-5.30	1.18	1.24
1	A	138	VAL	C-O	-5.28	1.18	1.24
1	A	210	THR	C-O	-5.27	1.18	1.24
1	A	89	ASN	C-O	-5.27	1.17	1.24
1	A	122	ALA	C-O	-5.27	1.17	1.24
1	A	382	TYR	C-O	-5.27	1.17	1.24
1	A	123	HIS	CE1-NE2	-5.25	1.27	1.32
1	A	323	LYS	C-O	-5.24	1.17	1.24
1	A	145	ASP	C-O	-5.23	1.19	1.24
1	A	60	HIS	CE1-NE2	-5.22	1.27	1.32
1	A	151	HIS	C-O	-5.22	1.18	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	356	LEU	C-O	-5.21	1.17	1.24
1	A	167	HIS	CE1-NE2	-5.21	1.27	1.32
1	A	257	VAL	C-O	-5.20	1.18	1.24
1	A	293	ILE	C-O	-5.20	1.17	1.24
1	A	45	GLN	C-O	-5.20	1.17	1.24
1	A	29	ASN	C-O	-5.19	1.18	1.24
1	A	143	ARG	C-O	-5.17	1.17	1.23
1	A	209	GLN	C-O	-5.16	1.18	1.24
1	A	324	TYR	C-O	-5.16	1.18	1.23
1	A	341	ALA	C-O	-5.16	1.17	1.23
1	A	43	HIS	C-O	-5.15	1.17	1.24
1	A	97	ALA	C-O	-5.14	1.17	1.24
1	A	385	MET	C-O	-5.12	1.17	1.24
1	A	104	LYS	C-O	-5.11	1.17	1.24
1	A	281	GLU	CD-OE2	-5.11	1.15	1.25
1	A	95	ILE	C-O	-5.10	1.18	1.24
1	A	199	ALA	C-O	-5.10	1.18	1.24
1	A	406	VAL	C-O	-5.10	1.18	1.24
1	A	162	LEU	C-O	-5.09	1.18	1.24
1	A	349	VAL	C-O	-5.09	1.18	1.24
1	A	239	PHE	C-O	-5.08	1.17	1.23
1	A	142	TYR	C-O	-5.07	1.18	1.24
1	A	141	VAL	C-O	-5.05	1.18	1.23
1	A	112	ARG	C-O	-5.05	1.17	1.24
1	A	169	LEU	C-O	-5.03	1.18	1.24
1	A	290	ILE	C-O	-5.01	1.17	1.24
1	A	250	LEU	C-O	-5.00	1.18	1.24
1	A	346	TYR	C-O	-5.00	1.18	1.24

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	217	HIS	CB-CA-C	-8.16	92.93	109.99
1	A	316	VAL	N-CA-CB	7.95	119.20	110.53
1	A	219	PHE	CB-CA-C	-7.47	97.00	109.55
1	A	313	ASN	CA-CB-CG	-7.15	105.45	112.60
1	A	355	ASP	CA-CB-CG	6.21	118.81	112.60
1	A	270	ASP	CB-CA-C	5.65	119.31	111.63
1	A	316	VAL	CB-CA-C	5.50	118.31	111.15
1	A	276	PHE	CA-CB-CG	5.46	119.25	113.80
1	A	414	ARG	CA-C-N	5.21	131.09	121.70
1	A	414	ARG	C-N-CA	5.21	131.09	121.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	253	LEU	CA-C-N	5.08	125.69	122.18
1	A	253	LEU	C-N-CA	5.08	125.69	122.18
1	A	355	ASP	CB-CA-C	5.04	119.09	110.01

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	186	SER	Mainchain
1	A	217	HIS	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3362	0	3274	10	0
2	A	12	0	8	0	0
3	A	477	0	0	5	7
All	All	3851	0	3282	10	7

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:402:GLN:NE2	3:A:602:HOH:O	2.32	0.62
1:A:317:LYS:NZ	3:A:601:HOH:O	2.21	0.49
1:A:12:GLY:N	3:A:605:HOH:O	2.45	0.49
1:A:174:LYS:HD2	1:A:174:LYS:C	2.38	0.48
1:A:201:LYS:NZ	3:A:613:HOH:O	2.48	0.47
1:A:68:ARG:HD3	1:A:171:ALA:HB3	2.00	0.43
1:A:213:ARG:NH2	3:A:623:HOH:O	2.50	0.41
1:A:108:TYR:CZ	1:A:144:ARG:HG2	2.56	0.41
1:A:112:ARG:HH11	1:A:112:ARG:HD2	1.73	0.41
1:A:65:TYR:CD1	1:A:65:TYR:C	3.00	0.40

All (7) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:843:HOH:O	3:A:1000:HOH:O[3_545]	1.93	0.27
3:A:639:HOH:O	3:A:978:HOH:O[4_555]	1.95	0.25
3:A:836:HOH:O	3:A:911:HOH:O[3_545]	1.96	0.24
3:A:997:HOH:O	3:A:1050:HOH:O[3_555]	1.99	0.21
3:A:711:HOH:O	3:A:1023:HOH:O[3_545]	2.14	0.06
3:A:763:HOH:O	3:A:843:HOH:O[3_555]	2.16	0.04
3:A:869:HOH:O	3:A:1003:HOH:O[3_545]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	404/415 (97%)	398 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	365/370 (99%)	362 (99%)	3 (1%)	73	50

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	36	THR
1	A	310	SER
1	A	316	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	180	GLN
1	A	313	ASN
1	A	314	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PHE	A	501	-	11,12,12	1.49	4 (36%)	11,15,15	0.71	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PHE	A	501	-	-	0/8/8/8	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	PHE	OXT-C	-2.83	1.21	1.30
2	A	501	PHE	CE2-CD2	2.22	1.42	1.38
2	A	501	PHE	CD1-CG	2.08	1.43	1.38
2	A	501	PHE	O-C	2.03	1.28	1.22

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	404/415 (97%)	0.03	18 (4%)	38 41	9, 22, 44, 64	2 (0%)

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	415	PHE	5.9
1	A	316	VAL	5.6
1	A	12	GLY	4.7
1	A	66	VAL	4.0
1	A	314	GLN	3.4
1	A	60	HIS	2.8
1	A	315	LYS	2.7
1	A	217	HIS	2.7
1	A	310	SER	2.7
1	A	69	PHE	2.4
1	A	312	ILE	2.3
1	A	311	ASN	2.2
1	A	221	ASP	2.2
1	A	269	GLN	2.2
1	A	317	LYS	2.2
1	A	220	GLY	2.2
1	A	68	ARG	2.2
1	A	62	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PHE	A	501	12/12	0.98	0.04	13,14,15,19	0

6.5 Other polymers [i](#)

There are no such residues in this entry.