



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 12:35 PM UTC

PDB ID : 3P98 / pdb_00003p98
Title : The crystal structure of the extended spectrum beta-lactamase TEM-72 reveals inhibition by citrate
Authors : Docquier, J.D.; Benvenuti, M.; Calderone, V.; Rossolini, G.M.; Mangani, S.
Deposited on : 2010-10-16
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	286	 68% 20% • 8%
1	B	286	 68% 21% • 8%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CIT	B	1	-	X	-	-

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 4341 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-lactamase TEM-72.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	263	Total	C	N	O	S	0	0	0
			2028	1267	361	390	10			
1	B	263	Total	C	N	O	S	0	0	0
			2028	1267	361	390	10			

- Molecule 2 is CITRIC ACID (CCD ID: CIT) (formula: $C_6H_8O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			13	6	7		
2	B	1	Total	C	O	0	0
			13	6	7		

- Molecule 3 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	4	3		
3	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	123	Total	O	0	0
			123	123		
4	B	122	Total	O	0	0
			122	122		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Beta-lactamase TEM-72

Chain A: 



• Molecule 1: Beta-lactamase TEM-72

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	60.63Å 90.21Å 96.05Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.91 – 2.10 28.91 – 2.10	Depositor EDS
% Data completeness (in resolution range)	100.0 (28.91-2.10) 98.4 (28.91-2.10)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.81 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.201 , 0.283 (Not available) , 0.270	Depositor DCC
R_{free} test set	2794 reflections (9.02%)	wwPDB-VP
Wilson B-factor (Å ²)	19.1	Xtriage
Anisotropy	0.171	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 41.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4341	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 39.48 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.1534e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, CIT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.26	4/2062 (0.2%)	1.22	8/2791 (0.3%)
1	B	1.26	2/2062 (0.1%)	1.25	10/2791 (0.4%)
All	All	1.26	6/4124 (0.1%)	1.23	18/5582 (0.3%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	108	VAL	CA-CB	-6.48	1.48	1.55
1	A	119	VAL	CA-CB	5.90	1.61	1.54
1	A	47	ILE	C-O	-5.90	1.18	1.24
1	A	126	ALA	CA-CB	-5.65	1.44	1.53
1	B	119	VAL	CA-CB	5.53	1.61	1.54
1	A	278	ILE	CA-C	5.00	1.59	1.52

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	182	THR	CA-C-N	-7.14	111.18	119.32
1	B	182	THR	C-N-CA	-7.14	111.18	119.32
1	B	225	LEU	CA-C-N	-6.94	113.56	120.98
1	B	225	LEU	C-N-CA	-6.94	113.56	120.98
1	A	115	ASP	N-CA-CB	-6.49	100.49	110.42
1	A	70	SER	N-CA-C	6.38	120.52	112.87
1	A	218	GLY	CA-C-N	-6.25	114.45	120.83
1	A	218	GLY	C-N-CA	-6.25	114.45	120.83
1	B	106	SER	CA-C-N	6.01	125.89	119.28
1	B	106	SER	C-N-CA	6.01	125.89	119.28
1	B	26	HIS	CA-C-N	5.71	126.98	119.84
1	B	26	HIS	C-N-CA	5.71	126.98	119.84
1	A	127	ILE	N-CA-C	5.57	115.66	110.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	56	ILE	N-CA-C	5.55	116.17	108.84
1	A	98	SER	CA-C-N	5.22	127.59	120.54
1	A	98	SER	C-N-CA	5.22	127.59	120.54
1	A	35	ASP	CB-CA-C	-5.12	101.98	110.68
1	B	73	LYS	N-CA-C	5.10	117.74	111.82

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2028	0	2048	43	0
1	B	2028	0	2046	38	0
2	A	13	0	5	3	0
2	B	13	0	5	2	0
3	A	7	0	10	2	0
3	B	7	0	10	1	0
4	A	123	0	0	6	0
4	B	122	0	0	10	0
All	All	4341	0	4124	85	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:201:LEU:HB2	4:B:394:HOH:O	1.40	1.18
1:B:241:ARG:O	1:B:266:GLY:HA2	1.71	0.89
2:B:1:CIT:H41	2:B:1:CIT:O1	1.75	0.85
1:A:69:MET:HG2	1:A:238:SER:OG	1.77	0.84
1:A:216:VAL:HG22	4:A:294:HOH:O	1.78	0.83
1:B:171:GLU:HA	1:B:240:LYS:HD3	1.61	0.82
1:B:106:SER:O	1:B:110:GLU:HG2	1.80	0.81

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:270:THR:OG1	1:B:273:GLU:HG3	1.80	0.80
1:A:77:CYS:HG	1:A:123:CYS:HG	1.26	0.79
1:A:268:GLN:HE21	1:A:268:GLN:HA	1.52	0.75
2:B:1:CIT:O1	2:B:1:CIT:C4	2.38	0.71
1:A:194:LEU:HD22	1:A:208:ILE:HG22	1.78	0.65
1:A:69:MET:HE1	1:A:244:ARG:C	2.21	0.65
1:A:170:ASN:HB3	1:A:238:SER:HB3	1.78	0.65
1:B:26:HIS:N	1:B:58:GLU:OE1	2.30	0.64
1:A:168:GLU:O	1:A:171:GLU:HG3	1.98	0.63
1:A:35:ASP:O	1:A:39:LYS:HG2	1.99	0.62
1:B:69:MET:HG2	1:B:238:SER:HB3	1.82	0.59
1:A:237:ALA:N	2:A:1:CIT:O2	2.30	0.58
1:B:47:ILE:HG21	1:B:187:ALA:HB2	1.86	0.58
1:A:164:ARG:NH2	1:A:179:ASP:OD2	2.36	0.56
1:A:170:ASN:O	1:A:238:SER:HB2	2.06	0.56
1:A:53:SER:OG	1:A:55:LYS:HD3	2.07	0.55
1:A:98:SER:HB2	1:A:100:ASN:ND2	2.22	0.55
1:B:47:ILE:HD12	1:B:59:SER:HB3	1.88	0.55
1:A:112:HIS:HD2	1:A:115:ASP:OD2	1.90	0.54
1:A:225:LEU:HD23	3:A:290:PEG:H31	1.90	0.54
1:A:161:ARG:HD2	1:A:163:ASP:OD2	2.07	0.54
1:B:101:ASP:O	4:B:354:HOH:O	2.18	0.54
1:A:167:PRO:HD2	1:A:168:GLU:OE2	2.08	0.52
1:B:99:GLN:HG2	4:B:328:HOH:O	2.08	0.52
1:B:265:THR:HG23	1:B:266:GLY:N	2.26	0.50
1:A:32:LYS:HE3	1:A:280:GLU:HB3	1.93	0.49
1:A:276:ARG:O	1:A:280:GLU:HG3	2.12	0.49
1:A:69:MET:HE1	1:A:245:GLY:N	2.28	0.48
1:A:69:MET:CG	1:A:238:SER:OG	2.54	0.48
1:A:128:THR:HA	1:A:214:ASP:HA	1.95	0.48
1:A:237:ALA:H	2:A:1:CIT:C1	2.23	0.48
1:B:182:THR:O	1:B:183:PRO:C	2.53	0.48
1:A:98:SER:HB2	1:A:100:ASN:HD22	1.79	0.47
1:B:90:GLN:HG3	4:B:294:HOH:O	2.15	0.47
1:B:54:GLY:HA2	1:B:191:ARG:HD2	1.97	0.47
1:B:169:LEU:HD12	1:B:169:LEU:HA	1.15	0.47
1:B:197:GLU:N	1:B:197:GLU:OE2	2.48	0.47
1:B:258:ARG:NH2	1:B:289:TRP:O	2.45	0.47
1:B:126:ALA:O	1:B:130:SER:HA	2.15	0.47
1:A:83:ARG:NH1	4:A:352:HOH:O	2.37	0.46
1:A:118:THR:OG1	1:A:121:GLU:HG3	2.16	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:126:ALA:O	1:A:130:SER:HA	2.16	0.45
1:A:204:ARG:NH2	4:A:304:HOH:O	2.49	0.45
1:A:205:GLN:NE2	1:A:209:ASP:OD1	2.50	0.44
3:B:290:PEG:H21	4:B:382:HOH:O	2.17	0.44
1:B:26:HIS:HA	1:B:27:PRO:HD2	1.76	0.43
1:B:47:ILE:CG2	1:B:187:ALA:HB2	2.47	0.43
1:B:175:ASN:ND2	4:B:392:HOH:O	2.45	0.43
1:A:174:PRO:O	1:A:175:ASN:HB2	2.18	0.43
1:A:216:VAL:CG2	4:A:294:HOH:O	2.52	0.43
1:B:153:HIS:HD2	4:B:350:HOH:O	2.00	0.43
1:B:191:ARG:HD3	4:B:395:HOH:O	2.19	0.43
3:A:290:PEG:H11	4:A:387:HOH:O	2.17	0.43
1:B:90:GLN:N	1:B:90:GLN:CD	2.78	0.42
1:B:172:ALA:O	1:B:241:ARG:HD2	2.20	0.42
1:A:44:VAL:HG22	1:A:264:THR:HG22	2.02	0.42
1:A:225:LEU:HG	1:A:229:TRP:HB2	2.02	0.42
1:A:32:LYS:HA	1:A:32:LYS:HD2	1.67	0.42
1:B:161:ARG:O	1:B:179:ASP:HA	2.20	0.42
1:B:166:GLU:OE2	4:B:369:HOH:O	2.22	0.42
1:B:193:LEU:HD23	1:B:198:LEU:HD23	2.02	0.42
1:B:269:ALA:HB1	1:B:273:GLU:HB2	2.01	0.42
1:B:69:MET:O	1:B:72:PHE:HD2	2.02	0.42
1:B:93:ARG:O	1:B:118:THR:HA	2.19	0.42
1:B:282:GLY:O	1:B:286:ILE:HG12	2.19	0.41
1:A:43:ARG:HD2	1:A:64:GLU:OE1	2.20	0.41
1:B:27:PRO:O	1:B:31:VAL:HG23	2.20	0.41
1:A:120:ARG:HD3	4:A:347:HOH:O	2.20	0.41
1:A:237:ALA:O	2:A:1:CIT:C1	2.69	0.41
1:A:268:GLN:HE21	1:A:268:GLN:CA	2.29	0.40
1:B:67:PRO:HD2	4:B:398:HOH:O	2.20	0.40
1:B:252:PRO:HG2	1:B:255:LYS:HB2	2.03	0.40
1:B:229:TRP:CD1	1:B:252:PRO:HA	2.56	0.40
1:A:81:LEU:HD23	1:A:81:LEU:HA	1.86	0.40
1:A:42:ALA:HB1	1:A:265:THR:O	2.21	0.40
1:A:57:LEU:CD1	1:A:258:ARG:HD3	2.51	0.40
1:A:58:GLU:OE2	1:A:58:GLU:HA	2.20	0.40
1:B:152:LEU:HA	1:B:155:MET:HG2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	261/286 (91%)	253 (97%)	8 (3%)	0	100	100
1	B	261/286 (91%)	244 (94%)	17 (6%)	0	100	100
All	All	522/572 (91%)	497 (95%)	25 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	218/237 (92%)	201 (92%)	17 (8%)	11	9
1	B	218/237 (92%)	203 (93%)	15 (7%)	14	12
All	All	436/474 (92%)	404 (93%)	32 (7%)	13	10

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	32	LYS
1	A	55	LYS
1	A	58	GLU
1	A	98	SER
1	A	99	GLN
1	A	100	ASN
1	A	111	LYS
1	A	114	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	155	MET
1	A	161	ARG
1	A	201	LEU
1	A	208	ILE
1	A	238	SER
1	A	257	SER
1	A	259	ILE
1	A	267	SER
1	A	268	GLN
1	B	32	LYS
1	B	35	ASP
1	B	47	ILE
1	B	49	LEU
1	B	55	LYS
1	B	58	GLU
1	B	95	ILE
1	B	104	GLU
1	B	169	LEU
1	B	194	LEU
1	B	195	THR
1	B	197	GLU
1	B	208	ILE
1	B	257	SER
1	B	268	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	96	HIS
1	A	100	ASN
1	A	112	HIS
1	A	154	ASN
1	A	170	ASN
1	A	268	GLN
1	B	153	HIS
1	B	175	ASN
1	B	206	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	CIT	B	1	-	12,12,12	1.35	2 (16%)	17,17,17	2.36	10 (58%)
2	CIT	A	1	-	12,12,12	1.18	0	17,17,17	1.54	3 (17%)
3	PEG	B	290	-	6,6,6	0.58	0	5,5,5	0.13	0
3	PEG	A	290	-	6,6,6	0.53	0	5,5,5	0.62	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CIT	B	1	-	-	10/16/16/16	-
2	CIT	A	1	-	-	14/16/16/16	-
3	PEG	B	290	-	-	3/4/4/4	-
3	PEG	A	290	-	-	3/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	CIT	C3-C6	2.63	1.56	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	CIT	C4-C3	2.42	1.57	1.54

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	CIT	O6-C6-C3	4.46	121.69	113.14
2	A	1	CIT	O6-C6-C3	3.82	120.46	113.14
2	B	1	CIT	O4-C5-C4	3.27	124.70	114.35
2	B	1	CIT	O7-C3-C2	-3.05	102.42	109.38
2	B	1	CIT	O6-C6-O5	-2.96	114.39	123.86
2	B	1	CIT	C3-C4-C5	2.83	121.67	113.92
2	B	1	CIT	C4-C3-C6	2.66	115.91	110.03
2	B	1	CIT	O7-C3-C6	2.58	112.62	108.96
2	B	1	CIT	O1-C1-C2	-2.41	116.13	122.95
2	A	1	CIT	O4-C5-C4	2.17	121.21	114.35
2	A	1	CIT	O5-C6-C3	-2.07	118.08	122.09
2	B	1	CIT	O4-C5-O3	-2.06	118.02	123.33
2	B	1	CIT	O3-C5-C4	-2.01	117.27	122.95

There are no chirality outliers.

All (30) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1	CIT	C2-C3-C4-C5
2	A	1	CIT	O7-C3-C4-C5
2	A	1	CIT	C6-C3-C4-C5
2	A	1	CIT	C2-C3-C6-O5
2	A	1	CIT	C2-C3-C6-O6
2	A	1	CIT	O7-C3-C6-O5
2	A	1	CIT	O7-C3-C6-O6
2	B	1	CIT	O7-C3-C4-C5
2	B	1	CIT	C6-C3-C4-C5
2	B	1	CIT	C2-C3-C6-O5
2	B	1	CIT	C2-C3-C6-O6
2	B	1	CIT	O7-C3-C6-O5
2	B	1	CIT	O7-C3-C6-O6
2	B	1	CIT	C2-C3-C4-C5
3	A	290	PEG	C1-C2-O2-C3
3	A	290	PEG	O2-C3-C4-O4
3	B	290	PEG	O2-C3-C4-O4
2	A	1	CIT	C1-C2-C3-O7
2	A	1	CIT	C1-C2-C3-C6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	1	CIT	C1-C2-C3-C4
3	A	290	PEG	O1-C1-C2-O2
2	A	1	CIT	C4-C3-C6-O5
2	A	1	CIT	C4-C3-C6-O6
2	B	1	CIT	C1-C2-C3-O7
3	B	290	PEG	O1-C1-C2-O2
3	B	290	PEG	C1-C2-O2-C3
2	A	1	CIT	C3-C4-C5-O4
2	B	1	CIT	C3-C4-C5-O4
2	A	1	CIT	C3-C4-C5-O3
2	B	1	CIT	C3-C4-C5-O3

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1	CIT	2	0
2	A	1	CIT	3	0
3	B	290	PEG	1	0
3	A	290	PEG	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	263/286 (91%)	-0.03	0	100 100	9, 20, 31, 39	0
1	B	263/286 (91%)	0.02	1 (0%)	88 90	11, 20, 35, 47	0
All	All	526/572 (91%)	-0.01	1 (0%)	91 92	9, 20, 33, 47	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	27	PRO	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CIT	A	1	13/13	0.70	0.21	67,71,73,73	0
2	CIT	B	1	13/13	0.72	0.19	46,49,54,58	0
3	PEG	B	290	7/7	0.81	0.13	49,60,65,67	0
3	PEG	A	290	7/7	0.82	0.15	42,43,46,50	0

6.5 Other polymers [i](#)

There are no such residues in this entry.