



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 19, 2026 – 12:32 AM UTC

PDB ID : 5P9H / pdb_00005p9h
Title : BTK1 COCRYSTALLIZED WITH RN983
Authors : Gardberg, A.S.
Deposited on : 2016-09-20
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	279	29% 83% 11% 5%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2043 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tyrosine-protein kinase BTK.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	264	Total	C	N	O	S	0	1	0
			1973	1265	325	363	20			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	381	GLY	-	expression tag	UNP Q06187

- Molecule 2 is 6- {tert}-butyl-8-fluoranyl-2-[3-(hydroxymethyl)-4-[1-methyl-5-[[5-(1-methylpyridin-4-yl)pyridin-2-yl]amino]-6-oxidanylidene-pyridazin-3-yl]pyridin-2-yl]phthalazin-1-one (CCD ID: 7G7) (formula: C₃₄H₃₇FN₈O₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	0	0
			46	34	1	8	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	24	Total 24	O 24	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein kinase BTK



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	62.76Å 101.40Å 38.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.36 – 1.95 24.36 – 1.95	Depositor EDS
% Data completeness (in resolution range)	85.2 (24.36-1.95) 85.2 (24.36-1.95)	Depositor EDS
R_{merge}	0.23	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.96 (at 1.95Å)	Xtriage
Refinement program	BUSTER	Depositor
R, R_{free}	0.267 , 0.313 0.293 , 0.333	Depositor DCC
R_{free} test set	782 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	25.4	Xtriage
Anisotropy	0.283	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 69.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	2043	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 7G7

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.84	0/2020	1.41	11/2749 (0.4%)

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	471	PHE	CA-CB-CG	7.16	120.96	113.80
1	A	579	ASP	CA-CB-CG	7.01	119.61	112.60
1	A	530	ASN	CA-CB-CG	6.38	118.98	112.60
1	A	560	PRO	CA-C-N	5.93	128.25	120.60
1	A	560	PRO	C-N-CA	5.93	128.25	120.60
1	A	555	VAL	CA-C-N	5.55	125.38	120.10
1	A	555	VAL	C-N-CA	5.55	125.38	120.10
1	A	645	LYS	CA-C-N	5.20	127.11	120.56
1	A	645	LYS	C-N-CA	5.20	127.11	120.56
1	A	653	ASP	CA-C-N	5.10	127.67	120.42
1	A	653	ASP	C-N-CA	5.10	127.67	120.42

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1973	0	1764	10	0
2	A	46	0	0	0	0
3	A	24	0	0	0	0
All	All	2043	0	1764	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:592:SER:HB3	1:A:595:LYS:HB2	1.69	0.75
1:A:399:PRO:HB3	1:A:467:GLN:HE22	1.56	0.69
1:A:501:MET:HG2	1:A:535:VAL:HG21	1.82	0.62
1:A:588:TRP:O	1:A:592:SER:HB2	2.06	0.56
1:A:577:LYS:HA	1:A:580:ILE:HB	1.90	0.53
1:A:456:LYS:HA	1:A:536:LYS:HG2	1.92	0.50
1:A:398:ASP:HB3	1:A:401:ASP:HB2	1.97	0.47
1:A:565:PRO:HG3	1:A:578:SER:HA	1.98	0.45
1:A:498:LEU:HB3	1:A:590:ILE:HG22	1.99	0.44
1:A:432:ILE:HD11	1:A:472:ILE:HD12	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	263/279 (94%)	256 (97%)	7 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	187/248 (75%)	178 (95%)	9 (5%)	23	12

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	410	THR
1	A	429	ILE
1	A	572	SER
1	A	586	LEU
1	A	592	SER
1	A	596[A]	MET
1	A	596[B]	MET
1	A	649	SER
1	A	654	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	412	GLN
1	A	459	GLN
1	A	467	GLN
1	A	484	ASN
1	A	650	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	7G7	A	701	-	50,51,51	0.92	2 (4%)	64,76,76	2.15	19 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	7G7	A	701	-	-	0/20/34/34	0/6/6/6

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	701	7G7	C29-N33	2.26	1.41	1.37
2	A	701	7G7	C7-C6	2.02	1.53	1.51

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	701	7G7	C28-C29-N33	6.76	118.37	112.05
2	A	701	7G7	C38-N39-C34	5.38	122.88	117.83
2	A	701	7G7	C13-C14-N15	-4.95	119.19	126.06
2	A	701	7G7	C14-N15-N10	4.60	122.50	114.15
2	A	701	7G7	C17-C16-C12	-4.40	119.75	123.96
2	A	701	7G7	C31-N27-C28	4.24	122.85	119.26
2	A	701	7G7	C9-N26-N27	4.06	123.08	118.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	701	7G7	C1-C9-N26	3.52	120.39	115.33
2	A	701	7G7	C37-C38-N39	-3.41	120.32	124.47
2	A	701	7G7	C44-N43-C42	3.27	114.75	109.54
2	A	701	7G7	C2-C3-N4	-3.27	119.97	123.97
2	A	701	7G7	C3-N4-C5	3.13	122.13	115.05
2	A	701	7G7	C35-C36-C37	-2.32	118.86	121.18
2	A	701	7G7	F21-C16-C12	2.32	122.21	118.15
2	A	701	7G7	C12-C13-C14	2.28	120.11	118.06
2	A	701	7G7	C36-C37-C38	2.11	119.06	116.89
2	A	701	7G7	C31-N27-N26	2.11	117.75	114.40
2	A	701	7G7	O32-C28-C29	-2.09	119.40	122.00
2	A	701	7G7	C18-C17-C16	2.06	121.50	119.46


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	264/279 (94%)	1.59	81 (30%) 1 1	26, 58, 100, 121	1 (0%)

All (81) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	421	TRP	5.4
1	A	656	ASP	5.0
1	A	607	ALA	4.5
1	A	602	THR	4.4
1	A	448	VAL	4.2
1	A	461	TYR	4.2
1	A	402	LEU	3.8
1	A	452	LEU	3.8
1	A	465	THR	3.8
1	A	601	PHE	3.8
1	A	471	PHE	3.7
1	A	470	ILE	3.6
1	A	614	LEU	3.6
1	A	611	ALA	3.6
1	A	442	PHE	3.6
1	A	397	ILE	3.4
1	A	657	GLU	3.3
1	A	429	ILE	3.3
1	A	425	TYR	3.2
1	A	404	PHE	3.2
1	A	395	TRP	3.2
1	A	446	ALA	3.1
1	A	604	SER	3.1
1	A	646	ILE	3.1
1	A	609	HIS	3.1
1	A	631	TYR	3.0
1	A	403	THR	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	422	ARG	2.9
1	A	557	SER	2.9
1	A	603	ASN	2.9
1	A	606	THR	2.8
1	A	658	GLU	2.8
1	A	654	VAL	2.8
1	A	400	LYS	2.7
1	A	610	ILE	2.7
1	A	574	PHE	2.7
1	A	453	SER	2.7
1	A	415	VAL	2.7
1	A	450	MET	2.6
1	A	399	PRO	2.6
1	A	469	PRO	2.6
1	A	642	PRO	2.6
1	A	613	GLY	2.6
1	A	416	VAL	2.6
1	A	535	VAL	2.6
1	A	555	VAL	2.5
1	A	617	TYR	2.5
1	A	549	ASP	2.5
1	A	458	VAL	2.5
1	A	419	GLY	2.4
1	A	655	MET	2.4
1	A	451	ASN	2.4
1	A	476	TYR	2.4
1	A	426	ASP	2.4
1	A	443	ILE	2.3
1	A	615	ARG	2.3
1	A	424	GLN	2.3
1	A	405	LEU	2.3
1	A	628	THR	2.3
1	A	467	GLN	2.3
1	A	413	PHE	2.2
1	A	605	GLU	2.2
1	A	580	ILE	2.2
1	A	420	LYS	2.2
1	A	547	LEU	2.2
1	A	652	LEU	2.2
1	A	398	ASP	2.2
1	A	626	VAL	2.2
1	A	423	GLY	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	583	PHE	2.2
1	A	466	LYS	2.2
1	A	600	ARG	2.2
1	A	481	CYS	2.2
1	A	569	LEU	2.2
1	A	473	ILE	2.1
1	A	460	LEU	2.1
1	A	586	LEU	2.1
1	A	644	PHE	2.1
1	A	449	MET	2.1
1	A	612	GLN	2.1
1	A	571	TYR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	7G7	A	701	46/46	0.71	0.16	40,50,72,75	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 7G7 A 701:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.