



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 02:29 PM UTC

PDB ID : 8PAJ / pdb_00008paj
Title : Crystal Structure of a Squalene-Hopene cyclase from Archangium gephyra
Authors : Worthy, H.L.; Isupov, M.N.; Littlechild, J.A.; Mitchell, D.E.
Deposited on : 2023-06-08
Resolution : 1.46 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.46 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1756 (1.46-1.46)
Clashscore	190562	1795 (1.46-1.46)
Ramachandran outliers	187476	1776 (1.46-1.46)
Sidechain outliers	187428	1776 (1.46-1.46)
RSRZ outliers	180081	1756 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	365	86% 7% 6%
1	B	365	2% 86% 8% 6%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 12247 atoms, of which 5709 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Squalene-hopene/tetraprenyl-beta-curcumene cyclase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	342	Total	C	H	N	O	S	82	31	0
			5645	1810	2804	533	488	10			
1	B	342	Total	C	H	N	O	S	83	32	0
			5662	1816	2813	531	492	10			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	383	ALA	-	expression tag	UNP A0A0G2ZXK3
A	384	SER	-	expression tag	UNP A0A0G2ZXK3
B	383	ALA	-	expression tag	UNP A0A0G2ZXK3
B	384	SER	-	expression tag	UNP A0A0G2ZXK3

- Molecule 2 is IMIDAZOLE (CCD ID: IMD) (formula: C₃H₅N₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	N	0	0
			10	3	5	2		
2	A	1	Total	C	H	N	0	0
			10	3	5	2		
2	B	1	Total	C	H	N	0	0
			10	3	5	2		
2	B	1	Total	C	H	N	0	0
			10	3	5	2		


- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	1	0
			10	2	6	2		
3	A	1	Total	C	H	O	1	0
			10	2	6	2		
3	A	1	Total	C	H	O	1	0
			10	2	6	2		
3	A	1	Total	C	H	O	1	0
			10	2	6	2		
3	B	1	Total	C	H	O	1	0
			10	2	6	2		
3	B	1	Total	C	H	O	1	0
			10	2	6	2		
3	B	1	Total	C	H	O	1	0
			10	2	6	2		

- Molecule 4 is 3[N-MORPHOLINO]PROPANE SULFONIC ACID (CCD ID: MPO) (formula:

C₇H₁₅NO₄S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total 28	C 7	H 15	N 1	O 4	S 1	1	0
4	B	1	Total 28	C 7	H 15	N 1	O 4	S 1	1	0

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Cl	0	0
			1	1		
5	B	1	Total	Cl	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	380	Total	O	0	0
			380	380		
6	B	392	Total	O	0	0
			392	392		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Squalene-hopene/tetraprenyl-beta-curcumene cyclase



- Molecule 1: Squalene-hopene/tetraprenyl-beta-curcumene cyclase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	92.96Å 84.17Å 87.97Å 90.00° 94.20° 90.00°	Depositor
Resolution (Å)	43.87 – 1.46 43.87 – 1.46	Depositor EDS
% Data completeness (in resolution range)	100.0 (43.87-1.46) 99.9 (43.87-1.46)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.09 (at 1.46Å)	Xtriage
Refinement program	REFMAC 5.8.0411	Depositor
R, R_{free}	0.159 , 0.199 0.159 , 0.199	Depositor DCC
R_{free} test set	5954 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	17.1	Xtriage
Anisotropy	0.504	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 42.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	12247	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, EDO, MPO, IMD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.75	0/3015	0.98	1/4109 (0.0%)
1	B	0.75	0/3016	0.98	1/4110 (0.0%)
All	All	0.75	0/6031	0.98	2/8219 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	352	THR	CA-CB-OG1	-5.77	100.94	109.60
1	A	352	THR	CA-CB-OG1	-5.64	101.14	109.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	67	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2841	2804	2816	38	0
1	B	2849	2813	2819	29	0
2	A	10	10	10	0	0
2	B	10	10	10	1	0
3	A	16	24	24	0	0
3	B	12	18	18	0	0
4	A	13	15	15	5	0
4	B	13	15	15	2	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	380	0	0	3	1
6	B	392	0	0	2	0
All	All	6538	5709	5727	67	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:309[B]:LEU:HD11	1:A:370[B]:PHE:CE2	1.24	1.63
1:B:306[B]:ASP:OD2	1:B:310[B]:ARG:CZ	1.77	1.31
1:A:309[B]:LEU:CD1	1:A:370[B]:PHE:CE2	2.12	1.30
1:A:309[B]:LEU:HD11	1:A:370[B]:PHE:CD2	1.86	1.09
1:A:309[B]:LEU:HD11	1:A:370[B]:PHE:CZ	1.90	1.05
1:A:309[B]:LEU:CD1	1:A:370[B]:PHE:CZ	2.38	1.04
1:B:309[B]:LEU:HD11	1:B:370[B]:PHE:CE2	2.01	0.95
1:B:43[B]:ARG:HG2	1:B:47[B]:LYS:HE3	1.57	0.86
1:B:222[A]:SER:OG	1:B:224[A]:GLU:OE1	1.97	0.82
1:B:306[B]:ASP:OD2	1:B:310[B]:ARG:NH1	2.13	0.81
1:B:309[B]:LEU:HD11	1:B:370[B]:PHE:CD2	2.19	0.77
1:A:309[B]:LEU:HD13	1:A:370[B]:PHE:CZ	2.20	0.75
1:A:310[B]:ARG:HG2	1:A:314[B]:ARG:HH21	1.50	0.74
1:B:309[B]:LEU:CD1	1:B:370[B]:PHE:CE2	2.74	0.70
1:A:310[B]:ARG:HH11	1:A:314[B]:ARG:NH2	1.89	0.70
1:A:128[B]:ARG:HD2	1:A:193[B]:LEU:HD21	1.73	0.70
1:B:251:ASP:OD2	2:B:402:IMD:H4	1.91	0.70
1:B:309[B]:LEU:HD11	1:B:370[B]:PHE:CZ	2.27	0.69
1:B:306[B]:ASP:OD2	1:B:310[B]:ARG:NE	2.24	0.68

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:306[B]:ASP:OD2	1:B:310[B]:ARG:NH2	2.25	0.68
1:B:309[B]:LEU:CD1	1:B:370[B]:PHE:CZ	2.76	0.68
1:A:256[B]:THR:HG22	1:A:295:ASP:OD1	1.94	0.67
1:A:207[B]:ARG:HG3	1:A:208:PRO:HD3	1.78	0.64
1:B:210[B]:LEU:HD21	1:B:249:LEU:HD21	1.82	0.62
1:A:138:GLU:HG3	4:A:405:MPO:H62	1.82	0.60
1:B:298[A]:ARG:NH1	1:B:374:SER:O	2.35	0.59
1:A:138:GLU:CG	4:A:405:MPO:H62	2.33	0.59
1:A:128[B]:ARG:HD2	1:A:193[B]:LEU:CD2	2.34	0.57
1:A:207[B]:ARG:NH1	1:A:211:GLU:HG3	2.20	0.56
1:A:328[A]:MET:HE3	1:A:351:LEU:HD12	1.87	0.55
1:A:85[B]:ARG:NH2	6:A:505:HOH:O	2.39	0.55
1:B:289:LEU:HD13	1:B:370[A]:PHE:HE2	1.71	0.55
1:B:289:LEU:HD13	1:B:370[A]:PHE:CE2	2.43	0.54
1:A:310[A]:ARG:NH1	6:A:503:HOH:O	2.27	0.54
1:A:34:VAL:HG21	1:A:309[A]:LEU:HD21	1.89	0.53
1:A:289:LEU:HD13	1:A:370[A]:PHE:HE2	1.71	0.53
1:A:210[B]:LEU:HD23	1:A:249:LEU:HD21	1.91	0.53
1:A:138:GLU:HB2	4:A:405:MPO:H62	1.91	0.53
1:B:256[B]:THR:HG22	1:B:295:ASP:OD1	2.08	0.53
1:A:310[B]:ARG:NH1	1:A:314[B]:ARG:NH2	2.57	0.52
1:A:34:VAL:CG2	1:A:309[A]:LEU:HD21	2.40	0.52
1:B:34:VAL:HG21	1:B:309[A]:LEU:HD21	1.91	0.51
1:A:289:LEU:HD13	1:A:370[A]:PHE:CE2	2.46	0.49
1:A:183:TRP:CH2	1:A:330[A]:MET:HE2	2.47	0.48
1:B:330[A]:MET:HE2	1:B:351[A]:LEU:HD11	1.96	0.47
1:A:330[B]:MET:HE2	1:A:351:LEU:HG	1.97	0.47
1:B:85[B]:ARG:NH1	6:B:509:HOH:O	2.48	0.46
1:B:317[A]:GLN:NE2	6:B:505:HOH:O	2.22	0.46
1:A:210[B]:LEU:CD2	1:A:249:LEU:HD21	2.46	0.46
1:A:310[B]:ARG:HH11	1:A:314[B]:ARG:HH21	1.61	0.46
1:B:198:LEU:HB2	1:B:210[A]:LEU:HD21	1.97	0.46
1:A:30:ARG:HB2	6:A:844:HOH:O	2.15	0.46
1:B:34:VAL:CG2	1:B:309[A]:LEU:HD21	2.46	0.45
1:B:138:GLU:HB2	4:B:404:MPO:H61	1.98	0.44
1:A:120:ALA:HB2	1:A:153:TYR:CG	2.53	0.43
1:A:138:GLU:HB2	4:A:405:MPO:C6	2.47	0.43
1:A:198:LEU:HB2	1:A:210[A]:LEU:HD21	2.00	0.43
1:B:30[A]:ARG:HG3	1:B:32[A]:SER:H	1.83	0.43
1:B:120:ALA:HB2	1:B:153:TYR:CG	2.55	0.41
1:B:60:PHE:CE2	1:B:330[A]:MET:HE1	2.55	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:85[B]:ARG:HH21	1:A:85[B]:ARG:HD3	1.73	0.41
1:A:138:GLU:CB	4:A:405:MPO:H62	2.50	0.41
1:A:128[B]:ARG:CD	1:A:193[B]:LEU:HD21	2.45	0.41
1:B:138:GLU:HG3	4:B:404:MPO:H62	2.03	0.41
1:B:309[B]:LEU:HD11	1:B:370[B]:PHE:CG	2.53	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:598:HOH:O	6:A:598:HOH:O[2_556]	1.58	0.62

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	369/365 (101%)	365 (99%)	4 (1%)	0	100	100
1	B	369/365 (101%)	363 (98%)	6 (2%)	0	100	100
All	All	738/730 (101%)	728 (99%)	10 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	285/272 (105%)	285 (100%)	0	100	100
1	B	286/272 (105%)	284 (99%)	2 (1%)	76	54
All	All	571/544 (105%)	569 (100%)	2 (0%)	87	70

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	330[A]	MET
1	B	330[B]	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	260	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 15 ligands modelled in this entry, 2 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	MPO	B	404	-	13,13,13	0.85	1 (7%)	17,17,17	0.99	1 (5%)
3	EDO	A	403	-	3,3,3	0.97	0	2,2,2	0.43	0
3	EDO	A	406	-	3,3,3	0.10	0	2,2,2	0.29	0
3	EDO	B	405	-	3,3,3	0.21	0	2,2,2	0.07	0
3	EDO	B	406	-	3,3,3	0.21	0	2,2,2	0.09	0
3	EDO	B	403	-	3,3,3	0.22	0	2,2,2	0.20	0
2	IMD	A	401	-	5,5,5	0.42	0	5,5,5	0.56	0
2	IMD	A	404	-	5,5,5	0.65	0	5,5,5	0.43	0
4	MPO	A	405	-	13,13,13	0.61	0	17,17,17	1.34	3 (17%)
2	IMD	B	401	-	5,5,5	0.27	0	5,5,5	0.67	0
3	EDO	A	402	-	3,3,3	0.16	0	2,2,2	0.19	0
2	IMD	B	402	-	5,5,5	0.48	0	5,5,5	0.40	0
3	EDO	A	407	-	3,3,3	0.26	0	2,2,2	0.20	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MPO	B	404	-	-	2/7/15/15	0/1/1/1
3	EDO	A	403	-	-	1/1/1/1	-
3	EDO	A	406	-	-	1/1/1/1	-
3	EDO	B	405	-	-	1/1/1/1	-
2	IMD	A	404	-	-	-	0/1/1/1
3	EDO	B	406	-	-	1/1/1/1	-
2	IMD	A	401	-	-	-	0/1/1/1
3	EDO	B	403	-	-	1/1/1/1	-
4	MPO	A	405	-	-	4/7/15/15	0/1/1/1
2	IMD	B	401	-	-	-	0/1/1/1
3	EDO	A	402	-	-	1/1/1/1	-
2	IMD	B	402	-	-	-	0/1/1/1
3	EDO	A	407	-	-	1/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	404	MPO	O3-S1	2.85	1.57	1.47

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	404	MPO	O3-S1-C1	-3.22	99.71	106.00
4	A	405	MPO	O3-S1-O2	-2.68	104.69	111.40
4	A	405	MPO	O4-C6-C7	2.21	116.53	111.77
4	A	405	MPO	O2-S1-O1	2.17	120.87	113.82

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	405	MPO	C1-C2-C3-N1
4	B	404	MPO	C1-C2-C3-N1
3	A	406	EDO	O1-C1-C2-O2
3	B	406	EDO	O1-C1-C2-O2
4	A	405	MPO	C2-C1-S1-O2
4	A	405	MPO	C2-C1-S1-O3
4	B	404	MPO	S1-C1-C2-C3
4	A	405	MPO	C2-C1-S1-O1
3	A	403	EDO	O1-C1-C2-O2
3	B	403	EDO	O1-C1-C2-O2
3	A	407	EDO	O1-C1-C2-O2
3	A	402	EDO	O1-C1-C2-O2
3	B	405	EDO	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	404	MPO	2	0
4	A	405	MPO	5	0
2	B	402	IMD	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	342/365 (93%)	-0.47	5 (1%) 72 75	9, 18, 38, 76	31 (9%)
1	B	342/365 (93%)	-0.42	8 (2%) 61 63	7, 19, 35, 76	32 (9%)
All	All	684/730 (93%)	-0.45	13 (1%) 66 68	7, 18, 37, 76	63 (9%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	63	VAL	4.3
1	B	30[A]	ARG	4.0
1	B	64	VAL	3.9
1	A	170	PRO	3.8
1	A	63	VAL	3.6
1	A	64	VAL	3.6
1	B	174	GLY	3.4
1	B	348	LEU	2.9
1	A	65	GLY	2.9
1	A	167	ALA	2.8
1	B	167	ALA	2.8
1	B	62	PHE	2.0
1	B	132[A]	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MPO	A	405	13/13	0.70	0.23	30,41,44,49	28
4	MPO	B	404	13/13	0.79	0.18	30,37,41,42	28
3	EDO	A	406	4/4	0.81	0.12	30,48,49,51	1
3	EDO	B	405	4/4	0.83	0.10	30,47,50,51	1
3	EDO	B	406	4/4	0.83	0.11	30,43,44,44	1
3	EDO	A	407	4/4	0.84	0.10	30,50,53,55	1
3	EDO	A	403	4/4	0.87	0.10	30,34,36,36	1
2	IMD	A	404	5/5	0.88	0.11	29,32,34,35	0
3	EDO	B	403	4/4	0.88	0.10	30,38,39,39	1
3	EDO	A	402	4/4	0.89	0.10	30,46,46,46	1
2	IMD	A	401	5/5	0.90	0.09	27,28,29,29	0
2	IMD	B	402	5/5	0.92	0.08	30,32,33,33	0
2	IMD	B	401	5/5	0.93	0.08	25,26,28,28	0
5	CL	A	408	1/1	0.99	0.12	35,35,35,35	0
5	CL	B	407	1/1	0.99	0.06	27,27,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.