



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 02:14 AM UTC

PDB ID : 1PB1 / pdb_00001pb1
Title : A four location model to explain the stereospecificity of proteins.
Authors : Mesecar, A.D.; Koshland Jr., D.E.
Deposited on : 2003-05-14
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	416	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ICT	A	601	X	-	-	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3586 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Isocitrate dehydrogenase [NADP].


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	416	Total	C	N	O	S	0	0	0
			3213	2045	540	609	19			

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is ISOCITRIC ACID (CCD ID: ICT) (formula: C₆H₈O₇).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			13	6	7		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	337	Total 337	O 337	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: Isocitrate dehydrogenase [NADP]

Chain A:  85% 12%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	103.65Å 103.65Å 149.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.79 – 1.70	Depositor
% Data completeness (in resolution range)	92.7 (24.79-1.70)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.185 , 0.211	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3586	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ICT, GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.83	1/3274 (0.0%)	1.04	11/4427 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	380	MET	SD-CE	-5.91	1.64	1.79

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	10	GLN	N-CA-C	-8.31	98.22	110.52
1	A	337	ALA	N-CA-C	-7.40	99.33	110.28
1	A	69	TYR	N-CA-C	7.14	121.14	109.72
1	A	389	VAL	N-CA-C	6.50	118.07	108.65
1	A	23	VAL	N-CA-C	6.06	114.56	107.77
1	A	96	ARG	N-CA-C	5.55	122.63	110.80
1	A	223	ASP	N-CA-C	5.50	119.15	112.23
1	A	125	TYR	N-CA-C	5.43	119.53	113.01
1	A	336	GLU	N-CA-C	5.40	117.29	108.76
1	A	391	TYR	N-CA-C	5.38	117.84	111.33
1	A	359	SER	N-CA-C	-5.36	105.48	112.23

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3213	0	3240	44	0
2	A	5	0	0	0	0
3	A	13	0	5	0	0
4	A	18	0	24	0	0
5	A	337	0	0	4	0
All	All	3586	0	3269	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:397:MET:HE2	1:A:400:ALA:HB2	1.47	0.94
1:A:169:SER:O	1:A:173:GLU:HG2	1.78	0.84
1:A:288:GLN:HE22	1:A:292:ARG:HD2	1.45	0.81
1:A:387:LYS:HG2	1:A:397:MET:HE1	1.73	0.68
1:A:401:LYS:HD3	1:A:403:LEU:HD21	1.76	0.68
1:A:4:LYS:HG3	1:A:78:TYR:CE2	2.29	0.67
1:A:2:GLU:H	1:A:2:GLU:CD	2.02	0.67
1:A:76:GLN:HG2	5:A:1025:HOH:O	1.93	0.67
1:A:288:GLN:NE2	1:A:292:ARG:HD2	2.12	0.63
1:A:344:LYS:HD2	1:A:345:TYR:CZ	2.34	0.62
1:A:229:HIS:CE1	1:A:281:ILE:HD13	2.37	0.59
1:A:345:TYR:CE2	1:A:350:LYS:HE2	2.39	0.58
1:A:350:LYS:HE3	1:A:391:TYR:CE2	2.39	0.57
1:A:1:MET:HE1	1:A:4:LYS:HE3	1.89	0.55
1:A:4:LYS:HG3	1:A:78:TYR:HE2	1.69	0.55
1:A:349:ASP:OD1	1:A:405:CYS:HB3	2.07	0.54
1:A:350:LYS:HE3	1:A:391:TYR:CD2	2.42	0.54
1:A:265:LYS:HB2	1:A:274:GLU:HG3	1.90	0.54
1:A:13:LYS:NZ	1:A:94:GLU:OE2	2.33	0.52
1:A:2:GLU:CD	1:A:2:GLU:N	2.67	0.52
1:A:169:SER:O	1:A:173:GLU:CG	2.56	0.51
1:A:20:LYS:HE2	5:A:1031:HOH:O	2.09	0.50
1:A:267:LYS:HD2	1:A:272:GLY:C	2.36	0.50
1:A:273:LYS:HD2	5:A:865:HOH:O	2.12	0.49
1:A:1:MET:HG3	1:A:2:GLU:O	2.12	0.49
1:A:30:PRO:HA	1:A:66:MET:O	2.13	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:151:ILE:HD11	1:A:313:LEU:HD12	1.95	0.49
1:A:4:LYS:N	1:A:78:TYR:OH	2.39	0.48
1:A:393:PHE:O	1:A:397:MET:HG2	2.14	0.47
1:A:93:ARG:O	1:A:93:ARG:HD2	2.13	0.47
1:A:387:LYS:HG2	1:A:397:MET:CE	2.44	0.46
1:A:167:ALA:O	1:A:168:ASP:HB2	2.15	0.46
1:A:30:PRO:HD2	1:A:97:VAL:O	2.16	0.45
1:A:1:MET:SD	1:A:2:GLU:O	2.75	0.44
1:A:4:LYS:CG	1:A:78:TYR:HE2	2.28	0.44
1:A:165:TRP:CD1	1:A:171:ASP:HB3	2.53	0.44
1:A:267:LYS:HD2	1:A:272:GLY:HA2	2.00	0.43
1:A:1:MET:CE	1:A:4:LYS:HE3	2.48	0.43
1:A:117:ALA:O	1:A:121:GLU:CG	2.68	0.42
1:A:1:MET:HE1	1:A:4:LYS:CE	2.49	0.42
1:A:37:ILE:HG21	1:A:351:VAL:HG11	2.00	0.42
1:A:37:ILE:HB	1:A:351:VAL:HG21	2.02	0.42
1:A:25:GLU:HG3	5:A:1117:HOH:O	2.20	0.41
1:A:4:LYS:CG	1:A:78:TYR:CE2	3.00	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	414/416 (100%)	399 (96%)	15 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	338/338 (100%)	325 (96%)	13 (4%)	29	13

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	2	GLU
1	A	4	LYS
1	A	20	LYS
1	A	54	GLU
1	A	157	GLU
1	A	173	GLU
1	A	187	LYS
1	A	256	GLU
1	A	267	LYS
1	A	273	LYS
1	A	274	GLU
1	A	344	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	80	GLN
1	A	115	ASN
1	A	135	GLN
1	A	143	HIS
1	A	287	GLN
1	A	288	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	A	706	-	5,5,5	0.53	0	5,5,5	0.41	0
4	GOL	A	703	-	5,5,5	0.73	0	5,5,5	0.18	0
3	ICT	A	601	-	12,12,12	2.52	4 (33%)	13,16,16	1.78	3 (23%)
2	SO4	A	801	-	4,4,4	0.70	0	6,6,6	0.25	0
4	GOL	A	704	-	5,5,5	0.58	0	5,5,5	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	703	-	-	2/4/4/4	-
4	GOL	A	704	-	-	2/4/4/4	-
3	ICT	A	601	-	2/2/5/5	7/16/16/16	-
4	GOL	A	706	-	-	2/4/4/4	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	601	ICT	O1-C1	4.65	1.35	1.22
3	A	601	ICT	O5-C6	4.38	1.35	1.22
3	A	601	ICT	O3-C5	3.38	1.33	1.22
3	A	601	ICT	C3-C2	2.91	1.58	1.54

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	601	ICT	C4-C3-C6	-3.26	106.06	111.25
3	A	601	ICT	O6-C6-C3	2.53	120.49	113.93
3	A	601	ICT	O4-C5-C4	2.33	121.23	114.00

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	601	ICT	C3
3	A	601	ICT	C2

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	703	GOL	O1-C1-C2-C3
4	A	704	GOL	O1-C1-C2-O2
4	A	704	GOL	O1-C1-C2-C3
4	A	706	GOL	C1-C2-C3-O3
4	A	706	GOL	O2-C2-C3-O3
4	A	703	GOL	O1-C1-C2-O2
3	A	601	ICT	C6-C3-C4-C5
3	A	601	ICT	C2-C3-C6-O5
3	A	601	ICT	C2-C3-C6-O6
3	A	601	ICT	C4-C3-C6-O5
3	A	601	ICT	C4-C3-C6-O6
3	A	601	ICT	C3-C4-C5-O3
3	A	601	ICT	C3-C4-C5-O4

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.