



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2026 – 05:57 PM UTC

PDB ID : 3PBE / pdb_00003pbe
Title : Crystal structure of the mutant W207F of human secretory glutaminyl cyclase
Authors : Huang, K.F.; Liaw, S.S.; Huang, W.L.; Chia, C.Y.; Lo, Y.C.; Chen, Y.L.; Wang, A.H.J.
Deposited on : 2010-10-20
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	329	2% 88% 9% ..
1	B	329	5% 86% 11% ..

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5733 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutaminyl-peptide cyclotransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	323	Total	C	N	O	S	5	0	0
			2607	1671	449	478	9			
1	B	323	Total	C	N	O	S	5	0	0
			2607	1671	449	478	9			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	207	PHE	TRP	engineered mutation	UNP Q16769
B	207	PHE	TRP	engineered mutation	UNP Q16769

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	289	Total	O	0	0
			289	289		
4	B	218	Total	O	0	0
			218	218		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glutaminyl-peptide cyclotransferase



- Molecule 1: Glutaminyl-peptide cyclotransferase



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	118.75Å 118.75Å 332.09Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 1.95 30.00 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.8 (30.00-1.95) 96.0 (30.00-1.95)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.92 (at 1.95Å)	Xtriage
Refinement program	CNS, REFMAC 5.5.0102	Depositor
R, R_{free}	0.143 , 0.190 0.172 , 0.167	Depositor DCC
R_{free} test set	3340 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	31.4	Xtriage
Anisotropy	0.077	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 37.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5733	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.59	12/2683 (0.4%)	1.15	4/3649 (0.1%)
1	B	1.45	13/2683 (0.5%)	1.11	4/3649 (0.1%)
All	All	1.52	25/5366 (0.5%)	1.13	8/7298 (0.1%)

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	308	ILE	CA-CB	14.32	1.61	1.54
1	B	308	ILE	CA-CB	9.11	1.58	1.54
1	A	241	MET	SD-CE	-8.59	1.58	1.79
1	A	162	VAL	CA-CB	8.24	1.58	1.54
1	A	100	ALA	CA-CB	7.87	1.65	1.53
1	A	252	ALA	CA-CB	6.71	1.63	1.53
1	A	46	ALA	CA-CB	6.32	1.61	1.53
1	A	40	LYS	CE-NZ	6.31	1.68	1.49
1	B	165	ALA	CA-CB	6.14	1.63	1.53
1	B	337	GLU	CG-CD	6.06	1.67	1.52
1	B	223	MET	CA-C	5.79	1.60	1.52
1	B	235	THR	CA-C	5.67	1.59	1.52
1	B	232	ALA	CA-CB	5.58	1.62	1.53
1	A	130	THR	C-O	5.50	1.31	1.24
1	A	337	GLU	CG-CD	5.42	1.65	1.52
1	B	315	VAL	CA-C	5.41	1.57	1.53
1	A	132	LYS	C-O	-5.24	1.18	1.24
1	B	337	GLU	CB-CG	5.24	1.68	1.52
1	B	55	GLN	CA-C	5.19	1.59	1.52
1	A	319	HIS	C-O	5.10	1.29	1.23
1	B	70	LEU	CG-CD1	5.10	1.69	1.52
1	B	74	LEU	CA-CB	5.08	1.58	1.52
1	A	171	ALA	C-O	5.06	1.29	1.24
1	B	221	ALA	N-CA	5.05	1.52	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	155	VAL	CA-CB	5.03	1.60	1.54

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	321	ILE	CA-C-N	-7.19	113.27	120.31
1	A	321	ILE	C-N-CA	-7.19	113.27	120.31
1	A	130	THR	CA-C-O	-6.54	111.40	119.18
1	B	131	ALA	CA-C-N	6.24	128.55	120.44
1	B	131	ALA	C-N-CA	6.24	128.55	120.44
1	A	337	GLU	CB-CG-CD	5.66	122.22	112.60
1	B	337	GLU	CB-CG-CD	5.50	121.94	112.60
1	B	130	THR	O-C-N	5.42	129.84	122.37

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2607	0	2528	21	0
1	B	2607	0	2528	21	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	289	0	0	4	0
4	B	218	0	0	6	0
All	All	5733	0	5056	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:40:LYS:CE	1:A:40:LYS:NZ	1.68	1.54
1:A:236:SER:H	1:A:239:HIS:CD2	2.00	0.80
1:A:236:SER:H	1:A:239:HIS:HD2	1.33	0.75
1:A:312:ARG:HH11	1:A:312:ARG:HG2	1.51	0.75
1:A:337:GLU:HG2	4:A:557:HOH:O	1.93	0.68
1:B:292:ARG:HD2	4:B:884:HOH:O	1.96	0.65
1:B:337:GLU:HG2	4:B:720:HOH:O	1.96	0.64
1:A:312:ARG:HH11	1:A:312:ARG:CG	2.13	0.61
1:B:177:LYS:NZ	4:B:855:HOH:O	2.18	0.58
1:B:258:PRO:HG2	1:B:260:PHE:CZ	2.39	0.57
1:A:112:GLN:HE21	1:A:113:THR:H	1.54	0.55
1:B:296:ASN:ND2	4:B:884:HOH:O	2.40	0.54
1:B:302:VAL:CG1	1:B:307:HIS:HE1	2.22	0.52
1:A:292:ARG:NE	4:A:877:HOH:O	2.32	0.52
1:A:296:ASN:O	1:A:297:TYR:HB3	2.10	0.52
1:A:40:LYS:NZ	1:A:40:LYS:CD	2.66	0.52
1:B:269:GLU:HA	1:B:272:GLN:HE21	1.75	0.51
1:B:305:ASP:HB3	1:B:307:HIS:CD2	2.47	0.50
1:A:41:ASN:ND2	4:A:409:HOH:O	2.44	0.50
1:A:258:PRO:HG2	1:A:260:PHE:CZ	2.47	0.50
1:A:112:GLN:HE21	1:A:112:GLN:HA	1.77	0.49
1:B:307:HIS:CD2	1:B:307:HIS:H	2.30	0.49
1:B:279:HIS:HD2	1:B:287:HIS:ND1	2.10	0.49
1:B:300:GLY:CA	4:B:896:HOH:O	2.61	0.48
1:B:296:ASN:ND2	1:B:296:ASN:H	2.12	0.47
1:B:74:LEU:HD13	1:B:336:GLU:HB2	1.97	0.47
1:B:276:HIS:HB2	1:B:292:ARG:HH22	1.80	0.46
1:B:295:GLN:O	1:B:297:TYR:N	2.48	0.46
1:A:210:GLN:HB2	4:A:874:HOH:O	2.16	0.46
1:A:132:LYS:HG3	1:A:190:ASP:OD2	2.16	0.45
1:B:49:ASN:OD1	1:B:49:ASN:C	2.59	0.45
1:A:241:MET:HE2	1:A:315:VAL:HG11	1.99	0.45
1:B:305:ASP:HB3	1:B:307:HIS:HD2	1.82	0.44
1:B:300:GLY:HA3	4:B:896:HOH:O	2.16	0.44
1:B:149:TRP:O	1:B:150:ASN:HB2	2.18	0.44
1:A:92:MET:SD	1:A:106:ILE:HD11	2.58	0.43
1:A:241:MET:HE2	1:A:241:MET:HB3	1.85	0.43
1:A:259:ASN:HD21	1:A:265:ALA:HA	1.84	0.43
1:A:36:TRP:N	1:A:37:PRO:CD	2.82	0.41
1:B:258:PRO:HG2	1:B:260:PHE:CE1	2.54	0.41
1:B:307:HIS:H	1:B:307:HIS:HD2	1.69	0.41
1:A:112:GLN:HE22	1:A:116:GLY:HA2	1.85	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	319/329 (97%)	310 (97%)	8 (2%)	1 (0%)	36	28
1	B	319/329 (97%)	308 (97%)	10 (3%)	1 (0%)	36	28
All	All	638/658 (97%)	618 (97%)	18 (3%)	2 (0%)	36	28

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	296	ASN
1	A	212	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	284/290 (98%)	282 (99%)	2 (1%)	76	76
1	B	284/290 (98%)	275 (97%)	9 (3%)	34	25
All	All	568/580 (98%)	557 (98%)	11 (2%)	50	45

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	112	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	312	ARG
1	B	34	SER
1	B	40	LYS
1	B	163	PRO
1	B	206	HIS
1	B	242	ASP
1	B	284	LEU
1	B	297	TYR
1	B	302	VAL
1	B	337	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	41	ASN
1	A	44	GLN
1	A	55	GLN
1	A	93	GLN
1	A	112	GLN
1	A	150	ASN
1	A	239	HIS
1	A	259	ASN
1	A	276	HIS
1	B	259	ASN
1	B	272	GLN
1	B	279	HIS
1	B	296	ASN
1	B	304	GLN
1	B	307	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	395	-	4,4,4	0.35	0	6,6,6	0.69	0
3	SO4	B	396	-	4,4,4	0.77	0	6,6,6	0.47	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	322/329 (97%)	-0.24	8 (2%) 58 65	22, 30, 47, 63	0
1	B	322/329 (97%)	0.22	17 (5%) 32 37	25, 37, 58, 80	0
All	All	644/658 (97%)	-0.01	25 (3%) 43 50	22, 32, 54, 80	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	33	ALA	4.9
1	B	189	PRO	4.1
1	B	297	TYR	3.8
1	A	297	TYR	3.7
1	B	33	ALA	3.7
1	A	149	TRP	3.3
1	A	207	PHE	3.0
1	B	190	ASP	2.9
1	B	149	TRP	2.7
1	B	301	GLY	2.6
1	B	337	GLU	2.5
1	B	130	THR	2.5
1	B	286	ASP	2.5
1	A	189	PRO	2.5
1	B	300	GLY	2.4
1	B	52	ALA	2.4
1	A	296	ASN	2.3
1	A	300	GLY	2.3
1	B	117	TYR	2.2
1	A	150	ASN	2.2
1	B	282	GLY	2.2
1	B	49	ASN	2.2
1	B	283	LEU	2.2
1	B	281	LEU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	314	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	B	396	5/5	0.96	0.10	38,42,43,44	0
3	SO4	A	395	5/5	0.97	0.07	47,47,51,51	0
2	ZN	B	392	1/1	0.99	0.02	29,29,29,29	0
2	ZN	A	391	1/1	1.00	0.01	27,27,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.