



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 03:43 PM UTC

PDB ID : 4PB4 / pdb_00004pb4
Title : D-threo-3-hydroxyaspartate dehydratase H351A mutant complexed with 2-amino maleic acid
Authors : Yasutake, Y.; Matsumoto, Y.; Wada, M.
Deposited on : 2014-04-11
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	390	% 87% 9% • •
1	B	390	4% 87% 12% •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	2KZ	A	403	-	X	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6277 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called D-threo-3-hydroxyaspartate dehydratase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	379	Total	C	N	O	S	0	6	0
			2857	1775	533	534	15			
1	B	389	Total	C	N	O	S	0	6	0
			2940	1826	552	544	18			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-9	GLY	-	expression tag	UNP B2DFG5
A	-8	HIS	-	expression tag	UNP B2DFG5
A	-7	HIS	-	expression tag	UNP B2DFG5
A	-6	HIS	-	expression tag	UNP B2DFG5
A	-5	HIS	-	expression tag	UNP B2DFG5
A	-4	HIS	-	expression tag	UNP B2DFG5
A	-3	HIS	-	expression tag	UNP B2DFG5
A	-2	ALA	-	expression tag	UNP B2DFG5
A	-1	MET	-	expression tag	UNP B2DFG5
A	0	SER	-	expression tag	UNP B2DFG5
A	351	ALA	HIS	engineered mutation	UNP B2DFG5
B	-9	GLY	-	expression tag	UNP B2DFG5
B	-8	HIS	-	expression tag	UNP B2DFG5
B	-7	HIS	-	expression tag	UNP B2DFG5
B	-6	HIS	-	expression tag	UNP B2DFG5
B	-5	HIS	-	expression tag	UNP B2DFG5
B	-4	HIS	-	expression tag	UNP B2DFG5
B	-3	HIS	-	expression tag	UNP B2DFG5
B	-2	ALA	-	expression tag	UNP B2DFG5
B	-1	MET	-	expression tag	UNP B2DFG5
B	0	SER	-	expression tag	UNP B2DFG5
B	351	ALA	HIS	engineered mutation	UNP B2DFG5

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (CCD ID: PLP) (formula: C₈H₁₀NO₆P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		
3	B	1	Total	Mg	0	0
			1	1		

- Molecule 4 is 2-amino maleic acid (CCD ID: 2KZ) (formula: C₄H₅NO₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			9	4	1	4		
4	B	1	Total	C	N	O	0	0
			9	4	1	4		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	215	Total	O	0	0
			215	215		
5	B	215	Total	O	0	0
			215	215		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: D-threo-3-hydroxyaspartate dehydratase



- Molecule 1: D-threo-3-hydroxyaspartate dehydratase



4 Data and refinement statistics

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	157.75Å 157.75Å 158.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.21 – 1.80 37.21 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.5 (37.21-1.80) 99.6 (37.21-1.80)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.06 (at 1.81Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.164 , 0.183 (Not available) , 0.167	Depositor DCC
R_{free} test set	4544 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	17.6	Xtriage
Anisotropy	0.007	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 38.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.002 for -l,-k,-h 0.004 for -h,l,k	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6277	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: 2KZ, PLP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.56	12/2921 (0.4%)	1.30	8/3962 (0.2%)
1	B	1.66	18/3010 (0.6%)	1.33	10/4080 (0.2%)
All	All	1.61	30/5931 (0.5%)	1.31	18/8042 (0.2%)

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	-6	HIS	CG-CD2	12.08	1.49	1.35
1	B	-6	HIS	CE1-NE2	11.44	1.44	1.32
1	B	204	ARG	CD-NE	-7.62	1.35	1.46
1	A	267	HIS	ND1-CE1	7.48	1.40	1.32
1	B	141	ARG	CA-C	7.41	1.62	1.52
1	B	172	HIS	ND1-CE1	7.23	1.39	1.32
1	A	321	HIS	ND1-CE1	7.18	1.39	1.32
1	A	172	HIS	ND1-CE1	6.99	1.39	1.32
1	B	145	GLY	N-CA	6.53	1.54	1.45
1	B	142	SER	CA-C	6.24	1.61	1.53
1	A	316	GLY	N-CA	6.02	1.51	1.45
1	A	140	HIS	ND1-CE1	5.96	1.38	1.32
1	A	172	HIS	CG-CD2	5.95	1.42	1.35
1	A	341	LEU	N-CA	5.86	1.53	1.46
1	B	92	LEU	CA-C	5.79	1.59	1.52
1	B	83	TYR	N-CA	5.66	1.53	1.46
1	A	233	VAL	CA-C	-5.58	1.46	1.52
1	B	107	LEU	N-CA	5.52	1.52	1.45
1	B	112	VAL	CA-C	5.36	1.59	1.52
1	A	318	ASN	CG-OD1	-5.33	1.13	1.23
1	B	202	ALA	N-CA	5.33	1.52	1.46
1	A	321	HIS	CG-CD2	5.30	1.41	1.35
1	B	116	GLN	C-O	5.28	1.30	1.24
1	B	92	LEU	C-O	5.26	1.29	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	307	ARG	N-CA	5.25	1.52	1.46
1	A	334	ASP	C-O	5.22	1.30	1.23
1	B	230	VAL	N-CA	5.12	1.52	1.46
1	B	141	ARG	CZ-NH2	5.07	1.40	1.33
1	B	-6	HIS	ND1-CE1	5.01	1.37	1.32
1	B	166	LEU	CA-CB	5.01	1.59	1.53

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	204	ARG	NE-CZ-NH2	-8.12	111.89	119.20
1	B	231	THR	OG1-CB-CG2	-6.74	95.82	109.30
1	A	310	PRO	CA-C-N	-6.40	113.12	122.46
1	A	310	PRO	C-N-CA	-6.40	113.12	122.46
1	B	149	THR	CA-C-N	-6.32	112.04	119.05
1	B	149	THR	C-N-CA	-6.32	112.04	119.05
1	A	55	ARG	NE-CZ-NH2	6.31	124.88	119.20
1	A	217	SER	N-CA-C	-6.27	99.54	109.07
1	B	217	SER	N-CA-C	-5.99	100.22	109.14
1	A	334	ASP	N-CA-C	5.90	119.32	111.30
1	B	140	HIS	N-CA-C	5.65	117.11	111.07
1	B	215	VAL	CA-C-N	-5.54	115.06	122.54
1	B	215	VAL	C-N-CA	-5.54	115.06	122.54
1	B	106	LYS	CD-CE-NZ	-5.22	95.20	111.90
1	A	338	ARG	CG-CD-NE	-5.21	100.55	112.00
1	B	142	SER	N-CA-C	-5.15	103.89	110.53
1	A	149	THR	CA-C-N	-5.07	113.42	119.05
1	A	149	THR	C-N-CA	-5.07	113.42	119.05

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2857	0	2868	22	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2940	0	2940	41	0
2	A	15	0	6	0	0
2	B	15	0	6	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	9	0	1	2	0
4	B	9	0	1	2	0
5	A	215	0	0	5	0
5	B	215	0	0	4	0
All	All	6277	0	5822	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:30:ARG:HH11	1:A:30:ARG:CG	1.53	1.20
1:B:245[B]:MET:CE	1:B:245[B]:MET:HA	1.72	1.17
4:B:403:2KZ:CB	5:B:712:HOH:O	1.90	1.17
1:B:245[B]:MET:HA	1:B:245[B]:MET:HE3	1.21	1.12
1:B:100[B]:ARG:HH11	1:B:100[B]:ARG:CG	1.61	1.11
4:A:403:2KZ:CB	5:A:709:HOH:O	2.02	1.07
1:A:30:ARG:HG3	1:A:30:ARG:NH1	1.33	1.03
1:B:100[B]:ARG:HH11	1:B:100[B]:ARG:HG2	0.86	1.01
1:B:100[B]:ARG:HG2	1:B:100[B]:ARG:NH1	1.67	0.97
1:B:245[B]:MET:HE3	1:B:248:ILE:HD12	1.63	0.81
1:A:2:GLN:CA	1:A:2:GLN:HE21	1.95	0.77
1:B:245[B]:MET:HA	1:B:245[B]:MET:HE2	1.66	0.77
1:A:2:GLN:HE21	1:A:2:GLN:HA	1.50	0.76
1:B:303:ASP:OD2	1:B:307:ARG:NH1	2.17	0.76
1:B:100[B]:ARG:CG	1:B:100[B]:ARG:NH1	2.33	0.74
1:A:2:GLN:HA	1:A:2:GLN:NE2	2.04	0.72
1:B:68[A]:LYS:HG2	1:B:377:LEU:HD21	1.70	0.72
1:B:307:ARG:HH11	1:B:307:ARG:HB2	1.55	0.71
1:A:94[A]:GLN:HG3	5:A:648:HOH:O	1.94	0.67
1:A:51:ALA:HA	1:A:54:GLN:HE21	1.59	0.67
1:B:245[B]:MET:CE	1:B:245[B]:MET:CA	2.55	0.66
1:B:245[B]:MET:HE3	1:B:245[B]:MET:CA	2.13	0.64
1:A:18:ASP:HB3	1:A:22:ARG:NH2	2.14	0.62
1:B:245[B]:MET:CE	1:B:248:ILE:HD12	2.30	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:219:PRO:HG3	1:B:245[B]:MET:HE1	1.82	0.60
1:B:291:ARG:HD2	1:B:291:ARG:H	1.66	0.60
1:B:334:ASP:OD2	1:B:337:THR:HG23	2.02	0.60
1:B:51:ALA:HA	1:B:54:GLN:HE21	1.65	0.60
1:B:291:ARG:HD2	1:B:291:ARG:N	2.16	0.60
4:B:403:2KZ:CA	5:B:712:HOH:O	2.34	0.59
1:A:18:ASP:OD1	5:A:659:HOH:O	2.17	0.59
1:B:140:HIS:HD2	5:B:614:HOH:O	1.85	0.59
1:B:245[B]:MET:HE2	1:B:245[B]:MET:CA	2.29	0.58
1:B:307:ARG:NH1	1:B:307:ARG:HB2	2.22	0.55
4:A:403:2KZ:CA	5:A:709:HOH:O	2.40	0.53
1:B:307:ARG:HH11	1:B:307:ARG:CB	2.20	0.53
1:A:30:ARG:CG	1:A:30:ARG:NH1	2.23	0.52
1:B:337:THR:HG23	5:B:693:HOH:O	2.10	0.52
1:B:132:TRP:CH2	1:B:212:VAL:HG11	2.46	0.49
1:A:318:ASN:OD1	1:B:141:ARG:HD3	2.11	0.49
1:A:333:ALA:O	1:A:334:ASP:C	2.55	0.48
1:B:43:LYS:HE3	2:B:401:PLP:C4A	2.44	0.48
1:B:245[B]:MET:HE2	1:B:250:VAL:HG22	1.96	0.47
1:A:68:LYS:HG2	1:A:377:LEU:HD21	1.97	0.47
1:A:132:TRP:CH2	1:A:212:VAL:HG11	2.52	0.45
1:B:237:VAL:HG11	1:B:245[A]:MET:HE3	1.98	0.45
1:A:85:VAL:O	1:A:86:SER:C	2.60	0.44
1:B:68[A]:LYS:HD3	1:B:377:LEU:HG	2.00	0.44
1:B:176:SER:HB3	1:B:188:LEU:HD23	2.00	0.44
1:A:11:PRO:HA	1:A:261:LEU:O	2.18	0.44
1:A:75:ALA:HB2	5:A:507:HOH:O	2.18	0.44
1:B:96:LEU:HD21	1:B:100[A]:ARG:CZ	2.47	0.44
1:A:291:ARG:H	1:A:291:ARG:HG2	1.49	0.43
1:B:30:ARG:HG2	1:B:31:MET:HE2	2.01	0.43
1:A:215:VAL:O	1:A:233:VAL:HA	2.18	0.43
1:A:170:LEU:HA	1:A:214:SER:O	2.19	0.42
1:B:-4:HIS:HA	1:B:-1:MET:HE2	2.00	0.42
1:B:-8:HIS:O	1:B:-4:HIS:CD2	2.73	0.42
1:A:141:ARG:HD3	1:B:318:ASN:OD1	2.20	0.42
1:A:309:MET:HE1	1:A:339:PHE:CZ	2.56	0.41
1:B:100[B]:ARG:NH1	1:B:127:GLU:OE2	2.52	0.41
1:B:-8:HIS:HB3	1:B:-4:HIS:CD2	2.56	0.41
1:B:338:ARG:HA	1:B:338:ARG:HD3	1.86	0.41
1:B:68[A]:LYS:CG	1:B:377:LEU:HD21	2.45	0.40
1:B:170:LEU:HA	1:B:214:SER:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	383/390 (98%)	373 (97%)	8 (2%)	2 (0%)	24	14
1	B	393/390 (101%)	383 (98%)	9 (2%)	1 (0%)	36	25
All	All	776/780 (100%)	756 (97%)	17 (2%)	3 (0%)	30	19

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	141	ARG
1	A	141	ARG
1	A	236	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	220/288 (76%)	217 (99%)	3 (1%)	59	52
1	B	151/288 (52%)	148 (98%)	3 (2%)	48	38
All	All	371/576 (64%)	365 (98%)	6 (2%)	53	47

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	GLN
1	A	30	ARG
1	A	268	GLN
1	B	291	ARG
1	B	334	ASP
1	B	337	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2	GLN
1	A	28	GLN
1	A	54	GLN
1	A	140	HIS
1	B	363	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PLP	A	401	4	15,15,16	2.42	4 (26%)	21,22,23	1.69	6 (28%)
4	2KZ	A	403	3,2	7,8,8	3.22	4 (57%)	8,10,10	2.44	4 (50%)
2	PLP	B	401	4	15,15,16	1.82	4 (26%)	21,22,23	1.71	5 (23%)
4	2KZ	B	403	3,2	7,8,8	2.11	2 (28%)	8,10,10	3.28	4 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	A	401	4	-	0/6/6/8	0/1/1/1
4	2KZ	A	403	3,2	-	3/6/8/8	-
2	PLP	B	401	4	-	0/6/6/8	0/1/1/1
4	2KZ	B	403	3,2	-	3/6/8/8	-

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	PLP	C5-C4	5.81	1.47	1.40
4	A	403	2KZ	OXT-C	-5.06	1.16	1.30
4	A	403	2KZ	OD2-CG	5.04	1.35	1.23
2	A	401	PLP	C3-C2	5.00	1.46	1.41
4	B	403	2KZ	OXT-C	-4.63	1.18	1.30
2	B	401	PLP	C3-C2	3.98	1.45	1.41
2	A	401	PLP	C3-C4	3.78	1.47	1.40
2	B	401	PLP	C3-C4	3.61	1.47	1.40
4	A	403	2KZ	OD1-CG	-3.25	1.22	1.30
4	A	403	2KZ	O-C	-2.69	1.16	1.22
2	B	401	PLP	C5-C4	2.67	1.43	1.40
2	B	401	PLP	C6-C5	2.22	1.42	1.37
4	B	403	2KZ	O-C	-2.06	1.17	1.22
2	A	401	PLP	O3-C3	-2.03	1.32	1.36

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	403	2KZ	OXT-C-CA	5.27	120.22	112.41
4	B	403	2KZ	OD1-CG-CB	5.10	128.54	113.40
4	A	403	2KZ	OXT-C-CA	4.66	119.31	112.41
4	B	403	2KZ	OD2-CG-CB	-4.47	110.33	124.02

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	PLP	C2A-C2-C3	-3.64	116.53	120.80
2	B	401	PLP	C2A-C2-N1	3.47	124.17	117.64
4	A	403	2KZ	OXT-C-O	-3.28	116.08	123.90
2	A	401	PLP	C2A-C2-N1	3.23	123.73	117.64
2	A	401	PLP	C6-N1-C2	3.14	124.89	119.20
2	B	401	PLP	C6-N1-C2	2.87	124.40	119.20
4	B	403	2KZ	O-C-CA	-2.83	117.86	123.66
2	A	401	PLP	C6-C5-C4	-2.75	115.84	118.10
4	A	403	2KZ	CB-CA-C	2.71	127.16	120.17
2	A	401	PLP	C2A-C2-C3	-2.67	117.67	120.80
2	A	401	PLP	O3-C3-C4	2.22	123.90	118.10
2	B	401	PLP	O2P-P-O4P	-2.19	100.97	106.67
4	A	403	2KZ	OD1-CG-CB	-2.09	107.18	113.40
2	A	401	PLP	O3P-P-O2P	2.04	115.47	107.80
2	B	401	PLP	O3P-P-O2P	2.04	115.44	107.80

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	403	2KZ	CA-CB-CG-OD1
4	A	403	2KZ	N-CA-CB-CG
4	B	403	2KZ	CA-CB-CG-OD1
4	B	403	2KZ	N-CA-CB-CG
4	A	403	2KZ	CA-CB-CG-OD2
4	B	403	2KZ	CA-CB-CG-OD2

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	403	2KZ	2	0
2	B	401	PLP	1	0
4	B	403	2KZ	2	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	379/390 (97%)	-0.50	3 (0%) 82 83	6, 18, 37, 62	6 (1%)
1	B	389/390 (99%)	-0.37	15 (3%) 43 43	7, 18, 41, 99	6 (1%)
All	All	768/780 (98%)	-0.43	18 (2%) 61 61	6, 18, 38, 99	12 (1%)

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	-8	HIS	6.8
1	B	-7	HIS	5.5
1	B	-5	HIS	4.7
1	B	-1	MET	4.3
1	B	-6	HIS	4.0
1	A	2	GLN	3.8
1	B	330	ALA	3.7
1	B	-4	HIS	3.6
1	B	1	MET	3.0
1	B	291	ARG	3.0
1	B	-2	ALA	3.0
1	B	333	ALA	3.0
1	B	0	SER	2.6
1	B	-3	HIS	2.5
1	A	147	ASP	2.4
1	B	142	SER	2.3
1	A	291	ARG	2.2
1	B	2	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MG	B	402	1/1	0.86	0.13	25,25,25,25	0
4	2KZ	A	403	9/9	0.89	0.09	22,25,34,34	0
4	2KZ	B	403	9/9	0.92	0.09	20,24,30,35	0
3	MG	A	402	1/1	0.96	0.12	25,25,25,25	0
2	PLP	A	401	15/16	0.98	0.04	15,16,18,21	0
2	PLP	B	401	15/16	0.99	0.04	13,14,17,20	0

6.5 Other polymers [i](#)

There are no such residues in this entry.