



Full wwPDB EM Validation Report ⓘ

Mar 27, 2026 – 06:17 PM UTC

PDB ID : 7PBP / pdb_00007pbp
EMDB ID : EMD-13298
Title : RuvAB branch migration motor complexed to the Holliday junction - RuvB
AAA+ state s5 [t2 dataset]
Authors : Fahrenkamp, D.; Goessweiner-Mohr, N.; Wald, J.; Marlovits, T.C.
Deposited on : 2021-08-02
Resolution : 3.20 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	15020 (2.70 - 3.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	315	5% 97% ..
1	B	315	.. 96% ..
1	C	315	5% 92% 6% .
1	D	315	13% 94% 5% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	315	22% 95% . .
1	F	315	12% 96% . .
2	G	54	85% 85% . 11%
2	H	54	81% 91% . 7%
3	U	15	93% 7%
4	V	15	13% 80% 20%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 32155 atoms, of which 15896 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Holliday junction ATP-dependent DNA helicase RuvB.

Mol	Chain	Residues	Atoms						AltConf	Trace
1	A	312	Total	C	H	N	O	S	0	0
			4910	1555	2455	412	472	16		
1	B	312	Total	C	H	N	O	S	0	0
			4910	1555	2455	412	472	16		
1	C	309	Total	C	H	N	O	S	0	0
			4895	1549	2451	412	467	16		
1	D	313	Total	C	H	N	O	S	0	0
			4934	1561	2468	416	473	16		
1	E	312	Total	C	H	N	O	S	0	0
			4910	1555	2455	412	472	16		
1	F	312	Total	C	H	N	O	S	0	0
			4910	1555	2455	412	472	16		

- Molecule 2 is a protein called Holliday junction ATP-dependent DNA helicase RuvA.

Mol	Chain	Residues	Atoms						AltConf	Trace
2	G	48	Total	C	H	N	O	S	0	0
			718	218	362	64	73	1		
2	H	50	Total	C	H	N	O	S	0	0
			760	234	379	74	72	1		

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	204	HIS	-	expression tag	UNP A0A0M0QTS9
G	205	HIS	-	expression tag	UNP A0A0M0QTS9
G	206	HIS	-	expression tag	UNP A0A0M0QTS9
G	207	HIS	-	expression tag	UNP A0A0M0QTS9
G	208	HIS	-	expression tag	UNP A0A0M0QTS9
G	209	HIS	-	expression tag	UNP A0A0M0QTS9
H	204	HIS	-	expression tag	UNP A0A0M0QTS9
H	205	HIS	-	expression tag	UNP A0A0M0QTS9
H	206	HIS	-	expression tag	UNP A0A0M0QTS9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	207	HIS	-	expression tag	UNP A0A0M0QTS9
H	208	HIS	-	expression tag	UNP A0A0M0QTS9
H	209	HIS	-	expression tag	UNP A0A0M0QTS9


- Molecule 3 is a DNA chain called random DNA.

Mol	Chain	Residues	Atoms						AltConf	Trace
3	U	15	Total	C	H	N	O	P	0	0
			479	147	167	63	87	15		

- Molecule 4 is a DNA chain called random DNA.


Mol	Chain	Residues	Atoms						AltConf	Trace
4	V	15	Total	C	H	N	O	P	0	0
			474	145	171	50	93	15		

- Molecule 5 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms						AltConf
5	A	1	Total	C	H	N	O	P	0
			39	10	12	5	10	2	
5	E	1	Total	C	H	N	O	P	0
			39	10	12	5	10	2	
5	F	1	Total	C	H	N	O	P	0
			39	10	12	5	10	2	

- Molecule 6 is PHOSPHOTHIOPHOSPHORIC ACID-ADENYLATE ESTER (CCD ID: AGS) (formula: C₁₀H₁₆N₅O₁₂P₃S).



Mol	Chain	Residues	Atoms							AltConf
6	B	1	Total	C	H	N	O	P	S	0
			45	10	14	5	12	3	1	
6	C	1	Total	C	H	N	O	P	S	0
			45	10	14	5	12	3	1	
6	D	1	Total	C	H	N	O	P	S	0
			45	10	14	5	12	3	1	


- Molecule 7 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
7	B	1	Total	Mg	0
			1	1	
7	C	1	Total	Mg	0
			1	1	
7	D	1	Total	Mg	0
			1	1	


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Holliday junction ATP-dependent DNA helicase RuvB




- Molecule 1: Holliday junction ATP-dependent DNA helicase RuvB



- Molecule 1: Holliday junction ATP-dependent DNA helicase RuvB




- Molecule 1: Holliday junction ATP-dependent DNA helicase RuvB




- Molecule 1: Holliday junction ATP-dependent DNA helicase RuvB









• Molecule 1: Holliday junction ATP-dependent DNA helicase RuvB



• Molecule 2: Holliday junction ATP-dependent DNA helicase RuvA





• Molecule 2: Holliday junction ATP-dependent DNA helicase RuvA



• Molecule 3: random DNA



• Molecule 4: random DNA



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	125425	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	30.7	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	4000	Depositor
Magnification	Not provided	
Image detector	GATAN K2 QUANTUM (4k x 4k)	Depositor
Maximum map value	0.151	Depositor
Minimum map value	-0.057	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.025	Depositor
Map size (Å)	392.40002, 392.40002, 392.40002	wwPDB
Map dimensions	360, 360, 360	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.09, 1.09, 1.09	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ADP, AGS, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.19	0/2498	0.41	0/3377
1	B	0.21	0/2498	0.43	0/3377
1	C	0.20	0/2486	0.41	0/3359
1	D	0.17	0/2509	0.42	0/3391
1	E	0.15	0/2498	0.40	0/3377
1	F	0.16	0/2498	0.38	0/3377
2	G	0.10	0/358	0.29	0/483
2	H	0.09	0/387	0.29	0/523
3	U	0.25	0/351	0.53	0/540
4	V	0.25	0/337	0.65	0/517
All	All	0.18	0/16420	0.41	0/22321

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	1
1	F	0	2
All	All	0	3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	222	ARG	Sidechain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	F	160	ARG	Sidechain
1	F	315	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2455	2455	2454	4	0
1	B	2455	2455	2453	6	0
1	C	2444	2451	2448	13	0
1	D	2466	2468	2467	10	0
1	E	2455	2455	2454	7	0
1	F	2455	2455	2454	4	0
2	G	356	362	361	1	0
2	H	381	379	378	1	0
3	U	312	167	168	1	0
4	V	303	171	171	2	0
5	A	27	12	12	0	0
5	E	27	12	12	0	0
5	F	27	12	12	0	0
6	B	31	14	12	0	0
6	C	31	14	12	0	0
6	D	31	14	12	0	0
7	B	1	0	0	0	0
7	C	1	0	0	0	0
7	D	1	0	0	0	0
All	All	16259	15896	15880	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:121:GLU:OE2	1:D:114:ARG:NH1	2.19	0.76
1:A:299:GLU:OE1	1:A:315:ARG:NH1	2.24	0.70

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:81:LYS:NZ	1:F:100:ASP:O	2.25	0.69
2:G:189:ALA:HB1	2:G:193:THR:OG1	2.00	0.61
1:F:167:PRO:O	1:F:171:ARG:NH1	2.43	0.51
1:C:78:VAL:HG12	1:C:79:ASN:N	2.27	0.50
1:B:254:ASP:C	1:B:254:ASP:OD1	2.55	0.49
1:C:98:LEU:HD13	1:C:132:ILE:HD11	1.95	0.49
1:D:313:THR:HG22	1:D:313:THR:O	2.12	0.49
1:F:186:LEU:HD11	1:F:215:GLY:HA2	1.93	0.48
1:C:311:THR:HG22	1:C:312:ARG:N	2.29	0.47
4:V:6:DC:H2"	4:V:7:DT:H71	1.95	0.47
1:B:121:GLU:OE1	1:C:114:ARG:HG3	2.16	0.46
1:B:212:ARG:NH1	1:B:253:VAL:O	2.49	0.46
1:E:143:ARG:NH1	2:H:162:GLU:OE2	2.48	0.46
1:A:290:GLU:OE1	1:A:293:THR:N	2.46	0.45
1:C:78:VAL:CG1	1:C:79:ASN:N	2.80	0.45
1:D:185:ASP:OD1	1:D:186:LEU:N	2.49	0.45
1:E:129:ASP:HB3	1:E:131:TYR:CE2	2.52	0.44
1:C:254:ASP:OD2	1:C:324:HIS:NE2	2.50	0.44
1:D:78:VAL:CG1	1:D:79:ASN:N	2.80	0.44
1:F:120:GLU:OE2	1:F:124:TYR:OH	2.33	0.44
1:E:167:PRO:O	1:E:171:ARG:NH1	2.51	0.44
1:C:164:LEU:O	1:C:169:ARG:NH1	2.52	0.43
1:E:254:ASP:OD2	1:E:324:HIS:NE2	2.52	0.43
1:B:186:LEU:HD11	1:B:215:GLY:HA2	1.99	0.43
1:C:159:THR:HG22	1:C:159:THR:O	2.18	0.43
3:U:11:DG:O6	4:V:5:DC:N4	2.52	0.43
1:C:134:ILE:CG2	1:C:135:MET:N	2.81	0.43
1:B:81:LYS:NZ	1:B:100:ASP:O	2.51	0.43
1:D:129:ASP:HB3	1:D:131:TYR:CE2	2.54	0.42
1:D:280:LEU:HD11	1:D:294:VAL:HG23	2.01	0.42
1:A:240:ASP:OD1	1:A:242:LYS:N	2.52	0.42
1:E:213:SER:O	1:E:214:ARG:HG2	2.19	0.42
1:E:241:ASP:OD1	1:E:242:LYS:N	2.52	0.42
1:A:164:LEU:O	1:A:169:ARG:NH1	2.53	0.42
1:D:311:THR:HG22	1:D:312:ARG:N	2.34	0.42
1:C:256:GLU:HG3	1:C:324:HIS:CE1	2.55	0.41
1:E:159:THR:O	1:E:159:THR:HG22	2.21	0.41
1:D:60:PRO:HG2	1:D:63:LEU:HD11	2.02	0.41
1:D:143:ARG:HG3	1:D:143:ARG:HH11	1.86	0.41
1:D:253:VAL:HG13	1:D:253:VAL:O	2.20	0.41
1:C:197:PHE:C	1:C:198:GLU:HG2	2.46	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:199:MET:SD	1:B:228:ARG:HG3	2.61	0.40
1:C:197:PHE:O	1:C:198:GLU:HG2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	310/315 (98%)	307 (99%)	3 (1%)	0	100	100
1	B	310/315 (98%)	306 (99%)	4 (1%)	0	100	100
1	C	305/315 (97%)	304 (100%)	1 (0%)	0	100	100
1	D	311/315 (99%)	306 (98%)	5 (2%)	0	100	100
1	E	310/315 (98%)	307 (99%)	3 (1%)	0	100	100
1	F	310/315 (98%)	309 (100%)	1 (0%)	0	100	100
2	G	46/54 (85%)	45 (98%)	1 (2%)	0	100	100
2	H	48/54 (89%)	47 (98%)	1 (2%)	0	100	100
All	All	1950/1998 (98%)	1931 (99%)	19 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	261/264 (99%)	261 (100%)	0	100	100
1	B	261/264 (99%)	260 (100%)	1 (0%)	84	86
1	C	261/264 (99%)	260 (100%)	1 (0%)	84	86
1	D	262/264 (99%)	262 (100%)	0	100	100
1	E	261/264 (99%)	261 (100%)	0	100	100
1	F	261/264 (99%)	261 (100%)	0	100	100
2	G	36/42 (86%)	36 (100%)	0	100	100
2	H	38/42 (90%)	38 (100%)	0	100	100
All	All	1641/1668 (98%)	1639 (100%)	2 (0%)	87	91

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	143	ARG
1	C	286	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	146	HIS
1	A	255	HIS
1	A	286	ASN
1	B	146	HIS
1	B	286	ASN
1	C	286	ASN
1	D	23	GLN
1	D	113	HIS
1	D	175	ASN
1	D	177	HIS
1	D	255	HIS
1	D	324	HIS
1	E	146	HIS
1	F	324	HIS
2	H	161	GLN
2	H	205	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 3 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
6	AGS	C	401	7	32,33,33	0.66	0	45,52,52	0.62	1 (2%)
5	ADP	F	600	-	28,29,29	1.35	5 (17%)	43,45,45	1.83	10 (23%)
6	AGS	B	401	7	32,33,33	0.74	1 (3%)	45,52,52	0.65	1 (2%)
5	ADP	A	600	-	28,29,29	1.37	5 (17%)	43,45,45	1.84	8 (18%)
6	AGS	D	401	7	32,33,33	0.54	0	45,52,52	0.61	1 (2%)
5	ADP	E	600	-	28,29,29	1.38	5 (17%)	43,45,45	1.83	9 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	AGS	C	401	7	-	2/21/38/38	0/3/3/3
5	ADP	F	600	-	-	1/16/32/32	0/3/3/3
6	AGS	B	401	7	-	3/21/38/38	0/3/3/3
5	ADP	A	600	-	-	1/16/32/32	0/3/3/3
6	AGS	D	401	7	-	5/21/38/38	0/3/3/3
5	ADP	E	600	-	-	2/16/32/32	0/3/3/3

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	E	600	ADP	C5-C4	4.42	1.47	1.39
5	A	600	ADP	C5-C4	4.28	1.46	1.39
5	F	600	ADP	C5-C4	4.18	1.46	1.39
5	A	600	ADP	C5-N7	-2.69	1.34	1.39
5	F	600	ADP	C5-N7	-2.66	1.34	1.39
5	E	600	ADP	C5-C6	2.66	1.48	1.41
5	A	600	ADP	C5-C6	2.47	1.47	1.41
5	E	600	ADP	C5-N7	-2.40	1.34	1.39
5	E	600	ADP	C8-N7	2.39	1.36	1.31
5	F	600	ADP	C5-C6	2.36	1.47	1.41
6	B	401	AGS	PB-O3A	-2.23	1.57	1.59
5	A	600	ADP	C8-N7	2.13	1.35	1.31
5	E	600	ADP	C4-N9	-2.13	1.33	1.37
5	F	600	ADP	C8-N7	2.12	1.35	1.31
5	F	600	ADP	C4-N9	-2.11	1.33	1.37
5	A	600	ADP	C4-N9	-2.03	1.33	1.37

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	600	ADP	C5-C4-N3	-6.05	118.38	126.72
5	F	600	ADP	C5-C4-N3	-5.86	118.65	126.72
5	E	600	ADP	C5-C4-N3	-5.80	118.73	126.72
5	A	600	ADP	N3-C4-N9	4.80	135.33	127.17
5	E	600	ADP	N3-C4-N9	4.39	134.63	127.17
5	F	600	ADP	N3-C4-N9	4.35	134.57	127.17
5	A	600	ADP	C2-N3-C4	3.92	121.41	111.83
5	E	600	ADP	C2-N3-C4	3.90	121.34	111.83
5	F	600	ADP	C2-N3-C4	3.85	121.24	111.83
5	F	600	ADP	N3-C2-N1	-3.72	122.95	128.58
5	E	600	ADP	C4-C5-N7	-3.66	106.40	110.58
5	A	600	ADP	N3-C2-N1	-3.64	123.06	128.58
5	E	600	ADP	N3-C2-N1	-3.64	123.07	128.58
5	F	600	ADP	C4-C5-N7	-3.64	106.42	110.58
5	A	600	ADP	C4-C5-N7	-3.48	106.61	110.58
6	D	401	AGS	PB-O3B-PG	-3.35	120.90	133.17
6	B	401	AGS	PB-O3B-PG	-3.13	121.73	133.17
6	C	401	AGS	PB-O3B-PG	-2.95	122.36	133.17
5	F	600	ADP	C5-N7-C8	2.74	107.76	103.45
5	A	600	ADP	C4-N9-C8	2.73	108.61	105.74
5	E	600	ADP	C4-N9-C8	2.73	108.61	105.74
5	A	600	ADP	C5-N7-C8	2.71	107.70	103.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	E	600	ADP	C5-N7-C8	2.69	107.69	103.45
5	E	600	ADP	C6-C5-N7	2.40	136.71	132.09
5	F	600	ADP	C4-N9-C8	2.33	108.18	105.74
5	E	600	ADP	N9-C8-N7	-2.30	110.67	113.94
5	A	600	ADP	N9-C8-N7	-2.22	110.79	113.94
5	F	600	ADP	N9-C8-N7	-2.17	110.85	113.94
5	F	600	ADP	C6-C5-N7	2.17	136.27	132.09
5	F	600	ADP	C2-N1-C6	2.05	122.10	118.73

There are no chirality outliers.

All (14) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	600	ADP	PA-O3A-PB-O2B
6	B	401	AGS	C5'-O5'-PA-O2A
6	D	401	AGS	C5'-O5'-PA-O2A
5	F	600	ADP	PA-O3A-PB-O3B
6	D	401	AGS	PA-O3A-PB-O1B
6	B	401	AGS	C5'-O5'-PA-O1A
6	D	401	AGS	C5'-O5'-PA-O1A
6	D	401	AGS	C5'-O5'-PA-O3A
6	C	401	AGS	PA-O3A-PB-O2B
6	D	401	AGS	PA-O3A-PB-O2B
5	E	600	ADP	PA-O3A-PB-O1B
5	E	600	ADP	PA-O3A-PB-O2B
6	B	401	AGS	PA-O3A-PB-O2B
6	C	401	AGS	PA-O3A-PB-O1B

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient


equivalents in the CSD to analyse the geometry.




Ligand AGS B 401




Ligand ADP A 600



Ligand AGS D 401



Ligand ADP E 600



5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-13298. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 180



Y Index: 180




Z Index: 180


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 162



Y Index: 156




Z Index: 176


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.025. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 73 nm^3 ; this corresponds to an approximate mass of 66 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.312 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.312 $\text{\AA}^{-1}$

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.20	-	-
Author-provided FSC curve	3.16	3.52	3.20
Unmasked-calculated*	-	-	-

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-13298 and PDB model 7PBP. Per-residue inclusion information can be found in section 3 on page 7.


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.025 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.025).

9.4 Atom inclusion [i](#)



At the recommended contour level, 69% of all backbone atoms, 69% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.025) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.6870	0.5280
A	0.7890	0.5680
B	0.8160	0.5770
C	0.7790	0.5620
D	0.6780	0.5200
E	0.5780	0.4810
F	0.6620	0.5200
G	0.1440	0.3150
H	0.2230	0.3160
U	0.8400	0.5700
V	0.6800	0.5060

1.0

0.0

<0.0