



# wwPDB X-ray Structure Validation Summary Report

Mar 10, 2026 – 07:22 PM UTC

PDB ID : 6PCC / pdb\_00006pcc  
Title : Crystal structure of beta-ketoadipyl-CoA thiolase mutant (H356A) in complex  
hexanoyl coenzyme A  
Authors : Sukritee, B.; Panjikar, S.  
Deposited on : 2019-06-17  
Resolution : 1.96 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*





The reported resolution of this entry is 1.96 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	423	 85% 10% 5%
1	B	423	 86% 8% 6%
1	C	423	 86% 7% 6%
1	D	423	 85% 9% 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	O8Y	A	1002	-	-	X	-
3	O8Y	B	502	-	-	X	-
3	O8Y	C	502	-	-	X	-
3	O8Y	D	502	-	-	X	-

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 13336 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-ketoacyl-CoA thiolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	403	2972	1849	539	566	18	0	3	0
1	B	398	2975	1849	539	567	20	0	7	0
1	C	396	2934	1825	529	561	19	0	5	0
1	D	402	2996	1863	543	572	18	0	7	0

There are 96 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-22	MET	-	initiating methionine	UNP Q88N39
A	-21	HIS	-	expression tag	UNP Q88N39
A	-20	HIS	-	expression tag	UNP Q88N39
A	-19	HIS	-	expression tag	UNP Q88N39
A	-18	HIS	-	expression tag	UNP Q88N39
A	-17	HIS	-	expression tag	UNP Q88N39
A	-16	HIS	-	expression tag	UNP Q88N39
A	-15	SER	-	expression tag	UNP Q88N39
A	-14	SER	-	expression tag	UNP Q88N39
A	-13	GLY	-	expression tag	UNP Q88N39
A	-12	VAL	-	expression tag	UNP Q88N39
A	-11	ASP	-	expression tag	UNP Q88N39
A	-10	LEU	-	expression tag	UNP Q88N39
A	-9	GLY	-	expression tag	UNP Q88N39
A	-8	THR	-	expression tag	UNP Q88N39
A	-7	GLU	-	expression tag	UNP Q88N39
A	-6	ASN	-	expression tag	UNP Q88N39
A	-5	LEU	-	expression tag	UNP Q88N39
A	-4	TYR	-	expression tag	UNP Q88N39
A	-3	HIS	-	expression tag	UNP Q88N39
A	-2	GLN	-	expression tag	UNP Q88N39

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP Q88N39
A	0	SER	-	expression tag	UNP Q88N39
A	356	ALA	HIS	engineered mutation	UNP Q88N39
B	-22	MET	-	initiating methionine	UNP Q88N39
B	-21	HIS	-	expression tag	UNP Q88N39
B	-20	HIS	-	expression tag	UNP Q88N39
B	-19	HIS	-	expression tag	UNP Q88N39
B	-18	HIS	-	expression tag	UNP Q88N39
B	-17	HIS	-	expression tag	UNP Q88N39
B	-16	HIS	-	expression tag	UNP Q88N39
B	-15	SER	-	expression tag	UNP Q88N39
B	-14	SER	-	expression tag	UNP Q88N39
B	-13	GLY	-	expression tag	UNP Q88N39
B	-12	VAL	-	expression tag	UNP Q88N39
B	-11	ASP	-	expression tag	UNP Q88N39
B	-10	LEU	-	expression tag	UNP Q88N39
B	-9	GLY	-	expression tag	UNP Q88N39
B	-8	THR	-	expression tag	UNP Q88N39
B	-7	GLU	-	expression tag	UNP Q88N39
B	-6	ASN	-	expression tag	UNP Q88N39
B	-5	LEU	-	expression tag	UNP Q88N39
B	-4	TYR	-	expression tag	UNP Q88N39
B	-3	HIS	-	expression tag	UNP Q88N39
B	-2	GLN	-	expression tag	UNP Q88N39
B	-1	GLY	-	expression tag	UNP Q88N39
B	0	SER	-	expression tag	UNP Q88N39
B	356	ALA	HIS	engineered mutation	UNP Q88N39
C	-22	MET	-	initiating methionine	UNP Q88N39
C	-21	HIS	-	expression tag	UNP Q88N39
C	-20	HIS	-	expression tag	UNP Q88N39
C	-19	HIS	-	expression tag	UNP Q88N39
C	-18	HIS	-	expression tag	UNP Q88N39
C	-17	HIS	-	expression tag	UNP Q88N39
C	-16	HIS	-	expression tag	UNP Q88N39
C	-15	SER	-	expression tag	UNP Q88N39
C	-14	SER	-	expression tag	UNP Q88N39
C	-13	GLY	-	expression tag	UNP Q88N39
C	-12	VAL	-	expression tag	UNP Q88N39
C	-11	ASP	-	expression tag	UNP Q88N39
C	-10	LEU	-	expression tag	UNP Q88N39
C	-9	GLY	-	expression tag	UNP Q88N39
C	-8	THR	-	expression tag	UNP Q88N39

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
C	-7	GLU	-	expression tag	UNP Q88N39
C	-6	ASN	-	expression tag	UNP Q88N39
C	-5	LEU	-	expression tag	UNP Q88N39
C	-4	TYR	-	expression tag	UNP Q88N39
C	-3	HIS	-	expression tag	UNP Q88N39
C	-2	GLN	-	expression tag	UNP Q88N39
C	-1	GLY	-	expression tag	UNP Q88N39
C	0	SER	-	expression tag	UNP Q88N39
C	356	ALA	HIS	engineered mutation	UNP Q88N39
D	-22	MET	-	initiating methionine	UNP Q88N39
D	-21	HIS	-	expression tag	UNP Q88N39
D	-20	HIS	-	expression tag	UNP Q88N39
D	-19	HIS	-	expression tag	UNP Q88N39
D	-18	HIS	-	expression tag	UNP Q88N39
D	-17	HIS	-	expression tag	UNP Q88N39
D	-16	HIS	-	expression tag	UNP Q88N39
D	-15	SER	-	expression tag	UNP Q88N39
D	-14	SER	-	expression tag	UNP Q88N39
D	-13	GLY	-	expression tag	UNP Q88N39
D	-12	VAL	-	expression tag	UNP Q88N39
D	-11	ASP	-	expression tag	UNP Q88N39
D	-10	LEU	-	expression tag	UNP Q88N39
D	-9	GLY	-	expression tag	UNP Q88N39
D	-8	THR	-	expression tag	UNP Q88N39
D	-7	GLU	-	expression tag	UNP Q88N39
D	-6	ASN	-	expression tag	UNP Q88N39
D	-5	LEU	-	expression tag	UNP Q88N39
D	-4	TYR	-	expression tag	UNP Q88N39
D	-3	HIS	-	expression tag	UNP Q88N39
D	-2	GLN	-	expression tag	UNP Q88N39
D	-1	GLY	-	expression tag	UNP Q88N39
D	0	SER	-	expression tag	UNP Q88N39
D	356	ALA	HIS	engineered mutation	UNP Q88N39

- Molecule 2 is COENZYME A (CCD ID: COA) (formula:  $C_{21}H_{36}N_7O_{16}P_3S$ ) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	P			S
2	A	1	Total	C	N	O	P	S	0	0
			48	21	7	16	3	1		
2	B	1	Total	C	N	O	P	S	0	0
			48	21	7	16	3	1		
2	C	1	Total	C	N	O	P	S	0	0
			48	21	7	16	3	1		
2	D	1	Total	C	N	O	P	S	0	0
			48	21	7	16	3	1		

- Molecule 3 is hexanal (CCD ID: O8Y) (formula: C<sub>6</sub>H<sub>12</sub>O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	6	1		
3	B	1	Total	C	O	0	0
			7	6	1		
3	C	1	Total	C	O	0	0
			7	6	1		
3	D	1	Total	C	O	0	0
			7	6	1		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula:  $C_3H_8O_3$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	C	1	Total	C	O	0	0
			6	3	3		
4	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	282	Total	O	0	0
			282	282		

*Continued on next page...*


*Continued from previous page...*


<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
5	B	280	Total 280	O 280	0	0
5	C	328	Total 328	O 328	0	0
5	D	325	Total 325	O 325	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Beta-ketoadipyl-CoA thiolase

Chain A: 




- Molecule 1: Beta-ketoadipyl-CoA thiolase

Chain B: 





- Molecule 1: Beta-ketoadipyl-CoA thiolase

Chain C: 



- Molecule 1: Beta-ketoadipyl-CoA thiolase

Chain D: 





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	111.39Å 116.40Å 127.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.94 – 1.96 19.94 – 1.96	Depositor EDS
% Data completeness (in resolution range)	95.3 (19.94-1.96) 95.3 (19.94-1.96)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.33 (at 1.96Å)	Xtrriage
Refinement program	REFMAC 5.8.0238	Depositor
R, $R_{free}$	0.176 , 0.208 (Not available) , 0.184	Depositor DCC
$R_{free}$ test set	968 reflections (0.81%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.0	Xtrriage
Anisotropy	0.049	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 43.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.003 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	13336	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.96% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CSO, GOL, COA, O8Y

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.61	0/3006	1.02	2/4066 (0.0%)
1	B	0.60	0/3001	1.04	4/4060 (0.1%)
1	C	0.61	0/2959	1.02	0/4004
1	D	0.63	0/3030	1.01	2/4099 (0.0%)
All	All	0.61	0/11996	1.02	8/16229 (0.0%)

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	16	ARG	CG-CD-NE	-7.67	95.13	112.00
1	B	32	VAL	N-CA-CB	6.15	115.25	110.45
1	D	228	GLU	CB-CG-CD	5.77	122.41	112.60
1	B	228	GLU	CB-CG-CD	5.76	122.40	112.60
1	A	399	ARG	CA-C-N	5.46	131.52	121.70

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2972	0	3006	41	0
1	B	2975	0	2998	23	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2934	0	2961	25	0
1	D	2996	0	3023	32	0
2	A	48	0	31	0	0
2	B	48	0	31	0	0
2	C	48	0	31	0	0
2	D	48	0	31	2	0
3	A	7	0	0	5	0
3	B	7	0	0	10	0
3	C	7	0	0	9	0
3	D	7	0	0	9	0
4	A	6	0	8	0	0
4	B	6	0	8	0	0
4	C	6	0	8	0	0
4	D	6	0	8	0	0
5	A	282	0	0	4	0
5	B	280	0	0	3	0
5	C	328	0	0	4	0
5	D	325	0	0	0	0
All	All	13336	0	12144	117	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 117 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:294:MET:HE2	1:A:388:GLY:HA2	1.32	1.10
1:B:65[A]:ARG:HE	3:C:502:O8Y:C5	1.71	1.02
1:A:65[A]:ARG:HE	3:D:502:O8Y:C5	1.75	0.98
1:A:163:MET:SD	1:A:294:MET:HE3	2.11	0.91
3:B:502:O8Y:C4	1:C:65[A]:ARG:HD2	2.00	0.91

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	403/423 (95%)	386 (96%)	16 (4%)	1 (0%)	43	36
1	B	399/423 (94%)	383 (96%)	14 (4%)	2 (0%)	24	16
1	C	395/423 (93%)	373 (94%)	20 (5%)	2 (0%)	24	16
1	D	406/423 (96%)	391 (96%)	14 (3%)	1 (0%)	43	36
All	All	1603/1692 (95%)	1533 (96%)	64 (4%)	6 (0%)	30	21

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	66	ASN
1	B	389	VAL
1	C	66	ASN
1	C	389	VAL
1	D	389	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	300/315 (95%)	291 (97%)	9 (3%)	36	27
1	B	300/315 (95%)	291 (97%)	9 (3%)	36	27
1	C	296/315 (94%)	286 (97%)	10 (3%)	32	23
1	D	303/315 (96%)	300 (99%)	3 (1%)	68	67
All	All	1199/1260 (95%)	1168 (97%)	31 (3%)	42	33

5 of 31 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	175	GLN
1	C	389	VAL
1	B	328	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	328	GLN
1	C	293	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	189	GLN
1	C	48	GLN
1	C	196	GLN
1	C	137	ASN
1	A	212	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	CSO	A	386	1	3,6,7	0.68	0	1,6,8	1.15	0
1	CSO	B	386[A]	1	3,6,7	0.80	0	1,6,8	1.06	0
1	CSO	D	386	1	3,6,7	0.79	0	1,6,8	0.38	0
1	CSO	C	386[B]	1	3,6,7	0.72	0	1,6,8	0.43	0
1	CSO	C	386[A]	1	3,6,7	0.71	0	1,6,8	1.20	0
1	CSO	B	386[B]	1	3,6,7	0.90	0	1,6,8	0.67	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSO	A	386	1	-	0/1/5/7	-
1	CSO	B	386[A]	1	-	0/1/5/7	-
1	CSO	D	386	1	-	0/1/5/7	-
1	CSO	C	386[B]	1	-	0/1/5/7	-
1	CSO	C	386[A]	1	-	0/1/5/7	-
1	CSO	B	386[B]	1	-	0/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	GOL	A	1003	-	5,5,5	0.11	0	5,5,5	0.42	0
3	O8Y	A	1002	-	5,6,6	0.71	0	4,5,5	0.31	0
2	COA	B	501	1	47,50,50	0.59	1 (2%)	69,75,75	0.71	2 (2%)
3	O8Y	D	502	-	5,6,6	0.68	0	4,5,5	0.52	0
3	O8Y	B	502	-	5,6,6	0.62	0	4,5,5	0.29	0
4	GOL	C	503	-	5,5,5	0.09	0	5,5,5	0.50	0
2	COA	A	1001	1	47,50,50	0.62	0	69,75,75	0.88	1 (1%)
4	GOL	B	503	-	5,5,5	0.12	0	5,5,5	0.32	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	O8Y	C	502	-	5,6,6	0.78	0	4,5,5	0.77	0
4	GOL	D	503	-	5,5,5	0.15	0	5,5,5	0.28	0
2	COA	D	501	1	47,50,50	0.65	1 (2%)	69,75,75	0.87	4 (5%)
2	COA	C	501	1	47,50,50	0.59	0	69,75,75	0.79	1 (1%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	1003	-	-	0/4/4/4	-
3	O8Y	A	1002	-	-	2/4/4/4	-
2	COA	B	501	1	-	5/48/64/64	0/3/3/3
3	O8Y	D	502	-	-	1/4/4/4	-
3	O8Y	B	502	-	-	2/4/4/4	-
4	GOL	C	503	-	-	0/4/4/4	-
2	COA	A	1001	1	-	0/48/64/64	0/3/3/3
4	GOL	B	503	-	-	0/4/4/4	-
3	O8Y	C	502	-	-	2/4/4/4	-
4	GOL	D	503	-	-	0/4/4/4	-
2	COA	D	501	1	-	4/48/64/64	0/3/3/3
2	COA	C	501	1	-	3/48/64/64	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	501	COA	P3B-O3B	2.75	1.64	1.59
2	B	501	COA	P3B-O3B	2.01	1.63	1.59

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1001	COA	P3B-O3B-C3B	-3.86	113.14	123.43
2	C	501	COA	P3B-O3B-C3B	-2.52	116.69	123.43
2	D	501	COA	O5A-P2A-O4A	2.32	123.21	112.44
2	B	501	COA	P3B-O3B-C3B	-2.26	117.38	123.43
2	D	501	COA	P3B-O3B-C3B	-2.26	117.41	123.43

There are no chirality outliers.

5 of 19 torsion outliers are listed below:




Mol	Chain	Res	Type	Atoms
2	B	501	COA	C5B-O5B-P1A-O2A
2	C	501	COA	C5B-O5B-P1A-O1A
2	C	501	COA	C5B-O5B-P1A-O3A
3	C	502	O8Y	C3-C4-C5-C6
3	D	502	O8Y	C3-C4-C5-C6


There are no ring outliers.




5 monomers are involved in 35 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1002	O8Y	5	0
3	D	502	O8Y	9	0
3	B	502	O8Y	10	0
3	C	502	O8Y	9	0
2	D	501	COA	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	402/423 (95%)	-0.16	1 (0%) 91 92	9, 20, 35, 71	3 (0%)
1	B	397/423 (93%)	-0.18	4 (1%) 79 84	8, 20, 37, 76	6 (1%)
1	C	395/423 (93%)	-0.29	0 100 100	7, 18, 32, 49	4 (1%)
1	D	401/423 (94%)	-0.32	0 100 100	7, 18, 29, 48	7 (1%)
All	All	1595/1692 (94%)	-0.24	5 (0%) 90 92	7, 19, 34, 76	20 (1%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	212	HIS	2.9
1	A	400	VAL	2.9
1	B	243	PRO	2.8
1	B	244	ASP	2.3
1	B	211	ALA	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	CSO	C	386[A]	7/8	0.96	0.07	16,16,21,24	7
1	CSO	C	386[B]	7/8	0.96	0.07	15,15,15,16	7
1	CSO	B	386[B]	7/8	0.98	0.05	20,20,21,24	7
1	CSO	A	386	7/8	0.98	0.07	19,19,21,33	0
1	CSO	B	386[A]	7/8	0.98	0.05	16,18,19,20	7
1	CSO	D	386	7/8	0.98	0.06	17,19,20,30	0

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	O8Y	A	1002	7/7	0.84	0.12	16,18,23,27	3
2	COA	D	501	48/48	0.87	0.10	22,30,57,69	1
3	O8Y	D	502	7/7	0.88	0.13	21,26,29,31	1
2	COA	B	501	48/48	0.89	0.09	27,39,64,77	1
2	COA	A	1001	48/48	0.89	0.09	22,28,62,76	1
3	O8Y	B	502	7/7	0.90	0.11	17,20,23,24	2
2	COA	C	501	48/48	0.90	0.09	22,32,52,57	1
4	GOL	A	1003	6/6	0.93	0.08	21,24,25,28	0
4	GOL	B	503	6/6	0.93	0.08	24,26,26,29	0
3	O8Y	C	502	7/7	0.95	0.09	17,20,22,25	2
4	GOL	C	503	6/6	0.95	0.07	16,18,19,21	0
4	GOL	D	503	6/6	0.95	0.07	18,19,20,22	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around O8Y A 1002:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around COA D 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around O8Y D 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around COA B 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around COA A 1001:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around O8Y B 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around COA C 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around GOL A 1003:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around GOL B 503:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around O8Y C 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around GOL C 503:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around GOL D 503:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.