



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 12:05 PM UTC

PDB ID : 3PEC / pdb_00003pec
Title : Siderocalin Recognitin of Carboxymycobactins: Interference by the immune system in intracellular iron acquisition by Mycobacteria tuberculosis
Authors : Clifton, M.C.
Deposited on : 2010-10-25
Resolution : 2.19 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	178	
1	B	178	
1	C	178	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	ZYG	C	188	X	-	-	-

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 4398 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Neutrophil gelatinase-associated lipocalin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	176	Total 1415	C 921	N 235	O 255	S 4	0	3	0
1	B	165	Total 1203	C 786	N 196	O 217	S 4	0	0	0
1	C	175	Total 1438	C 936	N 233	O 265	S 4	0	7	0

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	87	SER	CYS	engineered mutation	UNP P80188
B	87	SER	CYS	engineered mutation	UNP P80188
C	87	SER	CYS	engineered mutation	UNP P80188

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	C	1	Total C O 6 3 3	0	0
2	C	1	Total C O 6 3 3	0	0
2	C	1	Total C O 6 3 3	0	0

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0


- Molecule 4 is FE (III) ION (CCD ID: FE) (formula: Fe).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	1	Total Fe 1 1	0	0

- Molecule 5 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	C	4	Total Na 4 4	0	0

- Molecule 6 is 9-{hydroxy[(5S,6R)-6-hydroxy-6-[(1S)-3-[(3S)-1-hydroxy-2-oxoazepan-3-yl]amino}-1-methyl-3-oxopropyl]oxy}-5-([(4S)-2-(2-hydroxyphenyl)-4,5-dihydro-1,3-oxazol-4-yl]carbonyl)amino)hexyl]amino}-9-oxononanoic acid (CCD ID: ZYG) (formula: C₃₅H₅₃N₅O₁₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	C	1	Total C N O 52 35 12	0	0


- Molecule 7 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	87	Total O 87 87	0	0
7	B	20	Total O 20 20	0	0
7	C	121	Total O 121 121	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Neutrophil gelatinase-associated lipocalin

Chain A: 





- Molecule 1: Neutrophil gelatinase-associated lipocalin

Chain B: 



- Molecule 1: Neutrophil gelatinase-associated lipocalin

Chain C: 



4 Data and refinement statistics i

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	114.20Å 114.20Å 119.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.66 – 2.19 40.66 – 2.19	Depositor EDS
% Data completeness (in resolution range)	100.0 (40.66-2.19) 99.3 (40.66-2.19)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.04 (at 2.18Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.245 , 0.273 (Not available) , 0.281	Depositor DCC
R_{free} test set	3912 reflections (9.53%)	wwPDB-VP
Wilson B-factor (Å ²)	37.4	Xtriage
Anisotropy	0.024	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 47.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.011 for -h,l,k 0.006 for -l,-k,-h	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4398	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZYG, SO4, NA, GOL, FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/1461	0.61	0/1983
1	B	0.39	0/1235	0.63	0/1683
1	C	0.37	0/1493	0.60	0/2027
All	All	0.38	0/4189	0.61	0/5693

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1415	0	1387	18	0
1	B	1203	0	1034	9	0
1	C	1438	0	1407	16	0
2	A	24	0	32	0	0
2	C	18	0	24	0	0
3	A	5	0	0	0	0
3	C	10	0	0	0	0
4	C	1	0	0	0	0
5	C	4	0	0	0	0
6	C	52	0	49	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	87	0	0	0	0
7	B	20	0	0	0	0
7	C	121	0	0	0	0
All	All	4398	0	3933	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:34:VAL:HG21	1:C:144:LEU:HD22	1.52	0.91
1:B:153:ILE:HD13	1:B:166:ILE:HG21	1.61	0.82
1:A:34:VAL:HG21	1:A:144:LEU:HD12	1.67	0.74
1:A:34:VAL:HG23	1:A:137:LEU:HD13	1.76	0.67
1:A:140:ARG:HG2	1:C:23:GLN:NE2	2.12	0.64
1:C:120:MET:HE3	1:C:152:PHE:HD2	1.63	0.64
1:C:33:VAL:HG21	1:C:52:TYR:CZ	2.37	0.59
1:C:120:MET:HE1	1:C:155:PHE:CD2	2.39	0.57
1:C:120:MET:HE1	1:C:155:PHE:HD2	1.72	0.55
1:B:122:PHE:HD2	1:B:135:ILE:HD13	1.72	0.55
1:A:33:VAL:HG21	1:A:52:TYR:CE2	2.44	0.53
1:A:70:LEU:HD21	1:A:79:TRP:CZ2	2.44	0.53
1:C:34:VAL:HG23	1:C:137:LEU:HD13	1.90	0.52
1:C:33:VAL:HG21	1:C:52:TYR:CE2	2.44	0.52
1:C:120:MET:HE3	1:C:152:PHE:CD2	2.44	0.51
1:A:13:LEU:O	1:A:109:ARG:NH2	2.44	0.51
1:A:36:LEU:HD22	1:A:41:ILE:HG13	1.94	0.50
1:B:153:ILE:CD1	1:B:166:ILE:HG21	2.38	0.49
1:B:135:ILE:HD11	1:B:159:LEU:CD1	2.43	0.49
1:A:81:ARG:HG3	1:A:94:LEU:HD11	1.94	0.49
1:B:143:GLU:CB	1:B:144:LEU:CB	2.91	0.48
1:A:69:VAL:C	1:A:70:LEU:HD23	2.39	0.48
1:A:34:VAL:CG2	1:A:137:LEU:HD13	2.44	0.47
1:A:81:ARG:CG	1:A:94:LEU:HD11	2.45	0.47
1:C:143:GLU:O	1:C:144:LEU:HD12	2.15	0.46
1:B:108:VAL:HG22	1:B:123:PHE:CD1	2.51	0.46
1:A:70:LEU:HD21	1:A:79:TRP:CE2	2.51	0.45
1:C:31:TRP:CE3	1:C:138:TYR:HB3	2.51	0.45
1:C:119:ALA:HB3	1:C:138:TYR:HB2	1.99	0.44
1:B:25:ASN:HA	1:B:58:LEU:HD12	1.99	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:119:ALA:HB3	1:A:138:TYR:HB2	1.99	0.43
1:B:119:ALA:HB3	1:B:138:TYR:HB2	2.00	0.43
1:C:34:VAL:CG2	1:C:144:LEU:HD22	2.36	0.42
1:A:31:TRP:CE3	1:A:138:TYR:HB3	2.54	0.42
1:A:70:LEU:HD21	1:A:79:TRP:NE1	2.34	0.42
1:B:18:LEU:HD11	1:B:90:GLY:HA3	2.01	0.42
1:C:120:MET:CE	1:C:155:PHE:CD2	3.03	0.42
1:A:34:VAL:HG21	1:A:144:LEU:CD1	2.43	0.41
1:A:33:VAL:HG21	1:A:52:TYR:CZ	2.56	0.40
1:C:52:TYR:HB3	1:C:70:LEU:HG	2.03	0.40
1:A:66:VAL:HG21	1:A:83:PHE:CD1	2.57	0.40
1:C:49:GLN:HB3	1:C:167:VAL:HG11	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	177/178 (99%)	174 (98%)	3 (2%)	0	100	100
1	B	159/178 (89%)	152 (96%)	6 (4%)	1 (1%)	21	23
1	C	180/178 (101%)	176 (98%)	3 (2%)	1 (1%)	21	23
All	All	516/534 (97%)	502 (97%)	12 (2%)	2 (0%)	30	34

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	144	LEU
1	C	6	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	151/163 (93%)	144 (95%)	7 (5%)	24	32
1	B	104/163 (64%)	101 (97%)	3 (3%)	37	51
1	C	156/163 (96%)	150 (96%)	6 (4%)	29	40
All	All	411/489 (84%)	395 (96%)	16 (4%)	28	39

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	36	LEU
1	A	70	LEU
1	A	79	TRP
1	A	81	ARG
1	A	137	LEU
1	A	140	ARG
1	A	163	GLU
1	B	58	LEU
1	B	109	ARG
1	B	140	ARG
1	C	36	LEU
1	C	58	LEU
1	C	70	LEU
1	C	134	LYS
1	C	137	LEU
1	C	140	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	49	GLN
1	B	26	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 5 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	C	181	-	5,5,5	0.41	0	5,5,5	0.33	0
2	GOL	C	3731	-	5,5,5	0.38	0	5,5,5	0.23	0
2	GOL	C	180	-	5,5,5	0.38	0	5,5,5	0.22	0
2	GOL	A	182	-	5,5,5	0.39	0	5,5,5	0.23	0
3	SO4	C	186	-	4,4,4	0.24	0	6,6,6	0.09	0
2	GOL	A	179	-	5,5,5	0.38	0	5,5,5	0.29	0
3	SO4	A	183	-	4,4,4	0.24	0	6,6,6	0.09	0
2	GOL	A	181	-	5,5,5	0.38	0	5,5,5	0.09	0
6	ZYG	C	188	4	51,54,54	4.46	13 (25%)	54,71,71	4.48	23 (42%)
3	SO4	C	187	-	4,4,4	0.24	0	6,6,6	0.07	0
2	GOL	A	180	-	5,5,5	0.39	0	5,5,5	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	C	181	-	-	0/4/4/4	-
2	GOL	C	3731	-	-	2/4/4/4	-
2	GOL	C	180	-	-	2/4/4/4	-
2	GOL	A	182	-	-	4/4/4/4	-
2	GOL	A	179	-	-	1/4/4/4	-
6	ZYG	C	188	4	1/1/13/15	19/49/73/73	1/3/3/3
2	GOL	A	181	-	-	2/4/4/4	-
2	GOL	A	180	-	-	2/4/4/4	-

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	C	188	ZYG	C10-N11	12.18	1.44	1.27
6	C	188	ZYG	O38-C37	12.17	1.44	1.22
6	C	188	ZYG	C31-C30	-11.67	1.35	1.53
6	C	188	ZYG	O16-C15	10.55	1.43	1.23
6	C	188	ZYG	O28-C27	10.40	1.43	1.23
6	C	188	ZYG	O46-C45	9.77	1.44	1.23
6	C	188	ZYG	C45-N43	8.68	1.46	1.35
6	C	188	ZYG	O36-N35	7.09	1.45	1.40
6	C	188	ZYG	C15-N17	5.93	1.46	1.34
6	C	188	ZYG	C27-N29	5.54	1.45	1.34
6	C	188	ZYG	C30-N29	-5.18	1.35	1.45
6	C	188	ZYG	O14-C10	4.94	1.44	1.36
6	C	188	ZYG	C12-N11	-2.34	1.44	1.47

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	C	188	ZYG	O14-C10-N11	-14.43	101.49	118.10
6	C	188	ZYG	O38-C37-N35	-13.18	104.03	122.29
6	C	188	ZYG	C31-C30-C37	11.22	125.97	111.65
6	C	188	ZYG	O16-C15-C12	-9.50	107.12	120.19
6	C	188	ZYG	O28-C27-N29	-8.63	108.34	122.95
6	C	188	ZYG	O28-C27-C25	-8.44	109.20	121.54
6	C	188	ZYG	O16-C15-N17	-6.61	111.13	122.96
6	C	188	ZYG	O38-C37-C30	-6.25	108.66	121.04
6	C	188	ZYG	C8-C10-N11	-6.18	112.37	126.00
6	C	188	ZYG	C25-C27-N29	-5.62	108.58	116.25
6	C	188	ZYG	C32-C31-C30	5.42	124.79	114.23
6	C	188	ZYG	C37-C30-N29	5.17	119.64	108.73
6	C	188	ZYG	O46-C45-C47	-4.68	108.59	121.26

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	C	188	ZYG	C12-C15-N17	-4.43	109.94	115.71
6	C	188	ZYG	C13-O14-C10	3.54	109.56	105.54
6	C	188	ZYG	C13-C12-N11	-3.13	99.98	103.64
6	C	188	ZYG	O44-N43-C42	3.05	120.90	113.76
6	C	188	ZYG	C41-C42-N43	-2.73	105.50	111.11
6	C	188	ZYG	C31-C30-N29	2.52	114.38	109.76
6	C	188	ZYG	C33-C32-C31	-2.49	108.38	115.77
6	C	188	ZYG	C30-N29-C27	2.37	127.61	121.68
6	C	188	ZYG	C39-C18-N17	2.21	113.16	110.30
6	C	188	ZYG	C3-C8-C10	-2.06	119.00	121.85

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
6	C	188	ZYG	C30

All (32) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	180	GOL	C1-C2-C3-O3
2	A	182	GOL	C1-C2-C3-O3
6	C	188	ZYG	N11-C10-C8-C3
6	C	188	ZYG	O16-C15-N17-C18
6	C	188	ZYG	C39-C18-N17-C15
6	C	188	ZYG	C25-C27-N29-C30
6	C	188	ZYG	O28-C27-N29-C30
6	C	188	ZYG	C37-C30-N29-C27
6	C	188	ZYG	C41-C42-N43-O44
6	C	188	ZYG	C41-C42-N43-C45
6	C	188	ZYG	C47-C45-N43-C42
6	C	188	ZYG	C47-C45-N43-O44
6	C	188	ZYG	N43-C45-C47-C48
2	C	3731	GOL	O1-C1-C2-O2
2	A	181	GOL	O1-C1-C2-C3
2	C	180	GOL	O1-C1-C2-C3
2	C	3731	GOL	O1-C1-C2-C3
2	A	181	GOL	O1-C1-C2-O2
2	A	182	GOL	O2-C2-C3-O3
6	C	188	ZYG	C48-C49-C50-C51
2	A	180	GOL	O2-C2-C3-O3
6	C	188	ZYG	N11-C10-C8-C7
2	A	179	GOL	O2-C2-C3-O3

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	C	180	GOL	O1-C1-C2-O2
2	A	182	GOL	O1-C1-C2-C3
6	C	188	ZYG	C13-C12-C15-N17
6	C	188	ZYG	C52-C53-C54-O51
6	C	188	ZYG	C52-C53-C54-O52
2	A	182	GOL	O1-C1-C2-O2
6	C	188	ZYG	N11-C12-C15-N17
6	C	188	ZYG	C56-C22-C25-C27
6	C	188	ZYG	O46-C45-N43-O44

All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	C	188	ZYG	C30-C31-C32-C33-C34-C37-N35

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	176/178 (98%)	0.70	13 (7%) 20 18	22, 35, 50, 60	3 (1%)
1	B	165/178 (92%)	3.33	139 (84%) 0 0	51, 73, 104, 111	0
1	C	175/178 (98%)	0.49	15 (8%) 16 14	12, 26, 44, 60	7 (4%)
All	All	516/534 (96%)	1.47	167 (32%) 1 0	12, 37, 93, 111	10 (1%)

All (167) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	144	LEU	7.9
1	B	160	GLY	6.8
1	B	167	VAL	6.4
1	A	79	TRP	6.2
1	C	178	GLY	6.0
1	B	71	PHE	5.8
1	B	75	LYS	5.8
1	B	146	SER	5.7
1	B	7	LEU	5.7
1	B	166	ILE	5.6
1	B	63	SER	5.6
1	B	176	ILE	5.5
1	B	60	GLU	5.3
1	B	159	LEU	5.3
1	B	163	GLU	5.3
1	B	42	LEU	5.3
1	B	173	ASP	5.2
1	B	89	PRO	5.2
1	B	149	LYS	5.1
1	B	161	LEU	5.1
1	A	3	SER	5.1
1	B	52	TYR	5.0
1	B	74	LYS	5.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	172	ILE	5.0
1	B	155	PHE	4.9
1	B	80	ILE	4.9
1	B	87	SER	4.9
1	B	170	VAL	4.9
1	B	116	ASN	4.8
1	B	162	PRO	4.8
1	B	8	ILE	4.8
1	B	43	ARG	4.7
1	C	79	TRP	4.6
1	B	151	ASN	4.6
1	B	165	HIS	4.6
1	B	83	PHE	4.6
1	B	145	THR	4.5
1	B	110	VAL	4.5
1	B	92	PHE	4.4
1	B	168	PHE	4.4
1	B	58	LEU	4.3
1	B	152	PHE	4.3
1	B	97	ILE	4.3
1	B	69	VAL	4.2
1	B	37	ALA	4.2
1	B	34	VAL	4.2
1	B	56	TYR	4.1
1	B	153	ILE	4.1
1	B	157	LYS	4.1
1	B	55	ILE	4.1
1	B	53	ALA	4.0
1	B	36	LEU	4.0
1	B	107	LEU	4.0
1	B	94	LEU	4.0
1	B	164	ASN	4.0
1	B	13	LEU	3.9
1	B	70	LEU	3.9
1	B	95	GLY	3.8
1	C	4	THR	3.8
1	B	135	ILE	3.8
1	B	31	TRP	3.7
1	B	73	LYS	3.7
1	B	78	TYR	3.7
1	B	88	GLN	3.7
1	B	138	TYR	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	66	VAL	3.6
1	B	90	GLY	3.6
1	B	50	LYS	3.6
1	C	47	ASP	3.6
1	B	54	THR	3.6
1	B	81	ARG	3.6
1	B	67	THR	3.5
1	B	115	TYR	3.5
1	B	33	VAL	3.5
1	B	171	PRO	3.5
1	B	137	LEU	3.5
1	B	158	SER	3.5
1	B	84	VAL	3.5
1	B	133	PHE	3.4
1	B	76	CYS	3.4
1	B	79	TRP	3.4
1	B	174	GLN	3.4
1	B	96	ASN	3.3
1	B	114	ASN	3.3
1	B	64	TYR	3.2
1	B	132	TYR	3.2
1	B	28	GLN	3.2
1	B	41	ILE	3.2
1	B	150	GLU	3.2
1	B	129	ASN	3.2
1	B	103	LEU	3.2
1	B	40	ALA	3.1
1	B	59	LYS	3.1
1	B	169	PRO	3.1
1	B	68	SER	3.1
1	B	93	THR	3.1
1	B	140	ARG	3.0
1	B	30	LYS	3.0
1	B	98	LYS	3.0
1	B	136	THR	3.0
1	B	123	PHE	3.0
1	B	6	ASP	2.9
1	B	51	MET	2.9
1	B	82	THR	2.9
1	B	177	ASP	2.8
1	B	85	PRO	2.8
1	B	32	TYR	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	61	ASP	2.8
1	A	4	THR	2.8
1	B	39	ASN	2.8
1	B	9	PRO	2.8
1	C	44	GLU	2.8
1	C	101	PRO	2.8
1	B	117	GLN	2.7
1	B	77	ASP	2.7
1	B	86	GLY	2.7
1	B	156	SER	2.7
1	B	102	GLY	2.7
1	B	99	SER	2.7
1	B	23	GLN	2.6
1	B	22	PHE	2.6
1	B	18	LEU	2.6
1	A	45	ASP	2.6
1	B	111	VAL	2.6
1	B	57	GLU	2.6
1	C	48	PRO	2.6
1	B	62	LYS	2.6
1	C	70	LEU	2.6
1	C	143	GLU	2.6
1	B	128	GLN	2.5
1	B	143	GLU	2.5
1	B	20	GLN	2.5
1	B	104	THR	2.5
1	B	105	SER	2.5
1	B	141	THR	2.5
1	B	21	ASN	2.5
1	A	47	ASP	2.4
1	B	175	CYS	2.4
1	B	38	GLY	2.4
1	C	151	ASN	2.4
1	C	150	GLU	2.3
1	A	5	SER	2.3
1	C	45	ASP	2.3
1	B	108	VAL	2.3
1	B	122	PHE	2.3
1	A	20	GLN	2.3
1	B	65	ASN	2.3
1	B	25	ASN	2.2
1	B	154	ARG	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	100	TYR	2.2
1	B	121	VAL	2.2
1	B	126	VAL	2.2
1	B	49	GLN	2.2
1	B	17	PRO	2.2
1	C	163	GLU	2.2
1	C	100	TYR	2.2
1	A	46	LYS	2.2
1	A	128	GLN	2.1
1	C	81	ARG	2.1
1	A	99	SER	2.1
1	B	27	PHE	2.1
1	A	151	ASN	2.1
1	A	81	ARG	2.1
1	A	130	ARG	2.1
1	B	142	LYS	2.1
1	B	112	SER	2.1
1	B	131	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	SO4	A	183	5/5	0.49	0.18	97,97,97,97	0
5	NA	C	182	1/1	0.51	0.27	49,49,49,49	0
2	GOL	A	181	6/6	0.52	0.29	69,69,69,69	0
5	NA	C	185	1/1	0.59	0.31	55,55,55,55	0
2	GOL	C	3731	6/6	0.62	0.21	48,49,49,49	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GOL	A	182	6/6	0.67	0.20	60,60,60,61	0
2	GOL	C	180	6/6	0.70	0.21	56,57,57,57	0
6	ZYG	C	188	52/52	0.72	0.20	44,54,62,62	0
2	GOL	A	180	6/6	0.74	0.18	64,64,64,64	0
2	GOL	A	179	6/6	0.75	0.22	70,70,70,71	0
3	SO4	C	187	5/5	0.76	0.12	86,86,86,86	0
2	GOL	C	181	6/6	0.83	0.17	45,45,46,46	0
3	SO4	C	186	5/5	0.83	0.12	83,83,83,84	0
5	NA	C	183	1/1	0.89	0.12	42,42,42,42	0
5	NA	C	184	1/1	0.92	0.13	32,32,32,32	0
4	FE	C	179	1/1	0.95	0.06	55,55,55,55	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.