



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 06:58 AM UTC

PDB ID : 6PEC / pdb_00006pec
Title : Vaccine-elicited NHP FP-targeting antibody DF2F-e.01 in complex with HIV fusion peptide (residue 512-519)
Authors : Xu, K.; Liu, K.; Wang, Y.; Kwong, P.D.
Deposited on : 2019-06-20
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1187 (1.74-1.74)
Clashscore	190562	1207 (1.74-1.74)
Ramachandran outliers	187476	1200 (1.74-1.74)
Sidechain outliers	187428	1200 (1.74-1.74)
RSRZ outliers	180081	1188 (1.74-1.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	208	22% 88% 11%
2	H	216	14% 87% 12%
3	A	8	88% 12%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3693 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called antibody light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	208	Total	C	N	O	S	0	0	0
			1592	1001	261	325	5			

- Molecule 2 is a protein called antibody heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	216	Total	C	N	O	S	0	0	0
			1607	1016	267	320	4			

- Molecule 3 is a protein called HIV-1 fusion peptide residue 512-519.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	A	8	Total	C	N	O	0	0	0
			51	35	8	8			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	L	202	Total	O	0	0
			202	202		
4	H	230	Total	O	0	0
			230	230		
4	A	11	Total	O	0	0
			11	11		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: antibody light chain



- Molecule 2: antibody heavy chain



- Molecule 3: HIV-1 fusion peptide residue 512-519



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	54.76Å 88.38Å 121.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.71 – 1.75 35.71 – 1.75	Depositor EDS
% Data completeness (in resolution range)	98.3 (35.71-1.75) 98.4 (35.71-1.75)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.50 (at 1.75Å)	Xtriage
Refinement program	PHENIX 1.8.4_1496	Depositor
R, R_{free}	0.194 , 0.215 0.196 , 0.215	Depositor DCC
R_{free} test set	2987 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	26.6	Xtriage
Anisotropy	0.543	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 38.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3693	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	L	0.55	0/1627	0.91	3/2212 (0.1%)
2	H	0.52	0/1647	0.86	2/2247 (0.1%)
3	A	0.54	0/51	0.77	0/68
All	All	0.53	0/3325	0.89	5/4527 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	172	SER	CA-C-N	8.02	131.02	120.28
2	H	172	SER	C-N-CA	8.02	131.02	120.28
1	L	188	LYS	N-CA-C	-7.46	104.18	113.28
1	L	126	LYS	CA-C-N	6.24	128.64	120.28
1	L	126	LYS	C-N-CA	6.24	128.64	120.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1592	0	1544	20	0
2	H	1607	0	1578	18	0
3	A	51	0	53	1	0
4	A	11	0	0	0	0
4	H	230	0	0	4	2

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	L	202	0	0	2	2
All	All	3693	0	3175	38	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:128:GLY:HA2	1:L:183:LYS:HG2	1.37	1.06
1:L:128:GLY:HA2	1:L:183:LYS:CG	2.06	0.84
2:H:148:GLU:HG3	2:H:176:TYR:CE1	2.28	0.69
1:L:145:LYS:NZ	1:L:147:GLN:HE21	1.93	0.67
1:L:128:GLY:O	1:L:183:LYS:HG3	1.95	0.67
1:L:4:MET:HE1	1:L:90:GLN:HB2	1.81	0.63
1:L:186:TYR:O	1:L:192:TYR:OH	2.09	0.61
1:L:124:GLN:HE22	1:L:131:SER:HB2	1.67	0.59
2:H:148:GLU:HG3	2:H:176:TYR:CZ	2.38	0.59
2:H:22:CYS:HB3	2:H:78:PHE:HB3	1.85	0.58
2:H:120:SER:OG	4:H:301:HOH:O	2.07	0.57
1:L:210:ASN:O	1:L:211:ARG:HG3	2.06	0.56
1:L:145:LYS:HZ1	1:L:147:GLN:HE21	1.52	0.55
2:H:148:GLU:OE2	2:H:168:ALA:HB3	2.08	0.54
2:H:126:PRO:HD3	2:H:138:LEU:HD23	1.90	0.54
2:H:170:LEU:N	2:H:170:LEU:CD2	2.71	0.54
2:H:34:TRP:HZ2	2:H:94:ARG:HH21	1.58	0.52
2:H:77:GLN:OE1	4:H:302:HOH:O	2.18	0.51
2:H:94:ARG:NH2	4:H:312:HOH:O	2.43	0.50
1:L:128:GLY:C	1:L:183:LYS:HG3	2.36	0.50
2:H:40:PRO:HB2	2:H:43:LYS:HD3	1.94	0.50
1:L:151:ASP:OD1	1:L:190:LYS:HG2	2.11	0.49
2:H:206:LYS:HE2	4:H:331:HOH:O	2.10	0.49
2:H:100:TRP:CH2	2:H:100(A):LYS:HE2	2.50	0.47
1:L:128:GLY:CA	1:L:183:LYS:CG	2.88	0.47
1:L:208:SER:OG	4:L:301:HOH:O	1.98	0.46
2:H:47:TRP:CZ2	2:H:49:GLY:HA2	2.51	0.46
1:L:11:LEU:HD11	1:L:19:VAL:HG13	1.98	0.45
1:L:31:SER:HB2	4:L:373:HOH:O	2.15	0.45
1:L:94:TYR:CZ	3:A:513:VAL:HB	2.52	0.44
1:L:158:ASN:O	1:L:179:LEU:HD12	2.18	0.44
1:L:147:GLN:HB2	1:L:195:GLU:HG2	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:28:THR:HA	2:H:76:ASN:HD21	1.82	0.43
1:L:183:LYS:HG3	1:L:183:LYS:H	1.58	0.42
2:H:6:GLU:HG3	2:H:92:CYS:SG	2.60	0.42
2:H:170:LEU:N	2:H:170:LEU:HD23	2.36	0.41
2:H:122:PHE:HA	2:H:123:PRO:HD3	1.94	0.40
1:L:124:GLN:NE2	1:L:131:SER:H	2.18	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:L:460:HOH:O	4:H:401:HOH:O[1_455]	1.89	0.31
4:L:481:HOH:O	4:H:502:HOH:O[1_455]	2.05	0.15

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	206/208 (99%)	202 (98%)	4 (2%)	0	100	100
2	H	212/216 (98%)	211 (100%)	1 (0%)	0	100	100
3	A	6/8 (75%)	6 (100%)	0	0	100	100
All	All	424/432 (98%)	419 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	182/183 (100%)	177 (97%)	5 (3%)	39	16
2	H	183/183 (100%)	180 (98%)	3 (2%)	55	35
3	A	4/4 (100%)	4 (100%)	0	100	100
All	All	369/370 (100%)	361 (98%)	8 (2%)	45	22

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	4	MET
1	L	149	LYS
1	L	183	LYS
1	L	190	LYS
1	L	201	LEU
2	H	7	SER
2	H	170	LEU
2	H	189	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	37	GLN
1	L	89	GLN
1	L	124	GLN
1	L	147	GLN
2	H	192	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	H	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	H	127:SER	C	134:GLY	N	12.38

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	L	208/208 (100%)	1.04	45 (21%) 2 3	20, 34, 72, 80	0
2	H	216/216 (100%)	0.75	31 (14%) 6 8	19, 32, 54, 65	0
3	A	8/8 (100%)	0.13	0 100 100	22, 23, 28, 29	0
All	All	432/432 (100%)	0.88	76 (17%) 4 5	19, 32, 67, 80	0

All (76) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	125	LEU	6.8
2	H	170	LEU	5.9
1	L	186	TYR	5.5
1	L	184	ALA	5.1
1	L	191	VAL	4.8
1	L	185	ASP	4.5
2	H	171	GLN	4.3
1	L	130	ALA	4.2
1	L	121	SER	4.0
1	L	189	HIS	4.0
1	L	181	LEU	4.0
1	L	152	ASN	3.9
1	L	153	ALA	3.9
2	H	9	PRO	3.8
2	H	169	VAL	3.8
2	H	191	THR	3.7
1	L	211	ARG	3.5
2	H	213	PRO	3.4
2	H	176	TYR	3.4
1	L	187	GLU	3.4
1	L	192	TYR	3.4
2	H	134	GLY	3.3
1	L	154	LEU	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	L	210	ASN	3.3
1	L	188	LYS	3.2
1	L	190	LYS	3.2
2	H	127	SER	3.1
1	L	195	GLU	3.1
2	H	105	GLN	3.0
1	L	150	VAL	2.9
1	L	209	PHE	2.9
1	L	183	LYS	2.9
2	H	168	ALA	2.9
1	L	194	CYS	2.9
2	H	190	GLY	2.9
2	H	172	SER	2.9
2	H	6	GLU	2.8
1	L	119	PRO	2.7
2	H	189	LEU	2.7
1	L	128	GLY	2.7
1	L	155	GLN	2.7
2	H	173	SER	2.7
1	L	179	LEU	2.7
1	L	193	ALA	2.6
2	H	174	GLY	2.5
2	H	212	GLU	2.5
1	L	127	SER	2.5
1	L	151	ASP	2.5
2	H	42	GLY	2.5
1	L	148	TRP	2.4
1	L	145	LYS	2.4
1	L	180	THR	2.4
2	H	24	VAL	2.4
1	L	120	PRO	2.3
1	L	201	LEU	2.3
1	L	182	SER	2.3
1	L	202	SER	2.3
2	H	25	SER	2.3
2	H	5	GLN	2.2
1	L	129	THR	2.2
1	L	149	LYS	2.2
2	H	144	ASP	2.2
1	L	147	GLN	2.1
2	H	140	CYS	2.1
1	L	67	SER	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	L	156	SER	2.1
2	H	41	PRO	2.1
2	H	73	THR	2.1
2	H	195	ILE	2.1
1	L	134	CYS	2.1
2	H	148	GLU	2.1
1	L	157	GLY	2.1
2	H	158	ALA	2.1
2	H	188	SER	2.1
2	H	175	LEU	2.0
1	L	23	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.