



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 01:28 AM UTC

PDB ID : 6PEH / pdb_00006peh
Title : Crystal structure of rabbit monoclonal anti-HIV antibody 1C2
Authors : Liban, T.; Pancera, M.
Deposited on : 2019-06-20
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	237	10% 89% 6% .
1	H	237	8% 92% 7% .
2	B	215	16% 93% 6% .
2	L	215	14% 90% 9% .

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 13001 atoms, of which 6247 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 1C2 Fab Heavy Chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	H	220	Total	C	H	N	O	S	0	10	0
			3220	1032	1585	269	322	12			
1	A	227	Total	C	H	N	O	S	0	7	0
			3295	1053	1619	278	332	13			

- Molecule 2 is a protein called 1C2 Fab Light Chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	L	213	Total	C	H	N	O	S	0	3	0
			3115	995	1528	256	328	8			
2	B	214	Total	C	H	N	O	S	0	2	0
			3093	989	1515	257	325	7			

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	L	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	H	93	Total	O	0	0
			93	93		
4	L	58	Total	O	0	0
			58	58		
4	A	78	Total	O	0	0
			78	78		
4	B	39	Total	O	0	0
			39	39		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 1C2 Fab Heavy Chain




- Molecule 1: 1C2 Fab Heavy Chain



- Molecule 2: 1C2 Fab Light Chain



- Molecule 2: 1C2 Fab Light Chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, α , β , γ	71.80Å 98.61Å 162.98Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.32 – 2.30 43.32 – 2.30	Depositor EDS
% Data completeness (in resolution range)	81.9 (43.32-2.30) 81.9 (43.32-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.04 (at 2.29Å)	Xtriage
Refinement program	PHENIX (1.12_2829: ???)	Depositor
R, R_{free}	0.219 , 0.255 0.223 , 0.256	Depositor DCC
R_{free} test set	2129 reflections (4.05%)	wwPDB-VP
Wilson B-factor (Å ²)	31.9	Xtriage
Anisotropy	0.064	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 44.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	13001	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, PCA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.16	0/1727	0.35	0/2370
1	H	0.17	0/1701	0.36	0/2330
2	B	0.16	0/1623	0.38	0/2221
2	L	0.17	0/1636	0.37	0/2236
All	All	0.16	0/6687	0.36	0/9157

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1676	1619	1596	5	0
1	H	1635	1585	1548	1	0
2	B	1578	1515	1501	9	0
2	L	1587	1528	1507	9	1
3	A	5	0	0	0	1
3	L	5	0	0	0	0
4	A	78	0	0	0	1
4	B	39	0	0	2	1
4	H	93	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	L	58	0	0	1	0
All	All	6754	6247	6152	23	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:4:MET:HE3	2:B:90:TYR:HB3	1.71	0.72
1:A:164:ARG:NH2	2:B:137:ASN:OD1	2.31	0.60
1:A:48:ILE:HG23	1:A:67:LEU:HD13	1.84	0.60
1:A:66:ARG:NH1	1:A:86:ASP:OD2	2.38	0.57
2:L:125:VAL:O	2:L:182:SER:OG	2.26	0.54
2:L:130:VAL:HG11	2:L:185:TYR:HB2	1.88	0.54
2:L:131:THR:HG23	2:L:177:THR:CG2	2.40	0.51
2:B:39:LYS:NZ	2:B:81:ASP:O	2.27	0.49
2:L:187:SER:OG	2:L:188:HIS:ND1	2.46	0.48
1:H:211:SER:N	4:H:306:HOH:O	2.46	0.46
2:L:13:ALA:HB2	2:L:19:VAL:HG13	1.97	0.46
2:L:33:LEU:HD22	2:L:71:PHE:CG	2.51	0.46
2:B:138:LYS:O	4:B:302:HOH:O	2.21	0.45
2:B:119:PRO:O	4:B:301:HOH:O	2.21	0.45
2:L:80:CYS:SG	2:L:170[B]:CYS:HB3	2.57	0.45
2:L:130:VAL:HG11	2:L:185:TYR:CB	2.47	0.44
2:B:79:GLN:OE1	2:B:80:CYS:N	2.51	0.44
2:B:131:THR:HG23	2:B:177:THR:CG2	2.49	0.42
1:A:128:CYS:N	1:A:212:THR:OG1	2.53	0.42
2:B:50:ASP:O	2:B:51:THR:HB	2.20	0.41
2:B:13:ALA:HB2	2:B:19:VAL:HG22	2.03	0.41
1:A:52:TYR:CZ	1:A:52(B):GLY:HA3	2.56	0.41
2:L:45:LYS:NZ	4:L:401:HOH:O	2.36	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:199:THR:O	4:B:336:HOH:O[3_545]	2.11	0.09
3:A:301:SO4:O2	4:A:437:HOH:O[3_455]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	232/237 (98%)	226 (97%)	6 (3%)	0	100	100
1	H	224/237 (94%)	218 (97%)	6 (3%)	0	100	100
2	B	214/215 (100%)	200 (94%)	13 (6%)	1 (0%)	24	31
2	L	214/215 (100%)	198 (92%)	15 (7%)	1 (0%)	24	31
All	All	884/904 (98%)	842 (95%)	40 (4%)	2 (0%)	43	55

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	L	138	LYS
2	B	138	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	187/195 (96%)	181 (97%)	6 (3%)	34	51
1	H	184/195 (94%)	183 (100%)	1 (0%)	81	90
2	B	177/181 (98%)	175 (99%)	2 (1%)	65	81
2	L	181/181 (100%)	179 (99%)	2 (1%)	65	81
All	All	729/752 (97%)	718 (98%)	11 (2%)	59	75

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	75	THR
2	L	69	THR
2	L	96	PHE
1	A	3	SER
1	A	75	THR
1	A	130	ASP
1	A	136	VAL
1	A	193[A]	CYS
1	A	193[B]	CYS
2	B	137	ASN
2	B	138	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	H	80	GLN
1	A	32	ASN
2	B	22	ASN
2	B	207	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	PCA	A	2	1	7,8,9	2.00	1 (14%)	9,10,12	1.94	4 (44%)
1	PCA	H	2	1	7,8,9	1.85	1 (14%)	9,10,12	1.50	3 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PCA	A	2	1	-	0/0/11/13	0/1/1/1
1	PCA	H	2	1	-	0/0/11/13	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	2	PCA	CD-N	5.18	1.47	1.34
1	H	2	PCA	CD-N	4.76	1.46	1.34

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	2	PCA	O-C-CA	-3.10	116.80	124.77
1	A	2	PCA	OE-CD-CG	-2.73	121.85	126.72
1	A	2	PCA	CA-N-CD	-2.48	105.08	113.58
1	H	2	PCA	CB-CA-N	2.22	109.36	103.24
1	A	2	PCA	CB-CG-CD	-2.07	101.22	104.41
1	H	2	PCA	CA-N-CD	-2.02	106.67	113.58
1	H	2	PCA	OE-CD-CG	-2.01	123.14	126.72

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	301	-	4,4,4	0.24	0	6,6,6	0.07	0
3	SO4	L	301	-	4,4,4	0.23	0	6,6,6	0.10	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	301	SO4	0	1

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	226/237 (95%)	0.56	23 (10%) 12 13	11, 39, 72, 123	4 (1%)
1	H	219/237 (92%)	0.36	18 (8%) 17 19	8, 33, 83, 117	5 (2%)
2	B	214/215 (99%)	1.13	35 (16%) 4 5	26, 52, 86, 120	1 (0%)
2	L	213/215 (99%)	0.83	31 (14%) 6 6	18, 49, 76, 98	1 (0%)
All	All	872/904 (96%)	0.71	107 (12%) 8 9	8, 43, 82, 123	11 (1%)

All (107) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	212	THR	5.7
1	A	64	LYS	5.2
1	H	211	SER	5.2
2	B	1	ALA	5.1
1	A	63	ALA	5.0
2	L	210	ASP	4.9
1	A	130	ASP	4.7
2	B	168	ALA	4.7
1	A	128	CYS	4.6
1	A	213	CYS	4.5
1	H	132	PRO	4.4
2	B	155	THR	4.4
1	H	130	ASP	4.4
1	A	131	THR	4.4
1	H	129	GLY	4.3
1	A	132	PRO	4.1
2	B	12	SER	4.0
2	L	76	SER	4.0
2	B	95	SER	4.0
1	H	128	CYS	3.9
2	B	56	SER	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	L	62	PHE	3.7
1	A	211	SER	3.7
1	H	190	PRO	3.7
1	H	131	THR	3.6
1	H	160	THR	3.6
2	B	156	THR	3.6
1	A	129[A]	GLY	3.5
1	H	134	SER	3.5
2	B	167	SER	3.5
2	B	106	VAL	3.4
1	H	30	GLY	3.4
2	B	27	GLU	3.3
2	B	201	SER	3.3
2	B	13	ALA	3.2
1	A	212	THR	3.1
2	L	64	GLY	3.1
1	A	61	ASN	3.0
1	H	127	CYS	3.0
2	L	18	THR	3.0
2	L	203	VAL	3.0
2	L	202	VAL	3.0
1	H	100	ASP	2.9
2	L	58	VAL	2.8
1	H	161	ASN	2.8
1	H	186	SER	2.8
2	B	7	THR	2.8
2	B	144	THR	2.7
2	B	105	VAL	2.7
2	L	169	ASP	2.7
1	A	108	LEU	2.7
2	B	169	ASP	2.6
1	A	133	SER	2.6
1	A	134	SER	2.6
2	B	78	VAL	2.6
1	H	171	GLN	2.6
2	L	198	GLY	2.6
2	B	200	THR	2.6
2	B	2	ILE	2.6
1	A	190	PRO	2.5
1	H	135	THR	2.5
1	A	137	THR	2.5
2	B	4	MET	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	L	60	SER	2.4
2	L	13	ALA	2.4
1	A	43	LYS	2.4
2	L	49	TYR	2.4
2	L	153	THR	2.4
2	L	167	SER	2.4
2	B	187	SER	2.4
2	L	183	THR	2.4
1	A	210	PRO	2.3
2	L	201	SER	2.3
2	L	17	GLY	2.3
2	L	57	GLY	2.3
2	L	207	ASN	2.3
1	A	75	THR	2.3
2	L	110	PRO	2.3
2	B	199	THR	2.3
1	H	133	SER	2.2
1	A	62	TRP	2.2
2	L	187	SER	2.2
2	B	15	VAL	2.2
2	B	110	PRO	2.2
2	B	103	GLU	2.2
2	L	199	THR	2.2
2	L	140	PHE	2.2
2	B	18	THR	2.2
2	B	48	ILE	2.2
2	B	109	ASP	2.2
2	L	65	SER	2.2
2	L	111	VAL	2.1
2	B	143	VAL	2.1
2	B	16	GLY	2.1
2	L	151	GLY	2.1
1	A	172	SER	2.1
1	A	186	SER	2.1
1	A	189	GLN	2.1
2	B	95(A)	ASP	2.1
2	L	95	SER	2.0
2	B	114	SER	2.0
2	B	14	ALA	2.0
2	L	81	ASP	2.0
2	B	75	ILE	2.0
2	L	109	ASP	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	L	150	ASP	2.0
2	B	81	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	PCA	A	2	8/9	0.75	0.25	61,71,84,84	0
1	PCA	H	2	8/9	0.82	0.15	46,61,73,73	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	L	301	5/5	0.84	0.15	80,85,85,88	0
3	SO4	A	301	5/5	0.85	0.12	66,69,72,74	0

6.5 Other polymers [i](#)

There are no such residues in this entry.