



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 08:35 AM UTC

PDB ID : 8PE3 / pdb_00008pe3
Title : Structure of Csm6' from Streptococcus thermophilus in complex with cyclic hexa-adenylate (cA6)
Authors : McQuarrie, S.J.; Athukoralage, J.S.; McMahon, S.A.; Graham, S.; Ackerman, K.; Bode, B.E.; White, M.F.; Gloster, T.M.
Deposited on : 2023-06-13
Resolution : 1.96 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.96 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)
RNA backbone	3983	1010 (2.32-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	390	4% 85% 14%
1	B	390	6% 88% 9%
2	C	6	17% 50% 17% 17%
3	D	3	33% 33% 33%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	E	3	 67% 33%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6600 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CRISPR system endoribonuclease Csm6'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	386	Total	C	N	O	S	0	1	0
			3057	1949	517	585	6			
1	B	379	Total	C	N	O	S	0	1	0
			2986	1908	500	572	6			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP A0A0A7HFE6
A	-2	ALA	-	expression tag	UNP A0A0A7HFE6
A	-1	ASN	-	expression tag	UNP A0A0A7HFE6
A	0	ALA	-	expression tag	UNP A0A0A7HFE6
A	1	MET	-	expression tag	UNP A0A0A7HFE6
A	2	GLY	-	expression tag	UNP A0A0A7HFE6
A	21	ALA	SER	variant	UNP A0A0A7HFE6
A	281	LYS	GLU	variant	UNP A0A0A7HFE6
A	375	GLN	HIS	variant	UNP A0A0A7HFE6
B	-3	GLY	-	expression tag	UNP A0A0A7HFE6
B	-2	ALA	-	expression tag	UNP A0A0A7HFE6
B	-1	ASN	-	expression tag	UNP A0A0A7HFE6
B	0	ALA	-	expression tag	UNP A0A0A7HFE6
B	1	MET	-	expression tag	UNP A0A0A7HFE6
B	2	GLY	-	expression tag	UNP A0A0A7HFE6
B	21	ALA	SER	variant	UNP A0A0A7HFE6
B	281	LYS	GLU	variant	UNP A0A0A7HFE6
B	375	GLN	HIS	variant	UNP A0A0A7HFE6

- Molecule 2 is a RNA chain called Cyclic hexaadenosine monophosphate (cA6).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	6	Total	C	N	O	P	0	6	0
			132	60	30	36	6			

- Molecule 3 is a RNA chain called RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	3	Total	C	N	O	P	0	3	0
			66	30	15	18	3			
3	E	3	Total	C	N	O	P	0	3	0
			66	30	15	18	3			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	135	Total	O	0	0
			135	135		
4	B	142	Total	O	0	0
			142	142		
4	C	6	Total	O	0	0
			6	6		
4	D	4	Total	O	0	0
			4	4		
4	E	6	Total	O	0	0
			6	6		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CRISPR system endoribonuclease Csm6'




- Molecule 1: CRISPR system endoribonuclease Csm6'



- Molecule 2: Cyclic hexaadenosine monophosphate (cA6)



- Molecule 3: RNA



- Molecule 3: RNA

Chain E:

67%

33%



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	99.91Å 99.91Å 286.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	22.34 – 1.96 22.34 – 1.96	Depositor EDS
% Data completeness (in resolution range)	54.8 (22.34-1.96) 54.8 (22.34-1.96)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.32 (at 1.97Å)	Xtriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
R, R_{free}	0.198 , 0.236 0.196 , 0.234	Depositor DCC
R_{free} test set	2902 reflections (2.77%)	wwPDB-VP
Wilson B-factor (Å ²)	38.0	Xtriage
Anisotropy	0.092	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 39.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6600	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A23, ADN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.24	0/3113	0.50	2/4209 (0.0%)
1	B	0.22	0/3040	0.44	0/4107
2	C	0.46	0/149	1.42	2/230 (0.9%)
3	D	0.46	0/24	0.95	0/35
3	E	0.51	0/24	1.23	0/35
All	All	0.24	0/6350	0.53	4/8616 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	58	PRO	CA-N-CD	-11.15	96.39	112.00
2	C	4[A]	A	P-O3'-C3'	-7.09	109.57	120.20
2	C	3[A]	A	P-O3'-C3'	-5.43	112.06	120.20
1	A	58	PRO	N-CA-CB	-5.23	97.98	103.52

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3057	0	2987	36	0
1	B	2986	0	2898	21	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	132	0	55	5	0
3	D	66	0	25	1	0
3	E	66	0	22	3	0
4	A	135	0	0	2	0
4	B	142	0	0	2	0
4	C	6	0	0	0	0
4	D	4	0	0	0	0
4	E	6	0	0	1	0
All	All	6600	0	5987	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:58:PRO:HD2	1:A:59:ASN:H	1.22	1.01
1:A:58:PRO:HD2	1:A:59:ASN:N	1.89	0.78
1:B:120:ARG:HH11	1:B:125:ASN:HD22	1.34	0.76
1:A:175:LYS:HB3	1:B:84:GLN:HE21	1.56	0.69
1:A:1:MET:HE2	1:A:33:LYS:HE3	1.78	0.65
1:A:3:VAL:HG13	1:A:33:LYS:HB3	1.80	0.63
1:B:152:GLU:O	1:B:156:VAL:HG22	2.00	0.61
1:A:231:LYS:HA	1:A:234:ARG:HD2	1.85	0.58
1:A:264:SER:O	1:A:268:ILE:HG13	2.04	0.57
1:A:1:MET:HG2	1:A:98:ASP:OD1	2.04	0.56
1:A:290:GLU:O	1:A:293:GLU:HG3	2.05	0.56
1:A:152:GLU:O	1:A:156:VAL:HG12	2.06	0.56
1:A:46:GLN:O	1:A:50:GLU:HG3	2.06	0.55
1:B:1:MET:HE3	1:B:33:LYS:HE2	1.89	0.54
1:B:85:ARG:NH2	4:B:505:HOH:O	2.40	0.53
1:B:9:GLY:HA3	3:E:3[B]:A23:O1C	2.08	0.53
1:A:150:ILE:O	1:A:154:ILE:HG13	2.09	0.53
1:A:335:ALA:O	1:B:270:ASN:ND2	2.40	0.52
1:B:214:ARG:HG2	1:B:218:ARG:NH2	2.23	0.52
1:A:225:SER:HB2	1:B:261:VAL:HG12	1.91	0.52
1:A:58:PRO:CD	1:A:59:ASN:N	2.70	0.51
1:B:110:GLN:H	1:B:110:GLN:CD	2.19	0.50
3:D:2[B]:A:H1'	3:D:3[B]:A23:C8	2.42	0.50
1:B:321:ILE:O	1:B:325:ILE:HG13	2.11	0.50
1:B:302:LEU:HD21	1:B:324:ILE:HG22	1.94	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:269:LYS:HE2	1:A:331:ARG:CZ	2.43	0.48
1:B:271:LEU:HD23	1:B:356:LEU:HD21	1.96	0.48
1:A:190:GLU:HG3	1:A:382:LEU:HD11	1.96	0.47
1:A:58:PRO:CD	1:A:59:ASN:H	2.11	0.47
1:A:202:ASP:OD1	1:A:214:ARG:HD2	2.14	0.47
1:A:289:ASP:OD2	1:A:289:ASP:N	2.47	0.47
2:C:2[A]:A:H5'	4:E:809:HOH:O	2.15	0.47
1:B:267:ARG:HD3	4:B:596:HOH:O	2.15	0.46
1:A:57:ALA:HB1	1:A:58:PRO:HD3	1.96	0.46
1:A:131:VAL:HG12	3:E:1[B]:ADN:H8	1.99	0.45
1:B:29:TYR:CE1	1:B:127:LYS:HE2	2.52	0.44
2:C:1[A]:A:O2'	2:C:2[A]:A:P	2.76	0.43
1:A:10:ASP:HB2	2:C:1[A]:A:O2'	2.17	0.43
1:B:273:GLU:HG3	1:B:302:LEU:HD13	2.01	0.43
1:A:191:LYS:HG3	1:B:188:PHE:CZ	2.54	0.42
1:A:104:LEU:HD11	1:A:128:ALA:HB1	2.01	0.42
1:A:200:ILE:O	1:A:203:GLN:HG2	2.20	0.42
2:C:4[A]:A:C8	2:C:4[A]:A:H5''	2.54	0.42
1:A:255:GLN:HA	1:A:255:GLN:OE1	2.19	0.42
1:A:74:ASN:HB2	4:A:492:HOH:O	2.19	0.42
1:A:239:LYS:HG3	1:A:242:GLU:CD	2.45	0.42
1:A:118:ILE:HG23	1:A:122:ASN:ND2	2.35	0.41
1:A:331:ARG:HD3	1:A:331:ARG:C	2.46	0.41
1:B:190:GLU:HG3	1:B:382:LEU:HD11	2.03	0.41
1:A:130:GLN:NE2	4:A:401:HOH:O	2.37	0.41
1:A:86:PHE:CE1	1:A:111:ILE:HG12	2.56	0.41
1:A:207:PHE:HB2	1:A:208:PRO:HD2	2.01	0.41
1:A:85[B]:ARG:HA	1:A:85[B]:ARG:HD3	1.95	0.41
1:B:216:GLU:OE2	1:B:216:GLU:HA	2.21	0.40
1:B:252:ILE:HD13	1:B:268:ILE:HG13	2.02	0.40
2:C:4[A]:A:H5''	2:C:4[A]:A:H8	1.86	0.40
1:A:379:LYS:O	1:A:383:THR:HG23	2.20	0.40
1:B:75:GLU:HA	1:B:78:ILE:HD12	2.03	0.40
3:E:2[B]:A:N3	3:E:2[B]:A:H2'	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	383/390 (98%)	373 (97%)	9 (2%)	1 (0%)	36	28
1	B	374/390 (96%)	364 (97%)	8 (2%)	2 (0%)	24	16
All	All	757/780 (97%)	737 (97%)	17 (2%)	3 (0%)	30	21

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	107	ALA
1	A	107	ALA
1	B	160	ASN

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	327/356 (92%)	323 (99%)	4 (1%)	63	61
1	B	316/356 (89%)	316 (100%)	0	100	100
All	All	643/712 (90%)	639 (99%)	4 (1%)	78	79

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	171	ASP
1	A	306	SER
1	A	356	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	372	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	-1	ASN
1	A	130	GLN
1	A	203	GLN
1	A	260	ASN
1	A	336	HIS
1	B	18	HIS
1	B	84	GLN
1	B	122	ASN
1	B	125	ASN
1	B	222	ASN
1	B	255	GLN
1	B	270	ASN
1	B	348	GLN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	C	5/6 (83%)	3 (60%)	1 (20%)
3	D	1/3 (33%)	1 (100%)	0
3	E	1/3 (33%)	1 (100%)	0
All	All	7/12 (58%)	5 (71%)	1 (14%)

All (5) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	C	2[A]	A
2	C	4[A]	A
2	C	5[A]	A
3	D	3[B]	A23
3	E	3[B]	A23

All (1) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	C	4[A]	A

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	A23	E	3[B]	3	23,28,29	0.41	0	32,43,46	0.66	1 (3%)
3	A23	D	3[B]	3	23,28,29	0.42	0	32,43,46	0.62	1 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	A23	E	3[B]	3	-	3/7/35/36	0/4/4/4
3	A23	D	3[B]	3	-	1/7/35/36	0/4/4/4

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	3[B]	A23	O4'-C1'-C2'	-2.50	102.29	106.59
3	D	3[B]	A23	O4'-C1'-C2'	-2.23	102.74	106.59

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	E	3[B]	A23	O4'-C4'-C5'-O5'
3	E	3[B]	A23	C3'-C4'-C5'-O5'
3	E	3[B]	A23	C4'-C5'-O5'-P
3	D	3[B]	A23	C4'-C5'-O5'-P

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	3[B]	A23	1	0
3	D	3[B]	A23	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	386/390 (98%)	0.32	17 (4%) 39 45	21, 40, 64, 80	1 (0%)
1	B	379/390 (97%)	0.22	22 (5%) 29 33	20, 38, 64, 80	1 (0%)
2	C	6/6 (100%)	-0.06	0 100 100	16, 17, 17, 18	6 (100%)
3	D	1/3 (33%)	-0.29	0 100 100	17, 17, 17, 17	1 (100%)
3	E	1/3 (33%)	-0.20	0 100 100	16, 16, 16, 16	1 (100%)
All	All	773/792 (97%)	0.27	39 (5%) 34 40	16, 39, 64, 80	10 (1%)

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	335	ALA	7.1
1	A	334	VAL	5.8
1	A	336	HIS	4.1
1	B	337	SER	4.0
1	A	294	ASP	3.8
1	B	294	ASP	3.8
1	B	265	PHE	3.7
1	B	346	VAL	3.1
1	A	299	TYR	3.0
1	B	299	TYR	3.0
1	A	386	ASN	3.0
1	B	0	ALA	2.8
1	B	349	LEU	2.8
1	B	330	SER	2.7
1	B	303	PHE	2.7
1	B	326	ASP	2.6
1	B	331	ARG	2.6
1	A	58	PRO	2.6
1	A	327	MET	2.6
1	B	344	ASP	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	345	ALA	2.6
1	B	17	PHE	2.5
1	B	266	ILE	2.4
1	A	260	ASN	2.4
1	A	330	SER	2.4
1	B	386	ASN	2.3
1	B	329	SER	2.3
1	B	343	SER	2.3
1	A	331	ARG	2.2
1	A	266	ILE	2.2
1	A	259	GLY	2.2
1	A	265	PHE	2.1
1	B	58	PRO	2.1
1	B	345	ALA	2.1
1	A	337	SER	2.1
1	A	156	VAL	2.1
1	B	327	MET	2.1
1	B	267	ARG	2.1
1	B	332	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	A23	E	3[B]	25/26	0.96	0.08	27,31,33,33	25
3	A23	D	3[B]	25/26	0.97	0.08	31,33,36,37	25

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.