



Full wwPDB NMR Structure Validation Report ⓘ

Mar 27, 2026 – 04:48 AM UTC

PDB ID : 2PFU / pdb_00002pfu
Title : NMR structure determination of the periplasmic domain of ExbD from E.coli
Authors : Garcia-Herrero, A.; Peacock, S.R.; Howard, P.S.; Vogel, H.J.
Deposited on : 2007-04-05

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	99	43% 25% . 27%

2 Ensemble composition and analysis

This entry contains 19 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:63-A:134 (72)	0.66	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 2, 3, 4, 5, 6, 10, 12, 13, 14, 15, 16, 17, 18, 19
2	7, 8, 11
Single-model clusters	9

3 Entry composition


There is only 1 type of molecule in this entry. The entry contains 1536 atoms, of which 775 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Biopolymer transport exbD protein.

Mol	Chain	Residues	Atoms						Trace
1	A	99	Total	C	H	N	O	S	0
			1536	481	775	123	152	5	


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	43	MET	-	initiating methionine	UNP P0ABV2




4.2.3 Score per residue for model 3

- Molecule 1: Biopolymer transport exbD protein




4.2.4 Score per residue for model 4

- Molecule 1: Biopolymer transport exbD protein




4.2.5 Score per residue for model 5

- Molecule 1: Biopolymer transport exbD protein




4.2.6 Score per residue for model 6

- Molecule 1: Biopolymer transport exbD protein




4.2.7 Score per residue for model 7

- Molecule 1: Biopolymer transport exbD protein




4.2.8 Score per residue for model 8

- Molecule 1: Biopolymer transport exbD protein




4.2.9 Score per residue for model 9

- Molecule 1: Biopolymer transport exbD protein




4.2.10 Score per residue for model 10

- Molecule 1: Biopolymer transport exbD protein




4.2.11 Score per residue for model 11

- Molecule 1: Biopolymer transport exbD protein




4.2.12 Score per residue for model 12

- Molecule 1: Biopolymer transport exbD protein




4.2.13 Score per residue for model 13

- Molecule 1: Biopolymer transport exbD protein




4.2.14 Score per residue for model 14

- Molecule 1: Biopolymer transport exbD protein




4.2.15 Score per residue for model 15

- Molecule 1: Biopolymer transport exbD protein




4.2.16 Score per residue for model 16

- Molecule 1: Biopolymer transport exbD protein




4.2.17 Score per residue for model 17

- Molecule 1: Biopolymer transport exbD protein




4.2.18 Score per residue for model 18

- Molecule 1: Biopolymer transport exbD protein



4.2.19 Score per residue for model 19

- Molecule 1: Biopolymer transport exbD protein



5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics*.

Of the 20 calculated structures, 19 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	refinement	2.0
CYANA	structure solution	2.0

No chemical shift data was provided.

6 Model quality

6.1 Standard geometry

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	556	562	562	10±2
All	All	10564	10678	10678	182

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:106:ALA:HB1	1:A:110:VAL:HG21	0.81	1.49	18	3
1:A:86:MET:HE2	1:A:87:ILE:HD13	0.80	1.51	5	1
1:A:74:MET:HE1	1:A:86:MET:HE2	0.68	1.65	14	1
1:A:112:TYR:CD1	1:A:115:LEU:HD11	0.63	2.29	5	12
1:A:90:LEU:O	1:A:94:THR:HG22	0.62	1.94	10	1
1:A:105:ARG:HB2	1:A:133:VAL:HG11	0.62	1.72	10	6
1:A:68:VAL:HG13	1:A:74:MET:HG2	0.59	1.74	14	2
1:A:105:ARG:HG2	1:A:133:VAL:HG11	0.58	1.76	6	3
1:A:74:MET:CE	1:A:86:MET:HE2	0.56	2.29	14	1
1:A:86:MET:O	1:A:90:LEU:HD12	0.56	2.00	19	9
1:A:71:ASP:O	1:A:114:THR:HG21	0.56	2.01	12	6
1:A:90:LEU:HD13	1:A:90:LEU:C	0.55	2.26	18	2
1:A:105:ARG:HD2	1:A:133:VAL:HG21	0.55	1.78	15	1
1:A:129:LYS:O	1:A:130:ILE:HD12	0.55	2.02	15	2

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:112:TYR:CD1	1:A:115:LEU:HD22	0.54	2.37	4	3
1:A:122:LEU:HD13	1:A:127:TYR:CD2	0.54	2.38	8	3
1:A:90:LEU:HD22	1:A:127:TYR:OH	0.54	2.02	7	11
1:A:64:VAL:HG12	1:A:102:ILE:HG23	0.53	1.81	13	7
1:A:72:ASN:CG	1:A:114:THR:HG22	0.53	2.28	13	1
1:A:107:ASP:OD1	1:A:109:THR:HG22	0.53	2.03	17	1
1:A:115:LEU:HD12	1:A:116:MET:N	0.53	2.18	15	16
1:A:107:ASP:O	1:A:110:VAL:HG23	0.52	2.04	17	10
1:A:68:VAL:HG13	1:A:74:MET:CG	0.52	2.34	18	2
1:A:68:VAL:CG1	1:A:74:MET:HE3	0.52	2.34	16	6
1:A:72:ASN:OD1	1:A:74:MET:HE1	0.51	2.06	7	1
1:A:114:THR:O	1:A:118:VAL:HG22	0.50	2.06	11	3
1:A:68:VAL:HG13	1:A:74:MET:HG3	0.50	1.82	16	9
1:A:90:LEU:HD13	1:A:127:TYR:OH	0.50	2.06	10	7
1:A:66:LEU:HD21	1:A:90:LEU:HD11	0.50	1.82	19	2
1:A:90:LEU:O	1:A:90:LEU:HD22	0.50	2.07	9	1
1:A:105:ARG:CD	1:A:133:VAL:HG21	0.49	2.36	15	1
1:A:66:LEU:HD12	1:A:102:ILE:CG2	0.49	2.37	18	3
1:A:90:LEU:O	1:A:94:THR:HG23	0.48	2.08	17	1
1:A:68:VAL:CG1	1:A:74:MET:HE2	0.47	2.39	11	1
1:A:105:ARG:CB	1:A:133:VAL:HG11	0.47	2.39	10	4
1:A:104:PHE:O	1:A:133:VAL:HG12	0.47	2.10	16	10
1:A:90:LEU:HD13	1:A:91:ASN:N	0.47	2.24	9	2
1:A:86:MET:HG2	1:A:90:LEU:HD12	0.46	1.87	10	2
1:A:105:ARG:NE	1:A:133:VAL:HG21	0.46	2.25	15	1
1:A:107:ASP:OD2	1:A:109:THR:HG22	0.46	2.10	4	1
1:A:74:MET:HB2	1:A:81:VAL:HG23	0.46	1.87	2	2
1:A:87:ILE:HD11	1:A:121:THR:HG22	0.45	1.87	17	2
1:A:69:LYS:O	1:A:70:ALA:HB3	0.45	2.12	16	1
1:A:112:TYR:CE1	1:A:115:LEU:HD22	0.44	2.47	7	3
1:A:107:ASP:CG	1:A:110:VAL:HG23	0.43	2.38	6	1
1:A:116:MET:HE3	1:A:119:MET:HE2	0.43	1.91	11	1
1:A:87:ILE:HD12	1:A:125:ALA:HB1	0.43	1.91	15	1
1:A:87:ILE:HD11	1:A:121:THR:CG2	0.42	2.43	16	1
1:A:115:LEU:C	1:A:115:LEU:HD23	0.42	2.39	7	3
1:A:102:ILE:HD13	1:A:127:TYR:CE1	0.42	2.49	9	1
1:A:107:ASP:HB3	1:A:109:THR:HG22	0.41	1.92	16	1
1:A:112:TYR:HD1	1:A:115:LEU:HD11	0.41	1.74	3	1
1:A:115:LEU:HD12	1:A:115:LEU:C	0.41	2.41	6	1
1:A:90:LEU:C	1:A:90:LEU:HD22	0.41	2.40	9	1
1:A:89:ALA:O	1:A:93:LEU:HD12	0.41	2.15	16	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:68:VAL:HG12	1:A:73:SER:O	0.40	2.16	15	1
1:A:107:ASP:OD2	1:A:110:VAL:HG23	0.40	2.16	6	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	72/99 (73%)	62±1 (87±2%)	7±2 (9±3%)	3±1 (4±1%)	4	30
All	All	1368/1881 (73%)	1187 (87%)	127 (9%)	54 (4%)	4	30

All 5 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	70	ALA	16
1	A	127	TYR	15
1	A	72	ASN	11
1	A	77	GLY	11
1	A	71	ASP	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	62/86 (72%)	45±2 (73±3%)	17±2 (27±3%)	2	21
All	All	1178/1634 (72%)	858 (73%)	320 (27%)	2	21

All 40 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	68	VAL	19
1	A	82	THR	19
1	A	101	THR	19
1	A	74	MET	17
1	A	114	THR	15
1	A	90	LEU	14
1	A	88	THR	13
1	A	123	HIS	13
1	A	111	ASP	12
1	A	120	ASP	12
1	A	98	LYS	12
1	A	69	LYS	10
1	A	113	GLU	10
1	A	129	LYS	10
1	A	128	LEU	9
1	A	73	SER	9
1	A	117	LYS	9
1	A	84	GLU	7
1	A	108	LYS	7
1	A	79	ASP	6
1	A	124	GLN	6
1	A	105	ARG	6
1	A	71	ASP	6
1	A	122	LEU	6
1	A	116	MET	5
1	A	119	MET	5
1	A	91	ASN	5
1	A	95	GLU	5
1	A	132	LEU	5
1	A	78	ASN	4
1	A	67	SER	4
1	A	107	ASP	4
1	A	66	LEU	4
1	A	65	TYR	3
1	A	97	LYS	3
1	A	99	ASP	3
1	A	93	LEU	1
1	A	83	ASP	1
1	A	86	MET	1
1	A	72	ASN	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided