



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:30 AM UTC

PDB ID : 3PFK / pdb_00003pfk
Title : PHOSPHOFRUCTOKINASE. STRUCTURE AND CONTROL
Authors : Evans, P.R.; Hudson, P.J.
Deposited on : 1988-01-25
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	319	60% 32% 8%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2462 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PHOSPHOFRUCTOKINASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	319	Total	C	N	O	S	0	0	0
			2361	1477	422	454	8			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	95	GLN	GLU	conflict	UNP P00512

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	A	1	Total	O	P	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	91	Total	O	0	0
			91	91		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PHOSPHOFRUCTOKINASE



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	122.50Å 84.10Å 61.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.40 30.00 – 2.40	Depositor EDS
% Data completeness (in resolution range)	(Not available) (30.00-2.40) 99.0 (30.00-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.67 (at 2.40Å)	Xtriage
Refinement program	PROLSQ	Depositor
R, R_{free}	0.171 , (Not available) 0.170 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	30.5	Xtriage
Anisotropy	0.325	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 67.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2462	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.26% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.00	0/2397	2.06	84/3239 (2.6%)

There are no bond length outliers.

All (84) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	127	ASP	CA-CB-CG	-8.78	103.82	112.60
1	A	127	ASP	N-CA-C	8.39	121.36	111.71
1	A	277	GLU	CA-CB-CG	7.93	129.97	114.10
1	A	288	ASN	OD1-CG-ND2	7.55	130.15	122.60
1	A	21	ARG	CA-C-O	-7.43	112.67	120.55
1	A	295	ASP	CA-C-O	-7.32	112.71	121.05
1	A	152	LYS	O-C-N	7.27	130.44	122.15
1	A	290	GLN	O-C-N	7.21	131.64	123.27
1	A	160	HIS	CB-CA-C	-7.17	98.12	110.09
1	A	59	ASP	CA-CB-CG	7.10	119.70	112.60
1	A	154	ARG	NE-CZ-NH1	7.09	128.59	121.50
1	A	254	GLY	CA-C-O	6.86	126.68	122.22
1	A	114	GLU	N-CA-C	-6.70	104.20	112.38
1	A	98	VAL	N-CA-CB	6.70	119.05	111.21
1	A	58	GLY	N-CA-C	-6.66	103.05	112.37
1	A	48	ASN	OD1-CG-ND2	6.57	129.17	122.60
1	A	78	THR	CA-CB-OG1	-6.54	99.79	109.60
1	A	149	ALA	O-C-N	6.53	129.15	122.09
1	A	238	THR	O-C-N	6.42	131.13	122.59
1	A	230	PHE	CA-CB-CG	-6.41	107.39	113.80
1	A	92	HIS	N-CA-CB	6.38	120.03	110.65
1	A	295	ASP	N-CA-C	-6.26	100.31	110.20
1	A	14	PRO	CA-C-O	-6.24	114.56	121.23
1	A	222	GLU	CA-C-N	6.23	127.01	120.03

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	222	GLU	C-N-CA	6.23	127.01	120.03
1	A	88	GLN	O-C-N	6.23	128.72	122.12
1	A	144	ASN	CB-CA-C	6.17	121.04	110.79
1	A	104	GLY	N-CA-C	-6.13	105.21	113.24
1	A	117	PHE	O-C-N	6.12	128.10	121.12
1	A	280	GLY	N-CA-C	-6.10	102.19	111.10
1	A	135	PHE	CA-CB-CG	-6.06	107.74	113.80
1	A	236	GLU	CA-CB-CG	5.99	126.07	114.10
1	A	126	ILE	CA-C-N	5.98	128.89	120.28
1	A	126	ILE	C-N-CA	5.98	128.89	120.28
1	A	228	VAL	O-C-N	5.98	127.99	121.83
1	A	127	ASP	CA-C-O	-5.88	113.46	120.10
1	A	159	SER	CA-C-N	5.87	132.22	122.54
1	A	159	SER	C-N-CA	5.87	132.22	122.54
1	A	135	PHE	O-C-N	5.78	130.32	123.27
1	A	180	SER	CB-CA-C	-5.73	101.81	110.92
1	A	115	HIS	N-CA-CB	5.73	119.23	110.70
1	A	77	LYS	N-CA-C	-5.72	105.56	112.54
1	A	202	ILE	O-C-N	5.69	127.39	121.87
1	A	197	ASP	CA-CB-CG	-5.67	106.92	112.60
1	A	269	ALA	CA-C-N	5.64	127.77	120.44
1	A	269	ALA	C-N-CA	5.64	127.77	120.44
1	A	130	ILE	O-C-N	5.62	127.50	121.10
1	A	79	GLU	N-CA-C	-5.59	105.28	111.71
1	A	216	SER	N-CA-CB	-5.57	101.65	111.39
1	A	288	ASN	CA-CB-CG	-5.55	107.05	112.60
1	A	305	HIS	N-CA-CB	5.53	119.29	110.16
1	A	30	HIS	CA-CB-CG	-5.52	108.28	113.80
1	A	190	LEU	CA-C-O	-5.46	114.62	120.46
1	A	87	GLU	CB-CG-CD	5.44	121.84	112.60
1	A	142	ALA	CA-C-N	5.43	127.50	120.44
1	A	142	ALA	C-N-CA	5.43	127.50	120.44
1	A	210	GLU	O-C-N	5.43	127.66	122.07
1	A	129	ASP	CA-C-O	5.42	127.46	121.39
1	A	176	ILE	O-C-N	5.40	127.79	121.96
1	A	56	ASP	N-CA-C	-5.39	105.96	112.54
1	A	303	ASN	CA-CB-CG	5.37	117.97	112.60
1	A	135	PHE	CA-C-O	-5.36	114.49	120.66
1	A	97	LEU	N-CA-C	5.36	117.82	108.76
1	A	107	GLN	CB-CG-CD	-5.33	103.53	112.60
1	A	59	ASP	N-CA-C	5.25	118.96	112.24
1	A	83	LYS	CA-C-O	5.24	126.09	120.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	132	GLY	O-C-N	5.24	128.39	122.42
1	A	3	ARG	N-CA-CB	-5.23	102.03	110.71
1	A	187	GLU	CB-CG-CD	5.19	121.42	112.60
1	A	241	GLU	N-CA-C	-5.16	101.73	109.15
1	A	17	ASN	OD1-CG-ND2	5.15	127.75	122.60
1	A	165	VAL	N-CA-CB	5.14	117.94	111.46
1	A	268	GLY	CA-C-O	-5.12	115.48	120.75
1	A	231	GLY	CA-C-N	5.11	127.09	120.44
1	A	231	GLY	C-N-CA	5.11	127.09	120.44
1	A	107	GLN	OE1-CD-NE2	5.11	127.71	122.60
1	A	78	THR	N-CA-C	5.10	118.26	110.20
1	A	169	MET	CA-C-O	5.09	127.79	121.94
1	A	175	ASP	O-C-N	5.06	127.49	122.08
1	A	212	GLY	N-CA-C	5.04	122.37	115.27
1	A	257	THR	CA-CB-OG1	-5.03	102.05	109.60
1	A	188	THR	O-C-N	5.02	129.28	123.41
1	A	280	GLY	CA-C-O	-5.02	115.71	121.49
1	A	180	SER	O-C-N	5.01	127.44	122.08

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2361	0	2330	48	0
2	A	10	0	0	0	0
3	A	91	0	0	0	0
All	All	2462	0	2330	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:204:ARG:HH22	1:A:319:SER:HB2	1.50	0.76
1:A:78:THR:HG22	1:A:80:GLU:H	1.52	0.73
1:A:38:TYR:O	1:A:43:GLY:HA3	1.94	0.67
1:A:295:ASP:HB3	1:A:298:GLU:HG2	1.78	0.65
1:A:60:ILE:HD13	1:A:67:ILE:HD13	1.81	0.63
1:A:3:ARG:HG2	1:A:94:ILE:HA	1.80	0.62
1:A:44:LEU:HD11	1:A:89:LEU:HD13	1.82	0.61
1:A:89:LEU:HA	1:A:94:ILE:HD12	1.84	0.60
1:A:230:PHE:O	1:A:234:ILE:HD12	2.04	0.58
1:A:62:HIS:H	1:A:62:HIS:CD2	2.23	0.56
1:A:1:MET:HE2	1:A:95:GLN:C	2.31	0.55
1:A:111:LYS:O	1:A:115:HIS:HD2	1.89	0.55
1:A:78:THR:HG22	1:A:80:GLU:N	2.23	0.54
1:A:175:ASP:OD2	1:A:305:HIS:HE1	1.93	0.51
1:A:210:GLU:C	1:A:212:GLY:H	2.17	0.51
1:A:192:PRO:HD2	1:A:193:GLU:OE1	2.11	0.51
1:A:86:ILE:HG23	1:A:117:PHE:CE1	2.47	0.50
1:A:283:CYS:HB3	1:A:296:ILE:HG12	1.93	0.49
1:A:282:ARG:HA	1:A:294:HIS:O	2.13	0.49
1:A:128:ASN:OD1	1:A:135:PHE:HA	2.13	0.48
1:A:143:LEU:HD11	1:A:179:TRP:HB2	1.96	0.48
1:A:41:TYR:O	1:A:44:LEU:HB3	2.13	0.48
1:A:26:LYS:HE3	1:A:30:HIS:NE2	2.29	0.48
1:A:141:THR:HA	1:A:257:THR:HG23	1.95	0.48
1:A:160:HIS:HB3	1:A:162:ARG:HG3	1.95	0.48
1:A:247:LEU:HB2	1:A:251:GLN:NE2	2.29	0.48
1:A:86:ILE:HG23	1:A:117:PHE:CD1	2.50	0.47
1:A:32:VAL:HG21	1:A:276:LEU:HD21	1.97	0.46
1:A:197:ASP:HB3	1:A:200:ASP:OD1	2.17	0.45
1:A:44:LEU:CD1	1:A:89:LEU:HD13	2.46	0.45
1:A:210:GLU:C	1:A:212:GLY:N	2.76	0.44
1:A:129:ASP:HB3	1:A:173:ALA:CB	2.48	0.43
1:A:189:ILE:HG22	1:A:191:ILE:HG23	2.00	0.43
1:A:276:LEU:C	1:A:278:GLY:H	2.25	0.43
1:A:1:MET:HE1	1:A:118:PRO:HG2	1.99	0.43
1:A:232:ARG:C	1:A:232:ARG:HD2	2.44	0.43
1:A:14:PRO:CB	1:A:141:THR:HG21	2.49	0.42
1:A:113:THR:O	1:A:116:GLY:N	2.46	0.42
1:A:1:MET:HE2	1:A:96:GLY:N	2.35	0.42
1:A:82:GLN:O	1:A:86:ILE:HG13	2.20	0.41
1:A:213:LYS:HD2	1:A:320:ILE:HG21	2.03	0.41
1:A:197:ASP:OD2	1:A:199:ASN:HB2	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:255:SER:HA	1:A:256:PRO:HD2	1.81	0.41
1:A:311:MET:HE2	1:A:311:MET:HB3	1.95	0.40
1:A:19:ALA:O	1:A:23:VAL:HG23	2.21	0.40
1:A:175:ASP:OD2	1:A:305:HIS:CE1	2.73	0.40
1:A:1:MET:HG2	1:A:95:GLN:HB2	2.02	0.40
1:A:213:LYS:HE2	1:A:215:HIS:NE2	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	317/319 (99%)	308 (97%)	7 (2%)	2 (1%)	21	32

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	277	GLU
1	A	211	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	237/247 (96%)	219 (92%)	18 (8%)	12	21

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	7	LEU
1	A	48	ASN
1	A	59	ASP
1	A	66	THR
1	A	79	GLU
1	A	87	GLU
1	A	89	LEU
1	A	105	SER
1	A	107	GLN
1	A	112	LEU
1	A	117	PHE
1	A	148	ASP
1	A	153	ILE
1	A	187	GLU
1	A	232	ARG
1	A	310	ARG
1	A	319	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	62	HIS
1	A	115	HIS
1	A	209	HIS
1	A	288	ASN
1	A	305	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	A	322	-	4,4,4	0.85	0	6,6,6	1.30	1 (16%)
2	PO4	A	321	-	4,4,4	0.62	0	6,6,6	0.82	0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	A	322	PO4	O4-P-O2	2.58	115.94	107.91

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	319/319 (100%)	-0.30	0 100 100	13, 39, 71, 93	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PO4	A	321	5/5	0.91	0.10	67,67,71,71	0
2	PO4	A	322	5/5	0.96	0.07	49,50,56,59	0

6.5 Other polymers [i](#)

There are no such residues in this entry.