



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 05:39 AM UTC

PDB ID : 1PGQ / pdb_00001pgq
Title : CRYSTALLOGRAPHIC STUDY OF COENZYME, COENZYME ANALOGUE AND SUBSTRATE BINDING IN 6-PHOSPHOGLUCONATE DEHYDROGENASE: IMPLICATIONS FOR NADP SPECIFICITY AND THE ENZYME MECHANISM
Authors : Adams, M.J.; Phillips, C.; Gover, S.
Deposited on : 1994-07-18
Resolution : 3.17 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.17 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	2119 (3.20-3.16)
Ramachandran outliers	187476	2070 (3.20-3.16)
Sidechain outliers	187428	2069 (3.20-3.16)
RSRZ outliers	180081	2001 (3.20-3.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	482	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4113 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 6-PHOSPHOGLUCONATE DEHYDROGENASE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	473	Total	C	N	O	S	0	0	0
			3654	2332	634	666	22			

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is ADENOSINE-2'-MONOPHOSPHATE (CCD ID: 2AM) (formula: C₁₀H₁₄N₅O₇P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			23	10	5	7	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	421	Total	O	0	0
			421	421		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 6-PHOSPHOGLUCONATE DEHYDROGENASE



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	72.74Å 148.40Å 102.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 3.17 74.20 – 3.17	Depositor EDS
% Data completeness (in resolution range)	(Not available) ((Not available)-3.17) 91.9 (74.20-3.17)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.56 (at 3.15Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.169 , (Not available) 0.169 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	36.5	Xtriage
Anisotropy	0.179	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 85.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4113	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, 2AM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.72	2/3731 (0.1%)	1.71	48/5028 (1.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	88	VAL	CA-CB	6.23	1.58	1.53
1	A	253	ILE	CA-CB	5.84	1.61	1.54

All (48) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	76	ALA	N-CA-C	9.74	122.95	108.14
1	A	422	PHE	CA-CB-CG	9.28	123.08	113.80
1	A	377	ASP	CA-CB-CG	8.99	121.59	112.60
1	A	76	ALA	O-C-N	8.76	132.05	123.29
1	A	449	PHE	CA-CB-CG	8.06	121.86	113.80
1	A	452	HIS	N-CA-C	7.29	122.01	113.18
1	A	402	GLN	CA-C-O	-7.07	112.75	121.02
1	A	102	ASN	CA-CB-CG	7.06	119.66	112.60
1	A	75	LYS	N-CA-C	-7.01	99.28	109.59
1	A	139	SER	N-CA-C	-6.92	99.27	110.20
1	A	433	ARG	CA-CB-CG	6.90	127.91	114.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	458	ALA	N-CA-C	-6.76	98.61	109.96
1	A	365	CYS	N-CA-C	6.29	118.67	110.43
1	A	163	VAL	CB-CA-C	-6.25	102.66	111.21
1	A	20	ASN	CA-CB-CG	6.11	118.71	112.60
1	A	109	MET	O-C-N	6.09	128.43	122.09
1	A	440	ASN	CA-CB-CG	6.06	118.66	112.60
1	A	88	VAL	CA-C-O	-6.01	113.74	118.85
1	A	91	LEU	N-CA-C	5.93	119.15	109.59
1	A	378	ALA	N-CA-C	-5.91	104.76	111.14
1	A	87	LEU	N-CA-C	5.87	117.68	111.28
1	A	402	GLN	N-CA-C	5.83	120.93	111.37
1	A	278	THR	N-CA-C	5.83	118.85	111.69
1	A	33	ARG	CB-CG-CD	-5.81	97.94	111.30
1	A	88	VAL	CB-CA-C	-5.76	108.24	114.35
1	A	402	GLN	CG-CD-NE2	5.65	124.87	116.40
1	A	445	GLN	OE1-CD-NE2	-5.49	117.11	122.60
1	A	323	ARG	CA-CB-CG	5.43	124.97	114.10
1	A	264	LYS	CA-CB-CG	5.35	124.81	114.10
1	A	12	VAL	CB-CA-C	-5.32	105.16	111.97
1	A	471	HIS	N-CA-C	-5.32	102.21	109.71
1	A	85	GLU	O-C-N	5.31	127.55	122.03
1	A	78	GLN	CG-CD-NE2	5.28	124.33	116.40
1	A	345	ALA	N-CA-C	-5.28	105.42	111.07
1	A	139	SER	N-CA-CB	5.24	117.98	110.06
1	A	78	GLN	OE1-CD-NE2	-5.24	117.36	122.60
1	A	97	ILE	N-CA-C	-5.24	100.58	108.12
1	A	45	GLU	N-CA-C	5.23	118.44	111.75
1	A	404	SER	N-CA-C	-5.16	105.56	111.14
1	A	393	PHE	CA-C-N	5.14	127.44	120.44
1	A	393	PHE	C-N-CA	5.14	127.44	120.44
1	A	180	HIS	CA-CB-CG	-5.14	108.66	113.80
1	A	40	ASP	N-CA-C	-5.13	105.37	110.97
1	A	17	LEU	CA-C-N	5.11	127.19	120.60
1	A	17	LEU	C-N-CA	5.11	127.19	120.60
1	A	206	ASP	CA-CB-CG	5.11	117.71	112.60
1	A	453	THR	CA-CB-OG1	5.05	117.17	109.60
1	A	176	ASP	CA-CB-CG	5.02	117.62	112.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	273	TYR	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3654	0	3661	59	0
2	A	15	0	0	0	0
3	A	23	0	12	0	0
4	A	421	0	0	10	0
All	All	4113	0	3673	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:183:LYS:O	1:A:186:HIS:HB3	1.97	0.65
1:A:163:VAL:HG11	1:A:362:ARG:HD3	1.79	0.64
1:A:16:ASN:HD22	1:A:135:ARG:HD3	1.61	0.64
1:A:204:MET:HB2	1:A:215:MET:HE3	1.81	0.62
1:A:6:ALA:HB2	1:A:64:LEU:HD21	1.82	0.61
1:A:98:ILE:HG21	1:A:156:PHE:HZ	1.67	0.59
1:A:121:LEU:HB3	1:A:148:ALA:HB2	1.88	0.55
1:A:142:PRO:HG2	1:A:172:TRP:HA	1.89	0.55
1:A:430:ASP:O	1:A:434:HIS:HD2	1.90	0.54
1:A:362:ARG:HH11	1:A:362:ARG:HG2	1.73	0.53
1:A:58:GLU:HG3	1:A:90:LEU:HD11	1.92	0.52
1:A:350:TRP:HA	4:A:598:HOH:O	2.09	0.52
1:A:61:VAL:HG11	1:A:90:LEU:HD12	1.93	0.51
1:A:221:GLU:HG2	4:A:911:HOH:O	2.11	0.50
1:A:316:LYS:HG3	4:A:987:HOH:O	2.10	0.50
1:A:307:GLN:HB3	4:A:971:HOH:O	2.12	0.50
1:A:76:ALA:HB3	1:A:80:VAL:HG23	1.93	0.50
1:A:231:LEU:HB2	4:A:605:HOH:O	2.12	0.49
1:A:80:VAL:O	1:A:84:ILE:HG13	2.13	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:24:HIS:HA	4:A:719:HOH:O	2.12	0.48
1:A:21:MET:HG2	1:A:159:ILE:HG21	1.96	0.47
1:A:289:LEU:HD22	1:A:296:ARG:HD3	1.96	0.47
1:A:323:ARG:HG3	1:A:323:ARG:HH11	1.80	0.47
1:A:362:ARG:HG2	1:A:362:ARG:NH1	2.30	0.47
1:A:279:LEU:HD13	1:A:333:SER:HB2	1.97	0.46
1:A:81:ASP:HA	1:A:84:ILE:HD12	1.97	0.46
1:A:115:LEU:HD22	1:A:122:PHE:HB2	1.97	0.46
1:A:98:ILE:HA	1:A:123:VAL:O	2.16	0.46
1:A:471:HIS:HA	4:A:1000:HOH:O	2.16	0.45
1:A:125:SER:HB3	1:A:142:PRO:HA	1.99	0.45
1:A:321:ASP:HA	1:A:404:SER:HB3	1.99	0.45
1:A:9:GLY:O	1:A:14:GLY:HA3	2.16	0.45
1:A:222:TRP:CE2	1:A:323:ARG:HG2	2.52	0.45
1:A:229:SER:HB3	1:A:369:SER:HB3	1.99	0.45
1:A:227:LEU:HD12	1:A:323:ARG:HD2	1.99	0.44
1:A:10:LEU:HG	1:A:32:ASN:HB2	1.99	0.44
1:A:430:ASP:HA	1:A:433:ARG:HD3	1.99	0.44
1:A:131:GLU:HG3	4:A:1003:HOH:O	2.17	0.44
1:A:86:LYS:O	1:A:89:PRO:HD2	2.16	0.44
1:A:204:MET:HE1	1:A:219:PHE:CE2	2.53	0.44
1:A:377:ASP:O	1:A:381:ARG:HB2	2.18	0.43
1:A:73:LEU:HD23	1:A:73:LEU:HA	1.92	0.43
1:A:76:ALA:HB2	1:A:79:ALA:HB3	1.99	0.43
1:A:58:GLU:O	1:A:61:VAL:HG12	2.19	0.43
1:A:84:ILE:O	1:A:88:VAL:HG23	2.19	0.42
1:A:7:LEU:HD13	1:A:18:ILE:HG13	2.02	0.42
1:A:67:PRO:O	1:A:69:ARG:HG3	2.19	0.42
1:A:98:ILE:HG21	1:A:156:PHE:CZ	2.51	0.42
1:A:135:ARG:HA	1:A:135:ARG:HD2	1.87	0.42
1:A:217:LYS:HE2	4:A:1650:HOH:O	2.19	0.42
1:A:459:LYS:N	1:A:460:PRO:HD3	2.34	0.42
1:A:14:GLY:HA2	1:A:73:LEU:HD12	2.01	0.42
1:A:245:ASP:OD2	1:A:247:LYS:HB2	2.19	0.42
1:A:102:ASN:HD22	1:A:183:LYS:HE3	1.86	0.41
1:A:63:LYS:HE2	4:A:1659:HOH:O	2.19	0.41
1:A:205:LYS:HB2	1:A:215:MET:HE1	2.02	0.41
1:A:114:ASP:O	1:A:117:ASP:HB3	2.19	0.41
1:A:319:LEU:HD12	1:A:319:LEU:HA	1.81	0.41
1:A:417:ILE:HA	1:A:418:PRO:HD2	1.92	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	471/482 (98%)	436 (93%)	28 (6%)	7 (2%)	8	35

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	164	GLY
1	A	176	ASP
1	A	469	THR
1	A	261	GLY
1	A	2	GLN
1	A	137	GLY
1	A	9	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	378/386 (98%)	336 (89%)	42 (11%)	6	23

All (42) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	VAL
1	A	19	LEU
1	A	35	VAL
1	A	36	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	57	LEU
1	A	61	VAL
1	A	70	ILE
1	A	75	LYS
1	A	78	GLN
1	A	89	PRO
1	A	123	VAL
1	A	125	SER
1	A	142	PRO
1	A	145	ASN
1	A	163	VAL
1	A	176	ASP
1	A	184	MET
1	A	202	HIS
1	A	208	LEU
1	A	213	LYS
1	A	221	GLU
1	A	224	LYS
1	A	247	LYS
1	A	252	LYS
1	A	253	ILE
1	A	260	LYS
1	A	271	LEU
1	A	289	LEU
1	A	293	LYS
1	A	320	GLU
1	A	323	ARG
1	A	326	LEU
1	A	340	LEU
1	A	341	LEU
1	A	346	THR
1	A	351	THR
1	A	362	ARG
1	A	374	LYS
1	A	386	GLN
1	A	395	LYS
1	A	400	ASN
1	A	424	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	16	ASN
1	A	32	ASN
1	A	102	ASN
1	A	145	ASN
1	A	151	HIS
1	A	180	HIS
1	A	242	GLN
1	A	434	HIS
1	A	445	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	507	-	4,4,4	0.44	0	6,6,6	0.28	0
2	SO4	A	508	-	4,4,4	0.60	0	6,6,6	0.39	0
3	2AM	A	500	-	25,25,25	1.17	1 (4%)	38,38,38	2.33	8 (21%)
2	SO4	A	505	-	4,4,4	0.63	0	6,6,6	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	2AM	A	500	-	-	2/11/27/27	0/3/3/3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	500	2AM	C5-N7	-3.65	1.32	1.39

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	500	2AM	C5-C4-N3	-7.47	116.43	126.72
3	A	500	2AM	N3-C4-N9	6.55	138.31	127.17
3	A	500	2AM	C6-C5-C4	5.26	124.37	117.18
3	A	500	2AM	C6-C5-N7	-4.42	123.57	132.09
3	A	500	2AM	C2-N3-C4	2.90	118.92	111.83
3	A	500	2AM	O4'-C1'-N9	2.57	113.03	108.09
3	A	500	2AM	N3-C2-N1	-2.52	124.76	128.58
3	A	500	2AM	O3P-P-O2P	2.02	115.37	107.80

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	500	2AM	O4'-C4'-C5'-O5'
3	A	500	2AM	C2'-O2'-P-O1P

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	473/482 (98%)	-0.29	0 100 100	20, 29, 39, 55	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	A	507	5/5	0.83	0.23	52,52,53,53	0
3	2AM	A	500	23/23	0.92	0.09	41,45,49,52	0
2	SO4	A	508	5/5	0.93	0.11	51,51,52,52	0
2	SO4	A	505	5/5	0.96	0.09	29,29,31,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 2AM A 500:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.